


A solution to the Arf-Kervaire invariant problem I

Topologie Meeting
Oberwolfach

20 September, 2010



Mike Hill
University of Virginia
Mike Hopkins
Harvard University
Doug Ravenel
University of Rochester

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel




Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence



Vic Snaith and Bill Browder in 1981
Photo by Clarence Wilkerson

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire
formulation
- Questions raised by our
theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

A wildly popular dance craze



A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

A wildly popular dance craze



Drawing by Carolyn Snaith 1981
London, Ontario

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Mike Hill
Mike Hopkins
Doug Ravenel




Background and history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire
formulation
- Questions raised by our
theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence



Mike Hill, myself and Mike Hopkins
Photo taken by Bill Browder
February 11, 2010

Our main result

Our main theorem can be stated in three different but equivalent ways:

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.
- **Stable homotopy theoretic formulation:** It says that certain long sought hypothetical maps between high dimensional spheres do not exist.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.
- **Stable homotopy theoretic formulation:** It says that certain long sought hypothetical maps between high dimensional spheres do not exist.
- **Unstable homotopy theoretic formulation:** It says something about the EHP sequence, which has to do with unstable homotopy groups of spheres.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.
- **Stable homotopy theoretic formulation:** It says that certain long sought hypothetical maps between high dimensional spheres do not exist.
- **Unstable homotopy theoretic formulation:** It says something about the EHP sequence, which has to do with unstable homotopy groups of spheres.

The problem solved by our theorem is nearly 50 years old.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.
- **Stable homotopy theoretic formulation:** It says that certain long sought hypothetical maps between high dimensional spheres do not exist.
- **Unstable homotopy theoretic formulation:** It says something about the EHP sequence, which has to do with unstable homotopy groups of spheres.

The problem solved by our theorem is nearly 50 years old. There were several unsuccessful attempts to solve it in the 1970s.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result

Our main theorem can be stated in three different but equivalent ways:

- **Manifold formulation:** It says that a certain geometrically defined invariant $\Phi(M)$ (the Arf-Kervaire invariant, to be defined later) on certain manifolds M is always zero.
- **Stable homotopy theoretic formulation:** It says that certain long sought hypothetical maps between high dimensional spheres do not exist.
- **Unstable homotopy theoretic formulation:** It says something about the EHP sequence, which has to do with unstable homotopy groups of spheres.

The problem solved by our theorem is nearly 50 years old. There were several unsuccessful attempts to solve it in the 1970s. They were all aimed at proving the **opposite** of what we have proved.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence



Stable Homotopy Around the Arf-Kervaire Invariant, published in early 2009,

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work
The Arf-Kervaire formulation
Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence



Stable Homotopy Around the Arf-Kervaire Invariant, published in early 2009, **just before we proved our theorem.**

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work
The Arf-Kervaire formulation
Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence



Stable Homotopy Around the Arf-Kervaire Invariant, published in early 2009, **just before we proved our theorem.**

“As ideas for progress on a particular mathematics problem atrophy it can disappear.

Mike Hill
Mike Hopkins
Doug Ravenel



Our main result

Pontryagin's early work
The Arf-Kervaire formulation
Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence



Stable Homotopy Around the Arf-Kervaire Invariant, published in early 2009, **just before we proved our theorem.**

“As ideas for progress on a particular mathematics problem atrophy it can disappear. Accordingly I wrote this book to stem the tide of oblivion.”

Mike Hill
Mike Hopkins
Doug Ravenel



Our main result

- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Snaith's book (continued)



“For a brief period overnight we were convinced that we had the method to make all the sought after framed manifolds

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work
The Arf-Kervaire formulation
Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“For a brief period overnight we were convinced that we had the method to make all the sought after framed manifolds- a feeling which must have been shared by many topologists working on this problem.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“For a brief period overnight we were convinced that we had the method to make all the sought after framed manifolds- a feeling which must have been shared by many topologists working on this problem. All in all, the temporary high of believing that one had the construction

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“For a brief period overnight we were convinced that we had the method to make all the sought after framed manifolds- a feeling which must have been shared by many topologists working on this problem. All in all, the temporary high of believing that one had the construction was sufficient to maintain in me at least an enthusiastic spectator's interest in the problem.”

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“In the light of the above conjecture and the failure over fifty years to construct framed manifolds of Arf-Kervaire invariant one

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“In the light of the above conjecture and the failure over fifty years to construct framed manifolds of Arf-Kervaire invariant one this might turn out to be a book about things which do not exist.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“In the light of the above conjecture and the failure over fifty years to construct framed manifolds of Arf-Kervaire invariant one this might turn out to be a book about things which do not exist. This [is] why the quotations which preface each chapter contain a preponderance

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Snaith's book (continued)



“In the light of the above conjecture and the failure over fifty years to construct framed manifolds of Arf-Kervaire invariant one this might turn out to be a book about things which do not exist. This [is] why the quotations which preface each chapter contain a preponderance of utterances from the pen of Lewis Carroll.”

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)

Here is the stable homotopy theoretic formulation.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)

Here is the stable homotopy theoretic formulation.

Main Theorem

The Arf-Kervaire elements $\theta_j \in \pi_{2^{j+1}-2+n}(S^n)$ for large n do not exist for $j \geq 7$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)

Here is the stable homotopy theoretic formulation.

Main Theorem

The Arf-Kervaire elements $\theta_j \in \pi_{2^{j+1}-2+n}(S^n)$ for large n do not exist for $j \geq 7$.

The θ_j in the theorem is the name given to a hypothetical map between spheres for which the Arf-Kervaire invariant is nontrivial.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Here is the stable homotopy theoretic formulation.

Main Theorem

The Arf-Kervaire elements $\theta_j \in \pi_{2^{j+1}-2+n}(S^n)$ for large n do not exist for $j \geq 7$.

Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The θ_j in the theorem is the name given to a hypothetical map between spheres for which the Arf-Kervaire invariant is nontrivial. It follows from Browder's theorem of 1969 that such things can exist only in dimensions that are 2 less than a power of 2.

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j . They derived numerous consequences about homotopy groups of spheres.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j . They derived numerous consequences about homotopy groups of spheres. The possible nonexistence of the θ_j for large j was known as the **Doomsday Hypothesis**.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j . They derived numerous consequences about homotopy groups of spheres. The possible nonexistence of the θ_j for large j was known as the **Doomsday Hypothesis**.

After 1980, the problem faded into the background because it was thought to be too hard.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j . They derived numerous consequences about homotopy groups of spheres. The possible nonexistence of the θ_j for large j was known as the **Doomsday Hypothesis**.

After 1980, the problem faded into the background because it was thought to be too hard. Our proof is two giant steps away from anything that was attempted in the 70s.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Our main result (continued)



Mark Mahowald

Some homotopy theorists, most notably Mahowald, speculated about what would happen if θ_j existed for all j . They derived numerous consequences about homotopy groups of spheres. The possible nonexistence of the θ_j for large j was known as the **Doomsday Hypothesis**.

After 1980, the problem faded into the background because it was thought to be too hard. Our proof is two giant steps away from anything that was attempted in the 70s. We now know that the world of homotopy theory is very different from what they had envisioned then.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Mark Mahowald's sailboat

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history


Our main result

- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Mark Mahowald's sailboat



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

- Assume f is smooth.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

- Assume f is smooth. We know that any such map is can be continuously deformed to a smooth one.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

- Assume f is smooth. We know that any such map is can be continuously deformed to a smooth one.
- Pick a regular value $y \in S^n$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

- Assume f is smooth. We know that any such map can be continuously deformed to a smooth one.
- Pick a regular value $y \in S^n$. Its inverse image will be a smooth k -manifold M in S^{n+k} .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work on homotopy groups of spheres



Lev Pontryagin 1908-1988

Pontryagin's approach to maps $f : S^{n+k} \rightarrow S^n$ was

- Assume f is smooth. We know that any such map can be continuously deformed to a smooth one.
- Pick a regular value $y \in S^n$. Its inverse image will be a smooth k -manifold M in S^{n+k} .
- By studying such manifolds, Pontryagin was able to deduce things about maps between spheres.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Let D^n be the closure of an open ball around a regular value $y \in S^n$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Let D^n be the closure of an open ball around a regular value $y \in S^n$. If it is sufficiently small, then $V^{n+k} = f^{-1}(D^n) \subset S^{n+k}$ is an $(n+k)$ -manifold homeomorphic to $M \times D^n$ with boundary homeomorphic to $M \times S^{n-1}$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Let D^n be the closure of an open ball around a regular value $y \in S^n$. If it is sufficiently small, then $V^{n+k} = f^{-1}(D^n) \subset S^{n+k}$ is an $(n+k)$ -manifold homeomorphic to $M \times D^n$ with boundary homeomorphic to $M \times S^{n-1}$.

A local coordinate system around around the point $y \in S^n$ pulls back to one around M called a [framing](#).

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Let D^n be the closure of an open ball around a regular value $y \in S^n$. If it is sufficiently small, then $V^{n+k} = f^{-1}(D^n) \subset S^{n+k}$ is an $(n+k)$ -manifold homeomorphic to $M \times D^n$ with boundary homeomorphic to $M \times S^{n-1}$.

A local coordinate system around around the point $y \in S^n$ pulls back to one around M called a **framing**.

There is a way to reverse this procedure.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Let D^n be the closure of an open ball around a regular value $y \in S^n$. If it is sufficiently small, then $V^{n+k} = f^{-1}(D^n) \subset S^{n+k}$ is an $(n+k)$ -manifold homeomorphic to $M \times D^n$ with boundary homeomorphic to $M \times S^{n-1}$.

A local coordinate system around around the point $y \in S^n$ pulls back to one around M called a **framing**.

There is a way to reverse this procedure. A framed manifold $M^k \subset S^{n+k}$ determines a map $f : S^{n+k} \rightarrow S^n$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

To proceed further, we need to be more precise about what we mean by continuous deformation.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

$$h(x, 0) = f_1(x) \quad \text{and} \quad h(x, 1) = f_2(x).$$

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

$$h(x, 0) = f_1(x) \quad \text{and} \quad h(x, 1) = f_2(x).$$

If $y \in S^n$ is a regular value of h , then $h^{-1}(y)$ is a framed $(k+1)$ -manifold $N \subset S^{n+k} \times [0, 1]$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

$$h(x, 0) = f_1(x) \quad \text{and} \quad h(x, 1) = f_2(x).$$

If $y \in S^n$ is a regular value of h , then $h^{-1}(y)$ is a framed $(k+1)$ -manifold $N \subset S^{n+k} \times [0, 1]$ whose boundary is the disjoint union of $M_1 = f_1^{-1}(y)$ and $M_2 = f_2^{-1}(y)$.

Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

$$h(x, 0) = f_1(x) \quad \text{and} \quad h(x, 1) = f_2(x).$$

If $y \in S^n$ is a regular value of h , then $h^{-1}(y)$ is a framed $(k+1)$ -manifold $N \subset S^{n+k} \times [0, 1]$ whose boundary is the disjoint union of $M_1 = f_1^{-1}(y)$ and $M_2 = f_2^{-1}(y)$. This N is called a **framed cobordism** between M_1 and M_2 .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



To proceed further, we need to be more precise about what we mean by continuous deformation.

Two maps $f_1, f_2 : S^{n+k} \rightarrow S^n$ are **homotopic** if there is a continuous map $h : S^{n+k} \times [0, 1] \rightarrow S^n$ (called a **homotopy between f_1 and f_2**) such that

$$h(x, 0) = f_1(x) \quad \text{and} \quad h(x, 1) = f_2(x).$$

If $y \in S^n$ is a regular value of h , then $h^{-1}(y)$ is a framed $(k+1)$ -manifold $N \subset S^{n+k} \times [0, 1]$ whose boundary is the disjoint union of $M_1 = f_1^{-1}(y)$ and $M_2 = f_2^{-1}(y)$. This N is called a **framed cobordism** between M_1 and M_2 . When it exists the two closed manifolds are said to be **framed cobordant**.

Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Here is an example of a framed cobordism for $n = k = 1$.



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)



$$\Omega_k := \{\text{stably framed } k\text{-manifolds}\} / \text{cobordism}$$

Theorem: The above construction gives a bijection

$$\pi_{n+k}(S^n) \approx \Omega_k$$

where

$$\pi_{n+k}(S^n) := \{\text{maps } S^{n+k} \rightarrow S^n\} / \text{homotopy}$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$



$$\pi_n(S^n) = \mathbb{Z}$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=0$



$$\pi_n(S^n) = \mathbb{Z}$$

(the degree)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)


$k=0$



$$\pi_n(S^n) = \mathbb{Z}$$

(the degree)

$k=1$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)


$k=0$



$$\pi_n(S^n) = \mathbb{Z}$$

(the degree)

$k=1$



$$\pi_{n+1}(S^n) = \mathbb{Z}/2$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$ genus $M = 0 \Rightarrow M$ is a boundary

(since S^2 bounds a disk and
 $\pi_2(\mathrm{GL}_n(\mathbf{R}))=0$)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$ genus $M = 0 \Rightarrow M$ is a boundary

(since S^2 bounds a disk and
 $\pi_2(GL_n(\mathbf{R}))=0$)

Suppose the genus of M is
greater than 0.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$



choose an
embedded arc

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$



choose an
embedded arc

cut the surface open
and glue in disks

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

$k=2$



framed surgery

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

Obstruction: $\varphi : H_1(M; \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

Obstruction: $\varphi : H_1(M; \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$

Argument: Since the dimension of $H_1(M; \mathbb{Z}/2)$ is even, there is always a non-zero element in the kernel of φ , and so surgery can be performed.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's early work (continued)

Pontryagin (1930's)

Obstruction: $\varphi : H_1(M; \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$

Argument: Since the dimension of $H_1(M; \mathbb{Z}/2)$ is even, there is always a non-zero element in the kernel of φ , and so surgery can be performed.

Conclusion: $\Omega_2 = \pi_{n+2}(S^n) = 0$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's mistake for $k = 2$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's mistake for $k = 2$

The map $\varphi : H_1(M; \mathbf{Z}/2) \rightarrow \mathbf{Z}/2$ is **not** a homomorphism!

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

Pontryagin's mistake for $k = 2$

The map $\varphi : H_1(M; \mathbf{Z}/2) \rightarrow \mathbf{Z}/2$ is **not** a homomorphism!



Tuesday, April 21, 2009

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2

Let λ be a nonsingular anti-symmetric bilinear form on a free abelian group H of rank $2n$ with mod 2 reduction \overline{H} .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2

Let λ be a nonsingular anti-symmetric bilinear form on a free abelian group H of rank $2n$ with mod 2 reduction \overline{H} . It is known that \overline{H} has a basis of the form $\{a_i, b_i : 1 \leq i \leq n\}$ with

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2

Let λ be a nonsingular anti-symmetric bilinear form on a free abelian group H of rank $2n$ with mod 2 reduction \overline{H} . It is known that \overline{H} has a basis of the form $\{a_i, b_i: 1 \leq i \leq n\}$ with

$$\lambda(a_i, a_{i'}) = 0 \quad \lambda(b_j, b_{j'}) = 0 \quad \text{and} \quad \lambda(a_i, b_j) = \delta_{i,j}.$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2

Let λ be a nonsingular anti-symmetric bilinear form on a free abelian group H of rank $2n$ with mod 2 reduction \overline{H} . It is known that \overline{H} has a basis of the form $\{a_i, b_i: 1 \leq i \leq n\}$ with

$$\lambda(a_i, a_{i'}) = 0 \quad \lambda(b_j, b_{j'}) = 0 \quad \text{and} \quad \lambda(a_i, b_j) = \delta_{i,j}.$$

In other words, \overline{H} has a basis for which the bilinear form's matrix has the symplectic form

$$\begin{bmatrix} 0 & 1 & & & & & & & \\ 1 & 0 & & & & & & & \\ & & 0 & 1 & & & & & \\ & & 1 & 0 & & & & & \\ & & & & \ddots & & & & \\ & & & & & & 0 & 1 & \\ & & & & & & 1 & 0 & \end{bmatrix}.$$

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result
Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2 (continued)

A quadratic refinement of λ is a map $q : \bar{H} \rightarrow \mathbf{Z}/2$ satisfying

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2 (continued)

A quadratic refinement of λ is a map $q : \bar{H} \rightarrow \mathbf{Z}/2$ satisfying

$$q(x + y) = q(x) + q(y) + \lambda(x, y)$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2 (continued)

A quadratic refinement of λ is a map $q : \bar{H} \rightarrow \mathbf{Z}/2$ satisfying

$$q(x + y) = q(x) + q(y) + \lambda(x, y)$$

Its Arf invariant is

$$\text{Arf}(q) = \sum_{i=1}^n q(a_i)q(b_i) \in \mathbf{Z}/2.$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Arf invariant of a quadratic form in characteristic 2 (continued)

A quadratic refinement of λ is a map $q : \bar{H} \rightarrow \mathbf{Z}/2$ satisfying

$$q(x + y) = q(x) + q(y) + \lambda(x, y)$$

Its Arf invariant is

$$\text{Arf}(q) = \sum_{i=1}^n q(a_i)q(b_i) \in \mathbf{Z}/2.$$

In 1941 Arf proved that this invariant (along with the number n) determines the isomorphism type of q .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

From my stamp collection



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

From my stamp collection



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy


Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

From my stamp collection



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Money talks: Arf's definition republished in 2009



Cahit Arf 1910-1997

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension. Each $x \in H$ is represented by an immersion $i_x : S^{2m+1} \looparrowright M$ with a stably trivialized normal bundle.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension. Each $x \in H$ is represented by an immersion $i_x : S^{2m+1} \looparrowright M$ with a stably trivialized normal bundle. H has an antisymmetric bilinear form λ defined in terms of intersection numbers.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension. Each $x \in H$ is represented by an immersion $i_x : S^{2m+1} \looparrowright M$ with a stably trivialized normal bundle. H has an antisymmetric bilinear form λ defined in terms of intersection numbers.



Michel Kervaire 1927-2007

Kervaire defined a quadratic refinement q on its mod 2 reduction in terms of the trivialization of each sphere's normal bundle.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension. Each $x \in H$ is represented by an immersion $i_x : S^{2m+1} \looparrowright M$ with a stably trivialized normal bundle. H has an antisymmetric bilinear form λ defined in terms of intersection numbers.



Michel Kervaire 1927-2007

Kervaire defined a quadratic refinement q on its mod 2 reduction in terms of the trivialization of each sphere's normal bundle. The **Kervaire invariant** $\Phi(M)$ is defined to be the Arf invariant of q .

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold

Let M be a $2m$ -connected smooth closed framed manifold of dimension $4m + 2$. Let $H = H_{2m+1}(M; \mathbf{Z})$, the homology group in the middle dimension. Each $x \in H$ is represented by an immersion $i_x : S^{2m+1} \looparrowright M$ with a stably trivialized normal bundle. H has an antisymmetric bilinear form λ defined in terms of intersection numbers.



Michel Kervaire 1927-2007

Kervaire defined a quadratic refinement q on its mod 2 reduction in terms of the trivialization of each sphere's normal bundle. The **Kervaire invariant** $\Phi(M)$ is defined to be the Arf invariant of q .

For $m = 0$, Kervaire's q coincides with Pontryagin's φ .

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

What can we say about $\Phi(M)$?

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

What can we say about $\Phi(M)$?

- For $m = 0$ there is a framing on the torus $S^1 \times S^1 \subset \mathbf{R}^4$ with nontrivial Kervaire invariant.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m+2)$ -manifold (continued)

What can we say about $\Phi(M)$?

- For $m = 0$ there is a framing on the torus $S^1 \times S^1 \subset \mathbf{R}^4$ with nontrivial Kervaire invariant. Pontryagin used it in 1950 (after some false starts in the 30s) to show $\pi_{n+2}(S^n) = \mathbf{Z}/2$ for all $n \geq 2$.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

What can we say about $\Phi(M)$?

- For $m = 0$ there is a framing on the torus $S^1 \times S^1 \subset \mathbf{R}^4$ with nontrivial Kervaire invariant. Pontryagin used it in 1950 (after some false starts in the 30s) to show $\pi_{n+2}(S^n) = \mathbf{Z}/2$ for all $n \geq 2$.



A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- Kervaire (1960) showed it must vanish when $m = 2$.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- Kervaire (1960) showed it must vanish when $m = 2$. This enabled him to construct the first example of a topological manifold (of dimension 10) without a smooth structure.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω


How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- Kervaire (1960) showed it must vanish when $m = 2$. This enabled him to construct the first example of a topological manifold (of dimension 10) without a smooth structure.



A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.



Ed Brown



Frank Peterson
1930-2000

Brown-Peterson (1966) showed that it vanishes for all positive even m .

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.



Bill Browder

Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j .

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.



Bill Browder

Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result
Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

-



Bill Browder

Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result
Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.



Bill Browder

Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem. **This is the stable homotopy theoretic formulation of the problem.**

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result
Pontryagin's early work

The Arf-Kervaire formulation


Questions raised by our theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- 
Bill Browder
Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem. **This is the stable homotopy theoretic formulation of the problem.**
- θ_j is known to exist for $1 \leq j \leq 5$, i.e., in dimensions 2, 6, 14, 30 and 62.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- 
Bill Browder
Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem. **This is the stable homotopy theoretic formulation of the problem.**
- θ_j is known to exist for $1 \leq j \leq 5$, i.e., in dimensions 2, 6, 14, 30 and 62.
- In the decade following Browder's theorem, many topologists tried **without success** to construct framed manifolds with nontrivial Kervaire invariant in **all** dimensions 2 less than a power of 2.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof


The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.

- 
Bill Browder
Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem. **This is the stable homotopy theoretic formulation of the problem.**
- θ_j is known to exist for $1 \leq j \leq 5$, i.e., in dimensions 2, 6, 14, 30 and 62.
- In the decade following Browder's theorem, many topologists tried **without success** to construct framed manifolds with nontrivial Kervaire invariant in **all** dimensions 2 less than a power of 2.
- Our theorem says θ_j does **not** exist for $j \geq 6$.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The Kervaire invariant of a framed $(4m + 2)$ -manifold (continued)

More of what we can say about $\Phi(M)$.



Bill Browder

- Browder (1969) showed that it can be nontrivial only if $m = 2^{j-1} - 1$ for some positive integer j . This happens iff the element h_j^2 is a permanent cycle in the Adams spectral sequence. The corresponding element in $\pi_{n+2^{j+1}-2}(S^n)$ for large n is θ_j , the subject of our theorem. **This is the stable homotopy theoretic formulation of the problem.**
- θ_j is known to exist for $1 \leq j \leq 5$, i.e., in dimensions 2, 6, 14, 30 and 62.
- In the decade following Browder's theorem, many topologists tried **without success** to construct framed manifolds with nontrivial Kervaire invariant in **all** dimensions 2 less than a power of 2.
- Our theorem says θ_j does **not** exist for $j \geq 7$. The case $j = 6$ is still open.

A solution to the Arf-Kervaire invariant problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

Our main result

Pontryagin's early work

The Arf-Kervaire formulation

Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Questions raised by our theorem

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. We have no idea what their targets are.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. We have no idea what their targets are.

Unstable homotopy theoretic formulation.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres. Since they do not exist, a substitute for his conjecture is needed.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres. Since they do not exist, a substitute for his conjecture is needed. **We have no idea what it should be.**

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres. Since they do not exist, a substitute for his conjecture is needed. **We have no idea what it should be.**

Our method of proof offers a new tool, **the slice spectral sequence**, for studying the stable homotopy groups of spheres.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres. Since they do not exist, a substitute for his conjecture is needed. **We have no idea what it should be.**

Our method of proof offers a new tool, **the slice spectral sequence**, for studying the stable homotopy groups of spheres. We look forward to learning more with it in the future.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Questions raised by our theorem

Adams spectral sequence formulation. We now know that the h_j^2 for $j \geq 7$ are not permanent cycles, so they have to support nontrivial differentials. **We have no idea what their targets are.**

Unstable homotopy theoretic formulation. In 1967 Mahowald published an elaborate conjecture about the role of the θ_j (assuming that they all exist) in the unstable homotopy groups of spheres. Since they do not exist, a substitute for his conjecture is needed. **We have no idea what it should be.**

Our method of proof offers a new tool, **the slice spectral sequence**, for studying the stable homotopy groups of spheres. We look forward to learning more with it in the future. **We will illustrate it at the end of the talk.**

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire
formulation
- Questions raised by our
theorem

Our strategy

Ingredients of the proof

- The spectrum Ω
- How we construct Ω
- The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.
- Fiber sequences and cofiber sequences are the same, up to weak equivalence.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.
- Fiber sequences and cofiber sequences are the same, up to weak equivalence.
- While space X has a homotopy group $\pi_k(X)$ for each positive integer k ,



Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.
- Fiber sequences and cofiber sequences are the same, up to weak equivalence.
- While space X has a homotopy group $\pi_k(X)$ for each positive integer k , a spectrum X has an abelian homotopy group $\pi_k(X)$ **defined for every integer k** .



Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.
- Fiber sequences and cofiber sequences are the same, up to weak equivalence.
- While space X has a homotopy group $\pi_k(X)$ for each positive integer k , a spectrum X has an abelian homotopy group $\pi_k(X)$ **defined for every integer k** .

For the sphere spectrum S^0 , $\pi_k(S^0)$ is the usual homotopy group $\pi_{n+k}(S^n)$ for $n > k + 1$.



Ingredients of the proof

Our proof has several ingredients.

- We use methods of **stable homotopy theory**, which means we use spectra instead of topological spaces. Roughly speaking, spectra are to spaces as integers are to natural numbers. Instead of making addition formally invertible, we do the same for suspension.

This means

- Every spectrum X is equivalent to the suspension of another spectrum $Y = \Sigma^{-1}X$.
- X is equivalent to $\Omega\Sigma X$.
- Fiber sequences and cofiber sequences are the same, up to weak equivalence.
- While space X has a homotopy group $\pi_k(X)$ for each positive integer k , a spectrum X has an abelian homotopy group $\pi_k(X)$ **defined for every integer k** .

For the sphere spectrum S^0 , $\pi_k(S^0)$ is the usual homotopy group $\pi_{n+k}(S^n)$ for $n > k + 1$. The hypothetical θ_j is an element of this group for $k = 2^{j+1} - 2$.



Ingredients of the proof (continued)

More ingredients of our proof:

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We use [complex cobordism theory](#).

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We use **complex cobordism theory**. This is a branch of algebraic topology having deep connections with algebraic geometry and number theory.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We use **complex cobordism theory**. This is a branch of algebraic topology having deep connections with algebraic geometry and number theory. It includes some highly developed computational techniques that began with work by Milnor, Novikov and Quillen in the 60s.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We use **complex cobordism theory**. This is a branch of algebraic topology having deep connections with algebraic geometry and number theory. It includes some highly developed computational techniques that began with work by Milnor, Novikov and Quillen in the 60s. A pivotal tool in the subject is the theory of formal group laws.



John Milnor



Sergei Novikov



Dan Quillen

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We also make use of newer less familiar methods from **equivariant stable homotopy theory**.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We also make use of newer less familiar methods from **equivariant stable homotopy theory**. This means there is a finite group G (a cyclic 2-group) acting on all spaces in sight, and all maps are required to commute with these actions.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We also make use of newer less familiar methods from **equivariant stable homotopy theory**. This means there is a finite group G (a cyclic 2-group) acting on all spaces in sight, and all maps are required to commute with these actions. When we pass to spectra, we get homotopy groups indexed not just by the integers \mathbf{Z} , but by $RO(G)$, the real representation ring of G .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω
How we construct Ω
The slice spectral sequence

Ingredients of the proof (continued)

More ingredients of our proof:

- We also make use of newer less familiar methods from **equivariant stable homotopy theory**. This means there is a finite group G (a cyclic 2-group) acting on all spaces in sight, and all maps are required to commute with these actions. When we pass to spectra, we get homotopy groups indexed not just by the integers \mathbf{Z} , but by $RO(G)$, **the real representation ring of G** . Our calculations make use of this richer structure.



Peter May



John Greenlees



Gaunce Lewis
1949-2006

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω
The slice spectral sequence

The spectrum Ω

We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω
The slice spectral sequence

The spectrum Ω

We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence (which is a device for calculating homotopy groups) in which the image of each θ_j is nontrivial.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The spectrum Ω

We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence (which is a device for calculating homotopy groups) in which the image of each θ_j is nontrivial. This means that if θ_j exists, we will see its image in $\pi_*(\Omega)$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The spectrum Ω

We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence (which is a device for calculating homotopy groups) in which the image of each θ_j is nontrivial. This means that if θ_j exists, we will see its image in $\pi_*(\Omega)$.
- (ii) **Periodicity Theorem.** It is 256-periodic, meaning that $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The spectrum Ω

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence (which is a device for calculating homotopy groups) in which the image of each θ_j is nontrivial. This means that if θ_j exists, we will see its image in $\pi_*(\Omega)$.
- (ii) **Periodicity Theorem.** It is 256-periodic, meaning that $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
- (iii) **Gap Theorem.** $\pi_k(\Omega) = 0$ for $-4 < k < 0$.



We will produce a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective spectrum (meaning that it has nontrivial homotopy groups in arbitrarily large negative dimensions) with the following properties.

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence (which is a device for calculating homotopy groups) in which the image of each θ_j is nontrivial. This means that if θ_j exists, we will see its image in $\pi_*(\Omega)$.
- (ii) **Periodicity Theorem.** It is 256-periodic, meaning that $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
- (iii) **Gap Theorem.** $\pi_k(\Omega) = 0$ for $-4 < k < 0$. This property is our **zinger**. Its proof involves a new tool we call the slice spectral sequence.

The spectrum Ω (continued)

Here again are the properties of Ω

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω
The slice spectral sequence

The spectrum Ω (continued)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Here again are the properties of Ω

- (i) **Detection Theorem.** If θ_j exists, it has nontrivial image in $\pi_*(\Omega)$.
- (ii) **Periodicity Theorem.** $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
- (iii) **Gap Theorem.** $\pi_{-2}(\Omega) = 0$.

Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The spectrum Ω (continued)

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Here again are the properties of Ω

- (i) **Detection Theorem.** If θ_j exists, it has nontrivial image in $\pi_*(\Omega)$.
 - (ii) **Periodicity Theorem.** $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
 - (iii) **Gap Theorem.** $\pi_{-2}(\Omega) = 0$.
- (ii) and (iii) imply that $\pi_{254}(\Omega) = 0$.

Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

The spectrum Ω (continued)



Here again are the properties of Ω

- (i) **Detection Theorem.** If θ_j exists, it has nontrivial image in $\pi_*(\Omega)$.
 - (ii) **Periodicity Theorem.** $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
 - (iii) **Gap Theorem.** $\pi_{-2}(\Omega) = 0$.
- (ii) and (iii) imply that $\pi_{254}(\Omega) = 0$.

If $\theta_7 \in \pi_{254}(S^0)$ exists, (i) implies it has a nontrivial image in this group, so it cannot exist.

The spectrum Ω (continued)



Here again are the properties of Ω

- (i) **Detection Theorem.** If θ_j exists, it has nontrivial image in $\pi_*(\Omega)$.
 - (ii) **Periodicity Theorem.** $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
 - (iii) **Gap Theorem.** $\pi_{-2}(\Omega) = 0$.
- (ii) and (iii) imply that $\pi_{254}(\Omega) = 0$.

If $\theta_7 \in \pi_{254}(S^0)$ exists, (i) implies it has a nontrivial image in this group, so it cannot exist. The argument for θ_j for larger j is similar, since $|\theta_j| = 2^{j+1} - 2 \equiv -2 \pmod{256}$ for $j \geq 7$.

Background and history

Our main result
Pontryagin's early work
The Arf-Kervaire formulation
Questions raised by our theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω
The slice spectral sequence

How we construct Ω

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

To construct it we start with the complex cobordism spectrum MU .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

To construct it we start with the complex cobordism spectrum MU . It can be thought of as the set of complex points of an algebraic variety defined over the real numbers.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

To construct it we start with the complex cobordism spectrum MU . It can be thought of as the set of complex points of an algebraic variety defined over the real numbers. This means that it has an action of C_2 defined by complex conjugation.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

To construct it we start with the complex cobordism spectrum MU . It can be thought of as the set of complex points of an algebraic variety defined over the real numbers. This means that it has an action of C_2 defined by complex conjugation. The fixed point set of this action is the set of real points, known to topologists as MO , the unoriented cobordism spectrum.

Mike Hill
Mike Hopkins
Doug Ravenel



Background and history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω

How we construct Ω

- The slice spectral sequence

Our spectrum Ω will be the fixed point spectrum for the action of C_8 (the cyclic group of order 8) on an equivariant spectrum $\tilde{\Omega}$.

To construct it we start with the complex cobordism spectrum MU . It can be thought of as the set of complex points of an algebraic variety defined over the real numbers. This means that it has an action of C_2 defined by complex conjugation. The fixed point set of this action is the set of real points, known to topologists as MO , the unoriented cobordism spectrum. In this notation, U and O stand for the unitary and orthogonal groups.

Mike Hill
Mike Hopkins
Doug Ravenel



Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Ingredients of the proof
The spectrum Ω

The slice spectral sequence

How we construct Ω (continued)

Some people who have studied MU as a C_2 -spectrum:

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result

Pontryagin's early work

The Arf-Kervaire
formulation

Questions raised by our
theorem

Our strategy

Ingredients of the proof

The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

Some people who have studied MU as a C_2 -spectrum:



Peter Landweber

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

Some people who have studied MU as a C_2 -spectrum:



Peter Landweber



Shoro Araki
1930–2005

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

Some people who have studied MU as a C_2 -spectrum:



Peter Landweber



Igor Kriz and Po Hu



Shoro Araki
1930–2005

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

Some people who have studied MU as a C_2 -spectrum:



Peter Landweber



Igor Kriz and Po Hu



Shoro Araki
1930–2005



Nitu Kitchloo



Steve Wilson

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X . Here the action of H on G is by right multiplication, and the resulting object has an action of G by left multiplication.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X . Here the action of H on G is by right multiplication, and the resulting object has an action of G by left multiplication. As a set, $Y = X^{|G/H|}$, the $|G/H|$ -fold Cartesian power of X .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X . Here the action of H on G is by right multiplication, and the resulting object has an action of G by left multiplication. As a set, $Y = X^{|G/H|}$, the $|G/H|$ -fold Cartesian power of X . A general element of G permutes these factors, each of which is left invariant by the subgroup H .

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X . Here the action of H on G is by right multiplication, and the resulting object has an action of G by left multiplication. As a set, $Y = X^{|G/H|}$, the $|G/H|$ -fold Cartesian power of X . A general element of G permutes these factors, each of which is left invariant by the subgroup H .

In particular we get a C_8 -spectrum

$$MU^{(4)} = \text{Map}_{C_2}(C_8, MU).$$

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

How we construct Ω (continued)

To get a C_8 -spectrum, we use the following general construction for getting from a space or spectrum X acted on by a group H to one acted on by a larger group G containing H as a subgroup. Let

$$Y = \text{Map}_H(G, X),$$

the space (or spectrum) of H -equivariant maps from G to X . Here the action of H on G is by right multiplication, and the resulting object has an action of G by left multiplication. As a set, $Y = X^{|G/H|}$, the $|G/H|$ -fold Cartesian power of X . A general element of G permutes these factors, each of which is left invariant by the subgroup H .

In particular we get a C_8 -spectrum

$$MU^{(4)} = \text{Map}_{C_2}(C_8, MU).$$

This spectrum is not periodic, but it has a close relative $\tilde{\Omega}$ which is.

A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem


Our strategy

Ingredients of the proof
The spectrum Ω

How we construct Ω

The slice spectral sequence

A homotopy fixed point spectral sequence



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history


Our main result
Pontryagin's early work
The Arf-Kervaire
formulation
Questions raised by our
theorem

Our strategy

Ingredients of the proof
The spectrum Ω
How we construct Ω

The slice spectral sequence

The corresponding slice spectral sequence



A solution to the
Arf-Kervaire invariant
problem

Mike Hill
Mike Hopkins
Doug Ravenel



Background and
history

- Our main result
- Pontryagin's early work
- The Arf-Kervaire formulation
- Questions raised by our theorem

Our strategy

- Ingredients of the proof
- The spectrum Ω
- How we construct Ω

The slice spectral sequence