

A wildly popular dance craze




Drawing by Carolyn Snaith 1981
London, Ontario

1.1

Overview of the proof

MSRI Hot Topics Workshop
on the Kervaire Invariant

October 25, 2010



Mike Hill
University of Virginia
Mike Hopkins
Harvard University
Doug Ravenel
University of Rochester

1.2

1 Our strategy

Our strategy

Main Theorem. *The Arf-Kervaire elements $\theta_j \in \pi_{2j+1-2}(S^0)$ do not exist for $j \geq 7$.*

We will prove it by producing a map $S^0 \rightarrow \Omega$, where Ω is a nonconnective E_∞ -ring spectrum with the following properties.

1.3

- (i) **Detection Theorem.** It has an Adams-Novikov spectral sequence in which the image of each θ_j is nontrivial. This means that if θ_j exists, we will see its image in $\pi_*(\Omega)$. This will be the subject of a talk by myself on Thursday.
- (ii) **Periodicity Theorem.** It is 256-periodic, meaning that $\pi_k(\Omega)$ depends only on the reduction of k modulo 256. This will be the subject of a talk by Hu on Thursday.
- (iii) **Gap Theorem.** $\pi_k(\Omega) = 0$ for $-4 < k < 0$. This property is our **zinger**. Its proof involves a new tool we call the slice spectral sequence. This will be the subject of a talk by Gerhardt on Tuesday, modulo some technical theorems to be proved later.

How the theorem follows from the existence of Ω

Here again are the properties of Ω :

- (i) **Detection Theorem.** If θ_j exists, it has nontrivial image in $\pi_*(\Omega)$.
 - (ii) **Periodicity Theorem.** $\pi_k(\Omega)$ depends only on the reduction of k modulo 256.
 - (iii) **Gap Theorem.** $\pi_{-2}(\Omega) = 0$.
- (ii) and (iii) imply that $\pi_{254}(\Omega) = 0$.

If $\theta_7 \in \pi_{254}(S^0)$ exists, (i) implies it has a nontrivial image in this group, so it cannot exist. The argument for θ_j for larger j is similar, since $|\theta_j| = 2^{j+1} - 2 \equiv -2 \pmod{256}$ for $j \geq 7$.

1.4

2 The construction of Ω

How we construct the spectrum Ω

The construction of Ω requires the use of **equivariant stable homotopy theory**. **It will be the subject of talks by May and Greenlees today and tomorrow.**

Roughly speaking, an equivariant G -spectrum is a spectrum X with an action of the group G . **For us the group of interest will be C_8** . This leads to a fixed point spectrum X^G and a homotopy fixed point spectrum X^{hG} , with a map $X^G \rightarrow X^{hG}$.

For a G -space X , X^G is the subspace fixed by all of G , which is the same as the space of equivariant maps from a point to X , $\text{Map}_G(*, X)$. To get X^{hG} , we replace the point here by an free contractible G -space EG .

The homotopy type of $X^{hG} = \text{Map}_G(EG, X)$ is known to be independent of the choice of EG . The unique map $EG \rightarrow *$ leads to the map $X^G \rightarrow X^{hG}$.

1.5

How we construct the spectrum Ω (continued)

We construct a C_8 -spectrum $\tilde{\Omega}$ and show that

- $\tilde{\Omega}^{hC_8}$ satisfies the detection and periodicity theorems.
- $\tilde{\Omega}^{C_8}$ satisfies the gap theorem.

Hence our proof depends on a fourth property:

- (iv) **Fixed Point Theorem** The map $\tilde{\Omega}^{C_8} \rightarrow \tilde{\Omega}^{hC_8}$ is an equivalence. **This will be the subject of a Blumberg's talk on Friday.** This fixed point set is our spectrum Ω .

We will come back to the definition of $\tilde{\Omega}$ below.

1.6

3 MU and its equivariant relatives

MU and its equivariant relatives

The starting point for the construction of $\tilde{\Omega}$ is the action of C_2 on the complex cobordism spectrum MU given by complex conjugation. The resulting C_2 -equivariant spectrum is denoted by $MU_{\mathbf{R}}$ and is called **real cobordism theory**. This terminology follows Atiyah's definition of real K -theory, by which he meant complex K -theory equipped with complex conjugation.

Next we use a formal tool we call the **norm N_H^G** for inducing up from an H -spectrum to a G -spectrum when H is a subgroup of G . **This will be the subject of Miller's talk on Wednesday.**

1.7

[MU and its equivariant relatives \(continued\)](#)

For an H -space X , we have a G -space

$$\text{Map}_H(G, X),$$

where H acts on G by right multiplication and G acts on the mapping space via right multiplication in G . The underlying space here is the Cartesian product $X^{|G/H|}$. G permutes the factors the same way it permutes cosets, and each factor is invariant under H . [The norm functor is an analogous construction in the stable category.](#)

The case of interest to us is $X = MU_{\mathbf{R}}$, $H = C_2$ and $G = C_{2^{n+1}}$. This means that the underlying spectrum of $N_H^G X$ is $MU^{(2^n)}$, the 2^n -fold smash power of MU .

1.8

[MU and its equivariant relatives \(continued\)](#)

In order to proceed further we need to introduce [RO\(G\)-graded homotopy](#), where $RO(G)$ denotes the orthogonal representation ring of G . Let S^V denote the one point compactification of orthogonal representation V , and for a G -space or spectrum X define

$$\pi_V^G X = \text{Map}_G(S^V, X).$$

Note that when the action of G on V is trivial, an equivariant map $S^V = S^{\dim V} \rightarrow X$ must land in X^G , so

$$\pi_n^G X = \pi_n X^G.$$

In the stable category we can make sense of this for virtual as well as actual representations, so we get homotopy groups indexed by $RO(G)$, which we denote collectively by $\pi_*^G X$.

1.9

[MU and its equivariant relatives \(continued\)](#)

Recall that

$$\pi_*(MU) = \mathbf{Z}[x_1, x_2, \dots] \quad \text{with } |x_i| = 2i.$$

It turns out that any choice of generator $x_i : S^{2i} \rightarrow MU$ is the image of the forgetful functor of a map

$$S^{i\rho} \xrightarrow{\bar{x}_i} MU_{\mathbf{R}}.$$

Here ρ denotes the regular real representation of C_2 , which is the same thing as the complex numbers \mathbf{C} acted on by conjugation.

1.10

[MU and its equivariant relatives \(continued\)](#)

For $G = C_{2^{n+1}}$, the G -spectrum $N_{C_2}^G MU_{\mathbf{R}}$ is underlain by $MU^{(2^n)}$. $\pi_* MU^{(2^n)}$ is a graded polynomial algebra over \mathbf{Z} where

- there are 2^n generators in each positive even dimension $2i$.
- they are acted on transitively by G .

For a group generator $\gamma \in G$ and polynomial generator $r_i \in \pi_{2i}$, the set

$$\{\gamma^j r_i : 0 \leq j < 2^n\}$$

is algebraically independent, and $\gamma^{2^n} r_i = (-1)^i r_i$.

We define specific polynomial generators \bar{r}_i that are convenient for our purposes. [They are the subject of Kitchloo's talk on Wednesday.](#)

1.11

4 The slice filtration

The slice filtration

Now we introduce our main technical tool, [the slice filtration](#). It will be the subject of a more detailed talk by Hill tomorrow.

First we need to recall some things about the classical Postnikov tower. The m th Postnikov section $P^m X$ of a space or spectrum X is obtained by killing all homotopy groups of X above dimension m by attaching cells. The fiber of the map $X \rightarrow P^m X$ is $P_{m+1} X$, the m -connected cover of X .

These two functors have some universal properties. Let \mathcal{S} and $\mathcal{S}_{>m}$ denote the categories of spectra and m -connected spectra.

1.12

More about the Postnikov tower

The functor P^m is [Dror nullification](#) with respect to the subcategory $\mathcal{S}_{>m}$. This means the map $X \rightarrow P^m X$ is universal among maps from X to spectra which are $\mathcal{S}_{>m}$ -null in the sense that all maps to them from m -connected spectra are null. In other words,

- The spectrum $P^m X$ is $\mathcal{S}_{>m}$ -null.
- For any $\mathcal{S}_{>m}$ -null spectrum Z , the map $\mathcal{S}(P^m X, Z) \rightarrow \mathcal{S}(X, Z)$ is an equivalence.

Since $\mathcal{S}_{>m} \subset \mathcal{S}_{>m-1}$, there is a natural transformation $P^m \rightarrow P^{m-1}$, whose fiber is denoted by $P_m^m X$.

1.13

An equivariant Postnikov tower

In what follows G will be an arbitrary finite cyclic 2-group, and $g = |G|$. Let \mathcal{S}^G denote the category of G -equivariant spectra. We need an equivariant analog of $\mathcal{S}_{>m}$. Our choice for this is somewhat novel.

Recall that $\mathcal{S}_{>m}$ is the category of spectra built up out of spheres of dimension $> m$ using arbitrary wedges and mapping cones.

For a subgroup H of G with $|H| = h$ and an integer k , let

$$\widehat{S}(k\rho_H) = G_+ \wedge_H S^{k\rho_H}$$

where ρ_H denotes the regular real representation of H . Its underlying spectrum is a wedge of g/h spheres of dimension kh which are permuted by elements of G and are invariant under H .

1.14

An equivariant Postnikov tower (continued)

We will replace the set of sphere spectra by

$$\mathcal{A} = \left\{ \widehat{S}(k\rho_H), \Sigma^{-1}\widehat{S}(k\rho_H) : H \subset G, k \in \mathbf{Z} \right\}.$$

We will refer to the elements in this set as [slice cells](#) or simply as [cells](#). Note that $\Sigma^{-2}\widehat{S}(k\rho_H)$ (and larger desuspensions) are [not](#) cells. A [free cell](#) is one of the form $\widehat{S}(k\rho_{\{e\}})$, a wedge of g k -spheres permuted by G . Note that

$$\Sigma^{-1}\widehat{S}(k\rho_{\{e\}}) = \widehat{S}((k-1)\rho_{\{e\}}).$$

Nonfree cells are said to be [isotropic](#).

In order to define $\mathcal{S}_{>m}^G$, we need to assign a dimension to each element in \mathcal{A} . We do this in terms of the underlying wedge summands, namely

$$\dim \widehat{S}(k\rho_H) = kh \quad \text{and} \quad \dim \Sigma^{-1}\widehat{S}(k\rho_H) = kh - 1.$$

1.15

An equivariant Postnikov tower (continued)

Then $\mathcal{S}_{>m}^G$ is the category built up out of elements in \mathcal{A} of dimension $> m$ using arbitrary wedges, mapping cones and smash products with equivariant suspension spectra.

With this definition it is possible to construct functors P_{m+1}^G and P_m^m with the same formal properties as in the classical case. Thus we get a tower

$$\begin{array}{ccccccc} \dots & \longrightarrow & P_G^{m+1}X & \longrightarrow & P_m^m X & \longrightarrow & P_G^{m-1}X & \longrightarrow & \dots \\ & & \uparrow & & \uparrow & & \uparrow & & \\ & & {}^G P_{m+1}^{m+1}X & & {}^G P_m^m X & & {}^G P_{m-1}^{m-1}X & & \end{array}$$

in which the homotopy limit is X and the homotopy colimit is contractible.

1.16

5 The slice spectral sequence

The slice spectral sequence

$$\begin{array}{ccccccc} \dots & \longrightarrow & P_G^{m+1}X & \longrightarrow & P_m^m X & \longrightarrow & P_G^{m-1}X & \longrightarrow & \dots \\ & & \uparrow & & \uparrow & & \uparrow & & \\ & & {}^G P_{m+1}^{m+1}X & & {}^G P_m^m X & & {}^G P_{m-1}^{m-1}X & & \end{array}$$

We call this the **slice tower**. ${}^G P_m^m X$ is the n th **slice** and the decreasing sequence of subgroups of $\pi_*(X)$ is the **slice filtration**. We also get slice filtrations of the $RO(G)$ -graded homotopy $\pi_*^G(X)$ and the homotopy groups of fixed point sets $\pi_*(X^H)$.

There is an important difference between this tower and the classical one. In the classical case the map $X \rightarrow P^m X$ does not change homotopy groups in dimensions $\leq m$. **This is not true in this equivariant case.**

1.17

The slice spectral sequence (continued)

In the classical case, $P_m^m X$ is an Eilenberg-Mac Lane spectrum whose n th homotopy group is that of X . In our case, $\pi_*({}^G P_m^m X)$ need not be concentrated in dimension m .

This means the slice filtration leads to a (possibly noncollapsing) **slice spectral sequence** converging to $\pi_*(X)$ and its variants.

One variant has the form

$$E_2^{s,t} = \pi_{t-s}^G({}^G P_t^t X) \implies \pi_{t-s}^G(X).$$

Recall that $\pi_*^G(X)$ is by definition $\pi_*(X^G)$, the homotopy of the fixed point set.

This is the spectral sequence we will use to study $MU_{\mathbf{R}}^{(4)}$ and its relatives.

1.18

The slice spectral sequence (continued)

A large portion of our paper is devoted to proving that the slice spectral sequence has the desired properties. From now on we will drop the symbol G from the functors P^m , P_{m+1} and P_m^m .

Slice Theorem . *In the slice tower for $N_{C_2}^G MU_{\mathbf{R}}$, every odd slice is contractible and $P_{2m}^{2m} = \widehat{W}_m \wedge H\mathbf{Z}$, where \widehat{W}_m is a certain wedge of $2m$ -dimensional slice cells (to be named later) and $H\mathbf{Z}$ is the integer Eilenberg-Mac Lane spectrum. \widehat{W}_m never has any free summands.*

Our G -spectrum $\tilde{\Omega}$ (where $G = C_8$) is obtained from the E_∞ -ring spectrum $N_{C_2}^G MU_{\mathbf{R}}$ by inverting a certain element $D \in \pi_{19\rho_G}^G$. The choices of G and D are the simplest ones leading to a homotopy fixed point set with the detection property, as we will see in the talk on that topic on Thursday. The slice tower for $\tilde{\Omega}$ has similar properties to that of $N_{C_2}^G MU_{\mathbf{R}}$.

1.19

6 The gap theorem

Proving the gap theorem

The gap theorem follows from the fact that π_{-2}^G vanishes for each isotropic slice, i.e., for each one of the form

$$\widehat{S}(k\rho_H) \wedge H\mathbf{Z}$$


for nontrivial H . This will be explained by Gerhardt tomorrow.

In order to give a feel for these calculations we offer the following picture of $\pi_*^G S^{k\rho_G} \wedge H\mathbf{Z}$ for $G = C_8$ and various integers k .

1.20

Some C_8 slices

A picture of $\pi_*^G S^{k\rho_G} \wedge H\mathbf{Z}$ for $G = C_8$ and various integers k .



1.21

7 The periodicity theorem

Proving the Periodicity Theorem

We now outline the proof of the Periodicity Theorem, assuming the Slice Theorem. Details will be given in Hu's talk on Thursday, with some preparation in Kriz' talk tomorrow.

We establish some differentials in the slice spectral sequence and show that certain elements become permanent cycles after inverting a certain $D \in \pi_{19\rho_G}^G N_{C_2}^G MU_{\mathbf{R}}$. This leads to an equivariant self map

$$\Sigma^{256}\tilde{\Omega} \rightarrow \tilde{\Omega}.$$

It is an ordinary homotopy equivalence, and we will see that this implies formally that it induces an equivalence on homotopy fixed point sets.

1.22

Proving the Periodicity Theorem (continued)

The key tool for studying differentials in the slice spectral sequence is the geometric fixed point spectrum, denoted by $\Phi^G X$ for a G -spectrum X . This is an equivariant construction that will be explained later by Greenlees and May. It has much nicer properties than the usual fixed point spectrum X^G .

In particular it is known that for any cyclic 2-group G ,

$$\Phi^G N_{C_2}^G MU_{\mathbf{R}} = MO,$$

the unoriented cobordism spectrum. Its homotopy type has been well understood since Thom's work in the 50s.

Moreover there is a theorem saying that inverting a certain element in the slice spectral sequence converging to π_*X^G gives one converging to $\pi_*\Phi^GX$. Our knowledge of the latter group gives us a very good handle on the former spectral sequence.

1.23

Proving the Periodicity Theorem (continued)

Typically one proves theorems about differentials in such spectral sequences by means of some sort of extended power construction. In our case, all of the necessary geometry is encoded in the relation between π_*MU and π_*MO !

1.24

8 The detection theorem

Proving the detection theorem

The proof of the detection theorem is a calculation with the Adams-Novikov spectral sequence. It is the one part of our proof that could have been done 20 years ago. [In the first talk on Thursday, Behrens will introduce us to the necessary tools.](#)

He will also show us how they were used over 30 years ago to prove an odd primary analog (for $p > 3$) of our theorem. In that proof a key tool is a homomorphism from the E_2 -term of the Adams-Novikov spectral sequence for the sphere spectrum to the cohomology of C_p with certain coefficients. It is based on the fact that a formal group law of height $p - 1$ can have nontrivial automorphisms of order p .

[The proof of the detection theorem will be explained in the second talk on Thursday.](#) The key fact here is that at the prime 2 a formal group law of height 4 can have nontrivial automorphisms of order 8.

1.25

9 The slice and reduction theorems

Proving the slice theorem

Recall that a pivotal step in our proof is the Slice Theorem, which identifies the layers in the slice tower for MUR and its relatives. [On Friday Dugger will explain how it follows from the Reduction Theorem, which will be stated below.](#)

For each cyclic 2-group $G = C_{2^{n+1}}$ there is a certain equivariant (noncommutative) ring spectrum A which is a certain wedge of slice cells. It maps to $N_{C_2}^G MUR$ in such a way that the underlying wedge of spheres hits all of the underlying homotopy of $MU^{(2^n)}$. Thus both $N_{C_2}^G MUR$ and S^0 are A -modules.

Reduction Theorem. *The A -smash product $N_{C_2}^G MUR \wedge_A S^0$ is equivariantly equivalent to the integer Eilenberg-Mac Lane spectrum $H\mathbf{Z}$.*

The proof of this is the hardest calculation in our paper. [It will be the subject of the last two talks on Friday, to be given by Hill and Hopkins.](#)

1.26