


Noel J. Hicks

# Notes on Differential Geometry

with 25 figures and 100 problems



Revised and modernized edition by



TEXromancers



# TeXromancers

A collaborative typesetting project

---

This book is the product of a community effort. It was typeset by TeXromancers: an enthusiast group of mathematicians (for the most part), consisting of people organized on Discord. The reader should contact [amanzoo1@asu.edu](mailto:amanzoo1@asu.edu) if they are interested in joining the group.

---

## More books in this series:

- J. F Adams, *Stable Homotopy and Generalised Homology*
  - Available at <https://people.math.rochester.edu/faculty/doug/otherpapers/Adams-SHGH-latex.pdf>
- Noel J. Hicks, *Notes on Differential Geometry*
  - Available at <https://aareyanmanzoor.github.io/assets/hicks.pdf>
- Hideyuki Matsumura, *Commutative Algebra*
  - (Coming soon!)

Noel J. Hicks

# Notes on Differential Geometry

Revised and modernized edition by



TeXromancers

Typeset by T<sub>E</sub>Xromancers.

This digital book is meant for educational and scholarly purposes only, itself being a derivative work containing a transcription of a copyrighted publication. We claim no ownership of the original work's contents; all rights to the original book's contents belong to the respective publisher.

We believe that our use of the original work constitutes *fair use*, as defined under section 107 of the United States Copyright Act of 1976, wherein allowance is made for "fair use" for purposes such as criticism, comment, news reporting, teaching, scholarship, education and research.

Commercialization of this digital book is prohibited.

Original edition: 1965

This edition: 2022

# Preface to New Typesetting

This book is a re-LaTeX'ed version of the book “Notes on Differential Geometry” by Hicks. We're the T<sub>E</sub>Xromancers, a group of mostly math enthusiasts wanting to create more readable content for the math world. The typesetting credits go to: Aareyan Manzoor, George Coote, RokettoJanpu, Carl Sun, Andrew Lin, Nathaniel Alloway, Prakhar Agarwal, Shuayb Mohammed, Bastián Núñez, Arpit Mittal.

Here is a link to a dyslexia friendly version of the book: <https://aareyanmanzoor.github.io/assets/hicks-dyslexic.pdf>

Some notations in the original book are rather ancient or just not as popular, so we decided to change these to their modern counterpart. For example, set builders are written using  $[ ]$  in the original book; we decided to use  $\{ \}$ . We also used  $T_m M$ ,  $TM$  for the tangent space at  $m$ , the tangent bundle of  $M$ , rather than  $M_m$  and  $T(M)$ . We further used  $\Omega^p(M)$  to denote the space of  $p$ -forms, and  $T^{p,q}V$  to denote mixed tensors over some vector space  $V$ . We changed  $L_X Y$  to  $\mathcal{L}_X Y$  for the Lie derivative. However, we decided to keep the name *connexion* instead of the modern name *connection* as it has some charm.

The original book had a lot of inline equations to its detriment, and we decided to display mode some of these. A lot of things listed were made into an itemized list also. Some sentences were changed for better readability. A lot of the problems were split up into parts labeled by roman numbers.

We added citations and references with hyperlinks. References to e.g. theorems in the book are in blue, while citations to the bibliography is in red. The bibliography also has URLs now, for easy access. Some of the books in the bibliography had newer editions, so we went with those.

All the diagrams have been redrawn, special thanks to our artists: Yohan Wittgenstein, Bastián Núñez.

# Preface

The following paragraph presents a very brief history of differential geometry and the notation used in these notes.

Differential geometry is probably as old as any mathematical discipline and certainly was well launched after Newton and Leibniz had laid the foundations of calculus. Many results concerning surfaces in 3-space were obtained by Gauss in the first half of the nineteenth century, and in 1854 Riemann laid the foundations for a more abstract approach. At the end of that century, Levi-Civita and Ricci developed the concept of parallel translation in the classical language of tensors. This approach received a tremendous impetus from Einstein's work on relativity. During the early years of this century, E. Cartan initiated research and methods that were independent of a particular coordinate system (invariant methods). Chevalley's book [Che46] continued the clarification of concepts and notation, and it has had a remarkable effect on the current situation. The complete global synthesis of Cartan's approach was achieved when Ehresmann formulated a connexion<sup>1</sup> in terms of a fiber bundle. These notes utilize an invariant local method formulated by Koszul.

The first three chapters of this book provide a short course on classical differential geometry and could be used at the junior

---

<sup>1</sup>Editors' note: *connexion* means *connection* here and in the rest of the text. The latter is more common in modern usage, while the former likely comes from the corresponding French term. We have decided nonetheless to keep its use throughout the book.

level with a little outside reading in linear algebra and advanced calculus. The first six chapters can be used for a one-semester course in differential geometry at the senior-graduate level. Such a course would cover the main topics of classical differential geometry (except for the material in chapter 8) using modern language and techniques, and it would prepare a student for further study in the books of Helgason, Lang, Sternberg, etc. (see list in following paragraph). The entire book can be covered in a full year course. A selection of chapters could make up a “topics” course or a course on Riemannian geometry. For example, a course on manifolds and connexions could consist of chapters 1, 4, 5, 7 and sections 9.1, 9.3 and 9.4. The reader with a little experience should move through the first three chapters fairly quickly.

The problems are of several types: (a) those that provide explicit computations to test the understanding of the theory, (b) those that require the student to prove theorems similar to those in the text, (c) those that lead the student through supplementary material, some of which may be an integral part of the exposition, and (d) those that lead the student to books or papers in the literature. An introduction to bundle theory and the theory of Lie groups is covered via problem material. Our hope is to give the reader a solid understanding of the basic concepts and to stimulate him to further reading and thinking in differential geometry.

Besides the specific references found in the notes, we would like to mention the following general references:

- Point set topology: [Kel17], [HY12] and [PB14].
- Linear algebra: [Hal17] and [Jac13]
- Advanced calculus: [Buc03], [Kap03] and [HS13]
- Classical differential geometry: [Eis15], [HC99] and [Str61].
- Contemporary differential geometry: [AM12], [BC11], [Gug12], [Hel12], [KN63], [Lan14], [KN63] and [Ste64]



- History of differential geometry: [Str61] and [VW60]

We will use the following conventions: “iff” for “if and only if”; “□” for “Q.E.D.”; Cartan<sup>3</sup> will refer to the third reference in the bibliography under Cartan, and when there is only one reference for an author, we omit the superscript 1;  $\sum_{i=1}^n$ ,  $\sum_i$ ,  $\sum$  will all be used to indicate a sum is to be made, and in the latter two cases, we hope the omitted information (range or index of summation) is clear from the context.

At this time I would like to express my gratitude to former teachers N. Schwid and V.J. Varineau for their early encouragement, to Miss Margaret M. Genova and Miss Gillian D. Hodge for their help in typing the manuscript, and to L.M. Dickens for his contribution to the understanding of the illustrations. Finally, I am indebted to W. Ambrose and H. Samelson for sharing their insights via courses, notes, and conversations.

N.J. Hicks

*Ann Arbor, Michigan*  
May 1964



# Contents

<b>Preface to New Typesetting</b>	<b>5</b>
<b>Preface</b>	<b>7</b>
<b>1 Manifolds</b>	<b>15</b>
1.1 Manifolds . . . . .	15
1.2 Smooth Functions . . . . .	19
1.3 Vectors and Vector Fields . . . . .	21
1.4 The Jacobian of a Map . . . . .	26
1.5 Curves and Integral Curves . . . . .	27
1.6 Submanifolds . . . . .	30
<b>2 Hypersurfaces of <math>\mathbb{R}^n</math></b>	<b>39</b>
2.1 The Standard Connexion on $\mathbb{R}^n$ . . . . .	39
2.2 The Sphere Map and the Weingarten Map . . . . .	43
2.3 The Gauss Equation . . . . .	48
2.4 The Gauss Curvature and Codazzi-Mainardi Equations . . . . .	51
2.5 Examples . . . . .	52
2.6 Some Applications . . . . .	59
<b>3 Surfaces in <math>\mathbb{R}^3</math></b>	<b>65</b>
3.1 Smoothness and the Neighborhood of a non-Umbilic Point	65
3.2 Surfaces of Constant Curvature . . . . .	68
3.3 Parallel Surfaces (Normal Maps) . . . . .	72
3.4 Examples (Surfaces of Revolution) . . . . .	74

3.5	Lines of Curvature	76
<b>4</b>	<b>Tensors and Forms</b>	<b>81</b>
4.1	Tensors and Forms	81
<b>5</b>	<b>Connexions</b>	<b>93</b>
5.1	Invariant Viewpoint	93
5.2	Cartan Viewpoint	98
5.3	Coordinate Viewpoint	100
5.4	Difference Tensor of Two Connexions	101
5.5	Bundle Viewpoint	103
<b>6</b>	<b>Riemannian Manifolds and Submanifolds</b>	<b>109</b>
6.1	Length and Distance	109
6.2	Riemannian Connexion and Curvature	112
6.3	Curves in Riemannian Manifolds	115
6.4	Submanifolds	117
6.5	Hypersurfaces	120
6.6	Cartan Viewpoint and Coordinate Viewpoint	124
6.7	Canonical Spaces of Constant Curvature	126
6.8	Existence	129
<b>7</b>	<b>Operators on Forms and Integration</b>	<b>137</b>
7.1	Exterior Derivative	137
7.2	Contraction	140
7.3	Lie Derivative	141
7.4	General Covariant Derivative	144
7.5	Integration of Forms and Stokes' Theorem	148
7.6	Integration in a Riemannian Manifold	152
<b>8</b>	<b>Gauss-Bonnet Theory and Rigidity</b>	<b>159</b>
8.1	Gauss-Bonnet Formula	159
8.2	Index Theorem	167
8.3	Gauss-Bonnet Form	170
8.4	Characteristic Forms	173

8.5 Rigidity Problems . . . . .	177
<b>9 Existence Theory</b>	<b>183</b>
9.1 Involutive Distributions and the Frobenius Theorem . . .	183
9.2 The Fundamental Existence Theorem for Hypersurfaces	189
9.3 The Exponential Map . . . . .	193
9.4 Convex Neighborhoods . . . . .	197
9.5 Special Coordinate Systems . . . . .	200
9.6 Isothermal Coordinates and Riemannian Surfaces . . . .	203
<b>10 Topics in Riemannian Geometry</b>	<b>209</b>
10.1 Jacobi Fields and Conjugate Points . . . . .	209
10.2 First and Second Variation Formulae . . . . .	216
10.3 Geometric Interpretation of Riemannian Curvature . . .	225
10.4 The Morse Index Theorem . . . . .	228
10.5 Completeness . . . . .	237
10.6 Manifolds with Constant Riemannian Curvature . . . . .	243
10.7 Manifolds without Conjugate Points . . . . .	246
10.8 Manifolds with Non-Positive Curvature . . . . .	249
<b>Bibliography</b>	<b>255</b>
<b>Index</b>	<b>265</b>



# 1. Manifolds

In this chapter we define the fundamental concepts which we deal with throughout these notes. Specifically, the notions of manifold, function, and vector, and the concept of differentiability (smoothness), must be carefully digested for a solid foundation.

## 1.1 Manifolds

First some notation. Let  $\mathbb{R}$  be the set of real numbers. For an integer  $n > 0$ , let  $\mathbb{R}^n$  be the product space of ordered  $n$ -tuples of real numbers. Thus

$\mathbb{R}^n = \{(a_1, \dots, a_n) : a_i \in \mathbb{R}\}$ . For  $i = 1, \dots, n$ , let  $u_i$  be the natural coordinate (slot) functions of  $\mathbb{R}^n$ , i.e.,  $u_i: \mathbb{R}^n \rightarrow \mathbb{R}$  by  $u_i(a_1, \dots, a_n) = a_i$ . An open set of  $\mathbb{R}^n$  will be a set which is open in the standard metric topology induced by the standard metric function  $d$  on  $\mathbb{R}^n$ , thus if  $a = (a_1, \dots, a_n)$  and  $b = (b_1, \dots, b_n)$  are points in  $\mathbb{R}^n$ , then

$$d(a, b) = \left[ \sum_{i=1}^n (a_i - b_i)^2 \right]^{1/2}.$$

The concept of differentiability is based ultimately on the definition of a derivative in elementary calculus. Let  $r$  be an integer,  $r > 0$ . Recall from advanced calculus that a map  $f$  from an open set  $A \subset \mathbb{R}^n$  into  $\mathbb{R}$  is called  $C^r$  on  $A$  if it possesses continual partial derivatives on  $A$  of all orders  $\leq r$ . If  $f$  is merely continuous from  $A$  to  $\mathbb{R}$ , then  $f$  is  $C^0$  on  $A$ . If  $f$  is  $C^r$  on  $A$  for all  $r$ , then  $f$  is  $C^\infty$  on  $A$ . If  $f$  is real analytic on  $A$  (expandable in a power series in

the coordinate functions about each point of  $A$ ), then  $f$  is  $C^\omega$  on  $A$ . Henceforth, unless otherwise specified, we let  $r$  be  $\infty$ ,  $\omega$ , or an integer  $> 0$ .

A map  $f$  from an open set  $A \subset \mathbb{R}^n$  into  $\mathbb{R}^k$  ( $k$  an integer  $\geq 1$ ) is  $C^r$  on  $A$  if each of its slot functions  $f_i = u_i \circ f$  is  $C^r$  for  $i = 1, \dots, k$ ; thus for  $p$  in  $\mathbb{R}^n$ ,  $f(p) = (f_1(p), \dots, f_k(p)) \in \mathbb{R}^k$ .




Figure 1.1: Overlapping Coordinate Domains

We now define a manifold. Let  $M$  be a set. An  $n$ -coordinate pair on  $M$  is a pair  $(\phi, U)$  consisting of a subset  $U$  of  $M$  and a 1 to 1 map  $\phi$  of  $U$  onto an open set in  $\mathbb{R}^n$ . One  $n$ -coordinate pair  $(\phi, U)$  is  $C^r$  related to another  $n$ -coordinate pair  $(\psi, V)$  if the maps  $\phi \circ \theta^{-1}$  and  $\psi \circ \phi^{-1}$  are  $C^r$  maps wherever they are defined (thus their domains of definition must be open)<sup>1</sup>. A  $C^r$   $n$ -subatlas on  $M$  is a collection of  $n$ -coordinate pairs  $(\phi_h, U_h)$ , each of which is  $C^r$  related to every other member of the collection, and the union of sets  $U_h$  is  $M$ . A maximal collection of  $C^r$  related  $n$ -coordinate pairs is called a  $C^r$   $n$ -atlas. If a  $C^r$   $n$ -atlas contains a  $C^r$   $n$ -subatlas, we

---

<sup>1</sup>See Figure 1.1.



say that the subatlas induces or generates the atlas. Finally, an  $n$ -dimensional  $C^r$  manifold or a  $C^r$   $n$ -manifold is a set  $M$  together with a  $C^r$   $n$ -atlas. When  $r = 0$ ,  $M$  is customarily called a *locally Euclidean space* or a *topological manifold*, and only when  $r \neq 0$  is  $M$  called a *differentiable* or *smooth manifold*. An atlas on a set  $M$  is often called a *differentiable structure* or a *manifold structure* on  $M$ . Notice that one set may possess more than one differentiable structure (see example 4 below), however, a definition of “equivalent” differentiable structures is necessary before the study of different atlases on a set becomes meaningful (see [Mun66]).

Each  $n$ -coordinate pair  $(\phi, U)$  on a set  $M$  induces a set of  $n$  real valued functions on  $U$  defined by  $x_i = u_i \circ \phi$  for  $i = 1, \dots, n$ . The functions  $x_1, \dots, x_n$  are called *coordinate functions* or a *coordinate system* and  $U$  is called the *domain* of the coordinate system.

We list some examples:

1. Let  $M$  be  $\mathbb{R}^n$  with a  $C^r$   $n$ -subatlas equal to the pair consisting of  $\phi =$  the identity map and  $U = \mathbb{R}^n$ .
2. Let  $M$  be any open set of  $\mathbb{R}^n$  and let a  $C^r$   $n$ -subatlas be (the identity map,  $M$ ).
3. Let  $M = \text{GL}(n, \mathbb{R})$ , the group of non-singular  $\mathbb{R}$ -linear transformations of  $\mathbb{R}^n$  onto itself. Then  $M$  can be mapped 1:1 onto an open set in  $\mathbb{R}^{n^2}$  and thus a manifold structure can be defined on  $M$  via example 2. If  $(a_{ij})$  is a matrix representation of an element of  $M$  with respect to the usual base of  $\mathbb{R}^n$ , then map  $(a_{ij})$  into the  $n^2$ -tuple

$$(a_{11}, a_{12}, \dots, a_{1n}, a_{21}, a_{22}, \dots, a_{2n}, a_{31}, \dots, a_{nn}).$$

The image set of this map will be open since it is the inverse image of an open set by the determinant map, which is continuous (indeed it is  $C^\omega$  as a map on  $\mathbb{R}^{n^2}$ ).

4. Let  $M_1$  be the 1-dimensional  $C^1$  manifold of example 1, and let  $M_2 = \mathbb{R}$  with the  $C^1$  1-subatlas  $(x^3, \mathbb{R})$ , where  $x$  is the identity mapping on  $\mathbb{R}$ . Then  $M_1 \neq M_2$  since  $x^{1/3}$  is not  $C^1$  at the origin.
5. Let  $f$  be a  $C^r$  real valued function on  $\mathbb{R}^{n+1}$ , with  $r > 0$  and  $n > 0$ , and suppose the gradient of  $f$  does not vanish on an  $f$ -constant set  $M = \{p \in \mathbb{R}^{n+1} : f(p) = 0\}$ . Then at each point in  $M$ , choose any partial derivative of  $f$  that doesn't vanish, say the  $i$ th one. Apply the implicit function theorem to obtain a neighborhood of  $p$  (relative topology on  $M$ ) which projects in a 1:1 way into the  $u_i = 0$  hyperplane of  $\mathbb{R}^{n+1}$ . This can then be used to define a subatlas which makes  $M$  a  $C^r$   $n$ -manifold.  
 This example covers many classical hypersurfaces in  $\mathbb{R}^{n+1}$ , including spheres, planes, and cylinders.
6. The process in example 5 can easily be generalized to obtain  $C^r$   $(n-k)$ -manifolds from "constant sets" of a  $C^r$  map  $f: \mathbb{R}^n \rightarrow \mathbb{R}^k$  whose Jacobian matrix is of rank  $k$  on the constant set<sup>2</sup>.
7. Let  $F$  be a univalent map from an open set in  $\mathbb{R}^n$  into  $\mathbb{R}^m$ , with  $0 < n < m$ , and let  $M$  be the image of  $F$ . Then the  $n$ -coordinate pair  $(F^{-1}, M)$  defines a  $C^r$   $n$ -subatlas on  $M$ .

For further definitions, let  $M$  be a fixed  $C^r$   $n$ -manifold. An open set in  $M$  is a subset  $A$  of  $M$  such that  $\phi(A \cap U)$  is open in  $\mathbb{R}^n$  for every  $n$ -coordinate pair  $(\phi, U)$ . The reader can verify that  $M$  becomes a topological space with this definition of the open sets. If  $p \in M$ ,

---

<sup>2</sup>Suppose  $f: U \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  is a  $C^r$  map, where  $m > n$ . A point  $q \in f(U)$  is called a *regular value* of  $f$  if the Jacobian of  $f$  is surjective at each point  $p \in f^{-1}(\{q\})$ . Thus we can reword this example as follows: if  $q$  is a regular value of  $f$ , then  $M = f^{-1}(\{q\})$  is a  $(m-n)$ -dimensional submanifold of  $\mathbb{R}^m$ . Similarly if  $f: M \rightarrow N$  is a  $C^r$  map between two manifolds  $M$  and  $N$ , of dimension  $m$  and  $n$  resp. with  $m > n$ , a point  $q \in N$  is called a *regular value* of  $f$  if the differential  $f_*: \mathbf{T}_m M \rightarrow \mathbf{T}_{f(m)} N$  (cf. section 1.4) is surjective at each point  $p \in f^{-1}(\{q\})$ . Then for regular values  $q$ ,  $S = f^{-1}(\{q\})$  is a  $(m-n)$ -dimensional submanifold of  $M$ . This is a standard result and follows from the implicit and inverse function theorems, see e.g. [Hir76, Chapter 1, Theorem 3.2].

then a *neighborhood of  $p$*  is any open set containing  $p$ . Notice  $M$  need not be Hausdorff. The concept of Hausdorffness is irrelevant for much of local differential geometry. It becomes relevant in passing from a Riemannian metric to a distance function.

## 1.2 Smooth Functions

In this section let  $A$  be the domain of a function  $f$  and assume  $A$  is an open subset of the  $C^r$   $n$ -manifold  $M$ . If  $f$  is real valued, then  $f$  is  $C^s$  on  $A$  if  $f \circ \phi^{-1}$  is  $C^s$  on  $\phi(A \cap U)$  for every coordinate pair  $(\phi, U)$  on  $M$ . Note the independence of  $r$  and  $s$ . If  $N$  is a  $C^k$   $d$ -manifold and  $f$  is  $N$ -valued, then  $f$  is  $C^s$  on  $A$  if  $f$  is continuous and for every real valued function  $g$ , that is  $C^s$  on an open domain in  $N$ , the composite  $g \circ f$  is  $C^s$  on  $A \cap f^{-1}$  (domain of  $g$ ). Note the independence of  $r$ ,  $k$ , and  $s$ .




Figure 1.2: An Induced Map from  $\mathbb{R}^2$  into  $\mathbb{R}$

The local character of smoothness of a function is captured in the following definition. Suppose the domain of  $f$  is not necessarily open and  $f$  is  $N$ -valued. If  $p$  is in the domain of  $f$ , then  $f$  is  $C^s$  at

$p$  if there is a neighbourhood  $U$  of  $p$  with  $f$  defined on  $U$  such that  $f|_U$  is  $C^s$  on  $U$ . As a corollary, if  $f$  is  $C^s$  at every point in its domain then its domain is open.

Let us now specialize to  $C^\infty$  manifolds and  $C^\infty$  functions. This is done for convenience chiefly and it allows us to define a tangent vector in a very elegant way. Our concern in these notes is not with “the least possible assumptions” but rather with those concepts that arise naturally in a general situation. The restriction is not too drastic because of the following result due to Whitney: A  $C^r$  atlas on a set with  $r > 0$  contains a  $C^\infty$  atlas (see [Mun66]). There is an example of Kervaire which exhibits a  $C^0$  atlas on a set which admits no  $C^1$  atlas. For further work on the “equivalence” of differentiable structures see [MW97] and [Mil56], [Mun66] and [Mun59], and [Sma61].

### Problems

The following list of nine problems are recommended in order to familiarize oneself with the notion of a  $C^\infty$  map. In particular the problems are aimed at obtaining numbers 6 and 7 which are often useful. The list (remember  $A$  is open in  $M$ , which is a  $C^\infty$   $n$ -manifold);

1. The map  $f: A \rightarrow N$  is  $C^\infty$  on  $A$  iff  $f$  is  $C^\infty$  at each point  $p$  in  $A$ .
2. If  $f: A \rightarrow N$ ,  $f$  is  $C^\infty$  on  $A$ , and  $U$  is an open set contained in  $A$ , then  $f|_U$  is  $C^\infty$  on  $U$ .
3. Let  $U_h$  be a collection of open sets in  $M$  and let  $f_h: U_h \rightarrow N$  be  $C^\infty$  on  $U_h$  for each  $h$ . If  $f$  is a function whose domain is the union of all  $U_h$  and if  $f|_{U_h} = f_h$  for all  $h$ , then  $f$  is  $C^\infty$  on its domain.
4. If  $f: A \rightarrow \mathbb{R}^k$  is  $C^\infty$  on  $A \subset \mathbb{R}^n$  and  $g: B \rightarrow \mathbb{R}$  is  $C^\infty$  on the open set  $B \subset \mathbb{R}^k$ , then  $g \circ f$  is  $C^\infty$  on  $A \cap f^{-1}(B)$ .

5. If  $f: A \rightarrow N$  is  $C^\infty$  on  $A \subset M$  and  $(\phi, U)$  is a coordinate pair on  $M$ , then  $f \circ \phi^{-1}$  is  $C^\infty$  on  $\phi(A \cap U)$ .
6. Let  $P$  be a  $C^\infty$   $s$ -manifold. If  $F: A \rightarrow N$  is  $C^\infty$  on  $A \subset M$  and  $g: B \rightarrow P$  is  $C^\infty$  on the open set  $B \subset N$ , then  $g \circ f$  is  $C^\infty$  on  $A \cap f^{-1}(B)$ .
7. The map  $f: A \rightarrow N$  is  $C^\infty$  on  $A \subset M$  iff for every coordinate pair  $(\phi, U)$  in a subatlas on  $N$  the functions  $x_i \circ f$  are  $C^\infty$  on  $A \cap f^{-1}(U)$ , for  $i = 1, \dots, d$  and  $x_i = u_i \circ \phi$ .
8. If  $n \geq k$  and  $g: \mathbb{R}^n \rightarrow \mathbb{R}^n$  by  $g(a_1, \dots, a_n) = (a_1, \dots, a_k)$  then  $g$  is  $C^\infty$  on  $\mathbb{R}^n$ . If  $h: \mathbb{R}^k \rightarrow \mathbb{R}^n$  by  $h(a_1, \dots, a_k) = (a_1, \dots, a_k, 0, \dots, 0)$  then  $h$  is  $C^\infty$  on  $\mathbb{R}^k$ .
9. Let  $f$  and  $g$  be real valued functions that are  $C^\infty$  on the subsets  $A$  and  $B$  of  $M$ , respectively. Show that  $f+g$  and  $fg$  are  $C^\infty$  on  $A \cap B$ , where  $(f+g)(p) = f(p) + g(p)$  and  $(fg)(p) = f(p)g(p)$ .

For the record, we can and so do define a Lie group. A Lie group  $G$  is a group  $G$  whose underlying set is also a  $C^\infty$  manifold such that the group operations are  $C^\infty$ , i.e., the map  $\phi: G \times G \rightarrow G$  where  $\phi(g, h) = gh^{-1}$  is  $C^\infty$  (see problem 18 and 20).

One last bit of notation, let  $C^\infty(A, N)$  denote the set of  $C^\infty$  functions mapping an open set  $A$  in a manifold  $M$  into a manifold  $N$ .

### 1.3 Vectors and Vector Fields

The definition of a tangent vector generalizes the “directional derivative” in  $\mathbb{R}^n$ . If  $X$  is an ordinary (advanced calculus) vector at a point  $m$  in  $\mathbb{R}^n$  and  $f$  is a  $C^\infty$  function in a neighborhood of  $m$ , then define  $X_m f = X_m \cdot (\nabla f)_m$ , where  $\nabla f$  is the gradient vector field of  $f$ . From the properties of the “dot” product and the operator  $\nabla$ ,

it follows that

$$\begin{aligned} X_m(af + bg) &= aX_mf + bX_mg \\ X_m(fg) &= f(m)X_mg + g(m)X_mf, \end{aligned}$$

where  $g$  is a  $C^\infty$  function in a neighborhood of  $m$  and  $a$  and  $b$  are real numbers. Notice  $X$  is not normalized to be a unit vector. We generalize now to define a tangent vector on a manifold as an operator on  $C^\infty$  functions which obeys the above rules.

Let  $M$  be a  $C^\infty$   $n$ -manifold. Let  $m$  be in  $M$  and let  $C^\infty(m)$  denote the set of real valued functions that are  $C^\infty$  on some neighborhood of  $m$ . A tangent vector at  $m$  is a real valued function  $X$  on  $C^\infty(m)$  having the following properties:

- (1)  $X(f + g) = Xf + Xg$ ,  $X(bf) = b(Xf)$
- (2)  $X(fg) = (Xf)g(m) + f(m)(Xg)$

where  $f$  and  $g$  are in  $C^\infty(m)$ , and  $b$  is in  $\mathbb{R}$ . The set  $C^\infty(m)$  is almost a ring (there is a slight problem with domains), and thus a tangent vector is often called a derivation on  $C^\infty(m)$ .

The tangent space to  $M$  at  $m$ , denoted by  $\mathbf{T}_mM$ , is the set of all tangent vectors at  $m$ . It is a vector space over the real field where  $(X + Y)f = Xf + Yf$  and  $(bX)f = b(Xf)$  for  $X, Y$  in  $\mathbf{T}_mM$ ,  $f$  in  $C^\infty(m)$ , and  $b$  a real number.

Let  $x_1, \dots, x_n$  be a coordinate system about  $m$  (i.e.,  $m$  is in the domain of these coordinate functions). We define for each  $i$ , a coordinate vector at  $m$ , denoted  $\left(\frac{\partial}{\partial x_i}\right)_m$  by

$$\left(\frac{\partial}{\partial x_i}\right)_m f = \frac{\partial(f \circ \phi^{-1})}{\partial u_i}(\phi(m))$$

where  $x_i = u_i \circ \phi$  and the differentiation on the right side is as usual on  $\mathbb{R}^n$ . The verification of properties (1) and (2) above we leave to the reader. In a moment we show these coordinate vectors form a base for the tangent space at  $m$ .

**Lemma 1.1.** Let  $x_1, \dots, x_n$  be a coordinate system about  $m$  with  $x_i(m) = 0$  for all  $i$ . Then for every function  $f$  in  $C^\infty(m)$  there exist  $n$  functions  $f_1, \dots, f_n$  in  $C^\infty(m)$  with  $f_i(m) = \left(\frac{\partial}{\partial x_i}\right)_m f$  and  $f = f(m) + \sum_i x_i f_i$  in a neighborhood of  $m$ . (Note the equality in question is an equality between functions, and  $f(m)$  represents a constant function with value  $f(m)$ ; the sum is taken for  $i = 1, 2, \dots, n$ , and in the future this relevant range is to be understood.)

*Proof.* Let  $\phi$  be the coordinate map belonging to the  $x_i$ . Let  $F = f \circ \phi^{-1}$  and we know  $F$  is defined in a ball about the origin in  $\mathbb{R}^n$ , i.e., in a set

$B = \{p \in \mathbb{R}^n : \text{distance from origin to } p < r\}$ . For  $(a_1, \dots, a_n)$  in  $B$  we have

$$\begin{aligned} F(a_1, \dots, a_n) &= F(a_1, \dots, a_n) - F(a_1, \dots, a_{n-1}, 0) \\ &\quad + F(a_1, \dots, a_{n-1}, 0) - F(a_1, \dots, a_{n-2}, 0, 0) + \dots \\ &\quad + F(a_1, 0, \dots, 0) - F(0, \dots, 0) + F(0, \dots, 0) \\ &= \sum_i F(a_1, \dots, a_{i-1}, ta_i, 0, \dots, 0) \Big|_0^1 + F(0, \dots, 0) \\ &= F(0, \dots, 0) + \sum_i \int_0^1 \frac{\partial F}{\partial t_i}(a_1, \dots, a_{i-1}, ta_i, 0, \dots, 0) a_i dt \\ &= F(0, \dots, 0) + \sum_i a_i F_i(a_1, \dots, a_n), \end{aligned}$$

where

$$F_i(a_1, \dots, a_n) = \int_0^1 \frac{\partial F}{\partial u_i}(a_1, \dots, a_{i-1}, ta_i, 0, \dots, 0) dt$$

is  $C^\infty$  in  $B$  since  $\left(\frac{\partial}{\partial u_i}\right)$  is  $C^\infty$ . Let  $f_i = F_i \circ \phi$  and the lemma is proved. □

**Theorem 1.2.** Let  $M$  be a  $C^\infty$   $n$ -manifold and let  $x_1, \dots, x_n$  be a coordinate system about  $m$  in  $M$ . Then if  $X \in \mathbf{T}_m M$ ,  $X = \sum_i (X x_i) \left(\frac{\partial}{\partial x_i}\right)_m$ , and the coordinate vectors form a base for  $\mathbf{T}_m M$  which thus has

dimension  $n$ .

*Proof.* We first prove the stated representation. Take  $X \in \mathbf{T}_m M$  and  $f \in C^\infty(m)$ . If  $x_i(m) \neq 0$  for all  $i$ , let  $y_i = x_i - x_i(m)$ . Then apply the lemma to  $f$  with respect to the coordinate system  $y_1, \dots, y_n$  and notice  $\left(\frac{\partial f}{\partial y_i}\right)(m) = \left(\frac{\partial f}{\partial x_i}\right)(m)$ . Next we see if  $c$  a constant map then

$$X(c) = cX(1) = c(1X(1) + 1X(1)) = 2cX(1)$$

which implies  $cX(1) = 0$  and  $X(c) = 0$ . Thus

$$\begin{aligned} Xf &= X\left(f(m) + \sum_i y_i f_i\right) \\ &= \sum_i ((Xy_i)f_i(m) + y_i(m)(Xf_i)) \\ &= \sum_i X(x_i - x_i(m))f_i(m) \\ &= \sum_i (Xx_i)\left(\frac{\partial f}{\partial x_i}\right)(m) \end{aligned}$$

which proves the required representation. If  $Y = \sum_i a_i \left(\frac{\partial}{\partial x_i}\right) = 0$  then  $0 = Yx_j = a_j$ , thus the coordinate vectors are independent and span  $\mathbf{T}_m M$ .  $\square$

A vector field  $X$  on a set  $A$  is a mapping that assigns to each point  $p$  in  $A$  a vector  $X_p$  in  $M_p$ . A field  $X$  is  $C^\infty$  on  $A$  if  $A$  is open and for each real valued function  $f$  that is  $C^\infty$  on  $B$ , the function  $(Xf)(p) = X_p f$  is  $C^\infty$  on  $A \cap B$ . If  $X$  and  $Y$  are  $C^\infty$  vector fields on  $A$  their Lie bracket is a  $C^\infty$  vector field  $[X, Y]$  on  $A$  defined by  $[X, Y]_p f = X_p(Yf) - Y_p(Xf)$ .

If  $f$  and  $g$  are  $C^\infty$  functions, it is trivial that  $[X, Y](f+g) = [X, Y]f + [X, Y]g$ , and  $[X, Y](af) = a[X, Y]f$  for  $a$  in  $\mathbb{R}$ . To check the product



property, consider

$$\begin{aligned}
 [X, Y](fg) &= X(Y(fg)) - Y(X(fg)) \\
 &= X(fYg + gYf) - Y(fXg + gXf) \\
 &= fXYg + (Xf)(Yg) + (Xg)(Yf) + gXYf \\
 &\quad - fYXg - (Yf)(Xg) - (Yg)(Xf) - gYXf \\
 &= f[X, Y]g + g[X, Y]f.
 \end{aligned}$$

Thus  $[X, Y]$  is a vector field and the proof of its  $C^\infty$  nature we leave as a problem.

For later use, notice that  $[X, Y] = -[Y, X]$ ,  $[X, X] \neq 0$ , and the bracket is linear in each slot with respect to addition, i.e.,  $[X_1 + X_2, Y] = [X_1, Y] + [X_2, Y]$ . However,

$$[fX, gY] = f(Xg)Y - g(Yf)X + fg[X, Y]$$

and it is this property that prevents the bracket mapping from being a tensor (problem 10). Problem 13 gives a geometric interpretation of the bracket, and in section 9.1 there are applications involving integrability conditions. For example, if  $x_1, \dots, x_n$  is a coordinate system then  $\left[\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right] = 0$  for all  $i$  and  $j$  (since cross partial derivatives of  $C^\infty$  functions are equal), and actually this condition on  $n$  independent vector fields is sufficient to imply the fields are coordinate vector fields (section 9.1).

The bracket operation also satisfies the following expression which is called the *Jacobi identity*,

$$[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] = 0 \quad \text{(Jacobi Identity)}$$

where  $X, Y$ , and  $Z$  are  $C^\infty$  fields with a common domain.

## 1.4 The Jacobian of a Map

Let  $M$  and  $N$  be  $C^\infty$  manifolds of dimensions  $n$  and  $k$  respectively. We defined the above concept of a  $C^\infty$  map  $f$  from  $M$  into  $N$ . Such a map induces a linear transformation from each tangent space  $\mathbf{T}_m M$  into the tangent space  $\mathbf{T}_{f(m)} N$ . This linear map is called the *Jacobian map* or the *differential of  $f$*  and we denote it by  $f_*$  (often it is denoted  $df$ , but we reserve the symbol  $d$  for the exterior derivative<sup>3</sup> operator). Let  $X$  be in  $\mathbf{T}_m M$  and we define  $f_* X$  as a vector at  $f(m)$  and setting  $(f_* X)g = X(g \circ f)$ . It is trivial to check that  $f_* X$  is a vector at  $f(m)$  and the map  $f_*$  is linear.

By selecting a coordinate system  $x_1, \dots, x_n$  about  $m$  and another  $y_1, \dots, y_k$  about  $f(m)$ , we can determine a matrix representation for  $f_*$  which is called the *Jacobian matrix* of  $f_*$  with respect to the chosen coordinate systems. Let  $X_i = \frac{\partial}{\partial x_i}$ ,  $Y_j = \frac{\partial}{\partial y_j}$ , thus  $X_1, \dots, X_n$ , at  $m$ , form a base for  $\mathbf{T}_m M$  and we compute  $f_*$  by computing its action on this base. Namely,  $f_* X_i = \sum_j (f_* X_i) y_j Y_j$  by the representation theorem 1.2 above, hence the matrix in question is the matrix

$$((f_* X_i) y_j) = \left( \frac{\partial (y_j \circ f)}{\partial x_i} \right) \text{ for } 1 \leq i \leq n \text{ and } 1 \leq j \leq k.$$

The implicit function theorem and the inverse function theorem can be applied and formulated in this language. The former we postpone, since we do not really need it for some time (see Problem 16) but the latter is both useful and instructive. First a definition. A *diffeomorphism* is a map  $f: M \rightarrow N$  that is 1 to 1 and onto with both  $f$  and  $f^{-1}$   $C^\infty$ , and if such an  $f$  exists, then  $M$  is *diffeomorphic* to  $N$ .

**Theorem 1.3 (Inverse function).** Let  $M$  and  $N$  be  $C^\infty$   $n$ -manifolds and let  $f: M \rightarrow N$  be  $C^\infty$ . If for  $m$  in  $M$ , the Jacobian  $f_*$  at  $m$  is an isomorphism of  $\mathbf{T}_m M$  onto  $\mathbf{T}_{f(m)} N$ , then there is a neighborhood  $U$

---

<sup>3</sup>See sections 5.2 and 7.1

of  $m$  and a neighborhood  $V$  of  $f(m)$  such that  $f$  is a diffeomorphism from  $U$  to  $V$  (i.e.,  $f$  is a local diffeomorphism about  $m$ ).

We leave it to the reader to choose a coordinate system on both sides and apply the theorem from advanced calculus to obtain the result. Notice the  $C^\infty$  demand of  $f$  and  $f^{-1}$  implies the theorem could be stated as necessary as well as sufficient condition for the existence of a local inverse. If one only demands continuity of the inverse, then the map  $x \mapsto x^3$  provides a homeomorphism of  $\mathbb{R}$  onto  $\mathbb{R}$  whose Jacobian is singular at the origin.

Now consider the behavior of the Jacobian with respect to the composite maps. Let  $g$  be a  $C^\infty$  map of  $N$  into the  $C^\infty$  manifold  $L$ . Then at each  $m$  in  $M$ ,  $(g \circ f)_* = g_* \circ f_*$ , for if  $h$  is a  $C^\infty$  function about  $g(f(m))$  and  $X$  in  $\mathbf{T}_m M$  then  $((g \circ f)_* X)h = X(h \circ g \circ f) = (f_* X)(h \circ g) = (g_*(f_* X))h$ . In terms of coordinate systems, the above computation exhibits the chain rule and multiplicative behavior of Jacobian matrices. When  $f$  is a diffeomorphism of  $M$  into  $N$ , and  $X$  and  $Y$  are  $C^\infty$  fields on  $M$ , then  $f_* X$  and  $f_* Y$  are  $C^\infty$  fields on  $N$  with  $f_*[X, Y] = [f_* X, f_* Y]$ .

## 1.5 Curves and Integral Curves

In these notes curves will be viewed as a special case of mappings, thus we will deal with “parameterized curves” almost exclusively. A curve in  $M$  is a  $C^\infty$  map  $\sigma$  from an open subset of  $\mathbb{R}$  into  $M$ . Often we speak of a curve  $\sigma$  from  $[a, b]$  into  $M$  where  $[a, b]$  is a closed interval of real numbers, and in this case it is assumed the domain of  $\sigma$  is actually an open set in  $\mathbb{R}$  containing  $[a, b]$

Let  $\sigma$  be a curve in  $M$  with domain  $U$ . For each  $t$  in  $U$  define the tangent of  $\sigma$  at  $t$  to be the vector  $T(t)$ , or  $T_\sigma(t)$ , at  $\sigma(t)$  where  $T(t) = \sigma_* \left( \frac{d}{dt} \right)_t$  and  $\frac{d}{dt}$  denotes the usual differentiation operator of real valued  $C^\infty$  functions on  $\mathbb{R}$ . Thus if  $x, \dots, x_n$  a coordinate system

about  $\sigma(t)$ , then

$$T(t) = \sum_i \left( \frac{d(x_i \circ \sigma)}{dt} \right)_t \left( \frac{\partial}{\partial x_i} \right)_{\sigma(t)}.$$

By differentiating the coordinate parameter functions  $x_i \circ \sigma(t)$  one determines the coefficients of  $T(t)$  with respect to the coordinate vectors associated with the coordinate system. Notice this  $T(t)$  is the usual “velocity” vector associated with a parameterized curve in  $\mathbb{R}^3$ .




Figure 1.3: A Curve

Having the idea of curve and tangent vector we can give a geometric description of the Jacobian  $f_*$  associated with the map  $f : M \rightarrow N$ . For  $X$  in  $\mathbf{T}_m M$  choose any curve  $\sigma$  on  $M$  with  $\sigma(0) = m$  and  $T_\sigma(0) = X$ . Then  $f \circ \sigma$  is a curve on  $N$  with  $f \circ \sigma(0) = f(m)$  and indeed  $f_* X = T_{f \circ \sigma}(0)$ . Thus we “fill in the vector by a curve, map the curve to  $N$ , and take the new tangent vector.” This device is very useful if one knows geometrically the behavior of certain curves; e.g., let  $M = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1\}$ , let  $S$  be the unit sphere in  $\mathbb{R}^3$ , and let  $f : M \rightarrow S$  by  $f(x, y, z) = (x, y, 0)$ . The particular  $f$  just defined is called the “sphere map” or the “Gauss map” from

$M$  to  $S$ , since it essentially uses a unit normal vector field to  $M$  in its definition. Its Jacobian should be trivial to compute at each point from the above remarks.

We carry the idea of “filling in a vector” to a classical setting. Let  $X$  be a  $C^\infty$  vector field on the manifold  $M$ . A curve  $\sigma$  is an *integral curve* of  $X$  if whenever  $\sigma(t)$  is in the domain of  $X$  then  $T_{\sigma(t)} = X_{\sigma(t)}$ . Thus we say the curve  $\sigma$  “fits”  $X$ , and suggest the physical example of the velocity vector field (which gives  $X$ ) of a steady fluid flow and its streamlines (which give integral curves). The local existence of integral curves is guaranteed by the theory of ordinary differential equations.




Figure 1.4: An Integral Curve of a Vector Field

**Theorem 1.4.** Let  $X$  be a  $C^\infty$  vector field on  $M$  and let  $m$  be a point in the domain of  $X$ . Then for any real number  $b$  there exists a real number  $r > 0$  and a unique curve  $\sigma : (b - r, b + r) \rightarrow M$  such that  $\sigma(b) = m$  and  $\sigma$  an integral curve of  $X$ .

*Proof.* Let  $x_1, \dots, x_n$  be a coordinate system about  $m$  whose domain

$U$  is contained in the domain of  $X$ . Let  $X = \sum_i f_i \left( \frac{\partial}{\partial x_i} \right)$  define  $C^\infty$  real valued functions  $f_i$  on  $U$ . Then the condition that a curve  $\sigma$  be an integral curve of  $X$  becomes the condition

$$\frac{d(x_i \circ \sigma)}{dt} = f_i \circ \sigma$$

on the domain of  $\sigma$ , or writing (improperly) as usual  $x_i(t) = x_i \circ \sigma(t)$ , we have the system of first order ordinary differential equations

$$\frac{dx_i}{dt} = f_i(x_1, \dots, x_n),$$

for  $i = 1, \dots, n$ . Apply an existence and uniqueness theorem from differential equation theory to obtain  $r > 0$  and functions  $x_i(t)$  that define  $\sigma$  on the specified range with the required properties.  $\square$

Actually the theorem from differential equations gives much more than the above conclusion; it also includes the  $C^\infty$  dependence of solutions as we vary the initial parameter  $b$  and the point  $m$  (see section 9.3). We return to this later when discussing the existence of geodesics and the exponential map (sections 5.1 and 9.3). For global ramifications see [Pal57] or [Lan14].

It is convenient to define a broken  $C^\infty$  curve  $\sigma$  on an interval  $[a, b]$  to be a continuous map  $\sigma$  from  $[a, b]$  into  $M$  which is  $C^\infty$  on each of a finite number of subintervals  $[a, b_1], [b_1, b_2], \dots, [b_{k-1}, b]$ .

## 1.6 Submanifolds

A  $C^\infty$   $k$ -manifold is a submanifold of a  $C^\infty$   $n$ -manifold  $\bar{M}$  if for every point  $p$  in  $M$  there is a coordinate neighborhood  $\bar{U}$  of  $\bar{M}$  with coordinate functions  $\bar{x}_1, \dots, \bar{x}_n$  such that the set  $U = \{m \in \bar{U} : \bar{x}_{k+1}(m) = \dots = \bar{x}_n(m) = 0\}$  is a coordinate neighborhood of  $p$  in  $M$  with coordinate functions

$x_1 = \bar{x}_1|_U, \dots, x_k = \bar{x}_k|_U$ . These coordinate systems are called special or adapted coordinate systems.

Notice it is not required that  $M \cap \bar{U} = U$  so “slices” of  $M$  may approach other “slices” of  $M$  in  $\bar{M}$  (see problem 17) and hence the topology on  $M$  may not be the relative topology. The definition of submanifold implies  $M$  is a subset of  $\bar{M}$  and  $k \geq n$ . Letting  $i : M \rightarrow \bar{M}$  be the inclusion map, then  $i$  is  $C^\infty$  since  $\bar{x}_j \circ i$  are  $C^\infty$  maps for all special coordinate functions. The inclusion map is also an imbedding (see below) since the Jacobian  $i_*$  is non-singular, i.e.,  $i_*\left(\frac{\partial}{\partial x_j}\right)(p) = \frac{\partial}{\partial \bar{x}_j}(p)$  for  $j = 1, \dots, k$ . In these notes we will identify a tangent vector  $X$  in  $T_p M$  with its image in  $T_p \bar{M}$  unless there is a possibility of confusion (just as we identify  $p$  and  $i(p)$ ).

To make some more standard definitions, let  $M$  and  $\bar{M}$  be  $C^\infty$  manifolds and let  $f$  be a  $C^\infty$  map of  $M$  into  $\bar{M}$ . If  $f_*$  is non-singular (thus  $f_*$  has no kernel) at each point  $p$  of  $M$ , then  $f$  is called an *immersion* of  $M$  into  $\bar{M}$ . If in addition,  $f$  is univalent, then  $f$  is called an *imbedding* of  $M$  into  $\bar{M}$ . A subset  $M'$  of  $\bar{M}$  is called an *immersed submanifold* if there exists a manifold  $M$  and an immersion  $f : M \rightarrow \bar{M}$  such that  $f(M) = M'$ . (Thus an immersion is a “local imbedding with self-intersections”.) One can verify (problem 17) that if  $f : M \rightarrow \bar{M}$  is an imbedding and  $M' = f(M)$ , then by defining a differentiable structure of  $M'$  so  $f$  becomes a diffeomorphism,  $M'$  becomes a submanifold of  $\bar{M}$  (see [Hel12], p.23).

For examples of submanifolds see examples 5,6 and 7 at the end of section 1.1.

It is convenient to define a *base field* on a set  $A$  contained in an  $n$ -manifold to be a set of  $n$  vector fields that are independent at each point of  $A$ . When each field in a base field is  $C^\infty$ , then the base field is  $C^\infty$ . Since a set of coordinate fields is a  $C^\infty$  base field on the coordinate domain, we know  $C^\infty$  base fields always exist locally. A  $C^\infty$  base field does not necessarily exist over a whole manifold (consider the 2-sphere,  $S^2$ ); indeed, the manifold is called *parallelizable* if it admits a global  $C^\infty$  base field.

We now define a concept which we will often use. Let  $M$  be a submanifold of  $\bar{M}$  as described above. An  $\bar{M}$ -vector field  $Z$  that is

$C^\infty$  on  $M$  (or  $C^\infty$  on an open set  $A$  in  $M$ ) is a map that assigns to each  $p$  in  $M$  (or  $p$  in  $A$ ) a vector  $Z_p$  in  $\overline{M}_p$  such that if  $X_1, \dots, X_n$  is any  $C^\infty$  base field on a neighborhood  $\overline{U}$  of  $p$  and  $Z_m = \sum_{i=1}^n a_i(m)(X_i)_m$  for  $m$  in  $M \cap \overline{U}$  then the real valued functions  $a_i$  are  $C^\infty$  on  $M \cap U$  for all  $i$ . Notice  $Z_p$  is not necessarily tangent to  $M$ . Since the restriction to  $M$ , of a  $C^\infty$  function on  $\overline{M}$ , is a  $C^\infty$  function on  $M$ , it follows that if  $Z$  is  $C^\infty$  on  $\overline{M}$  then  $Z|_M$  is an  $\overline{M}$ -vector field that is  $C^\infty$  on  $M$ .

## Problems

(For problems 1 thru 9 see page 20)

10. (i) Let  $W_1, \dots, W_n$  be a  $C^\infty$  base field on an open set  $U$  in a manifold  $M$  and let  $X = \sum_{i=1}^n f_i W_i$  be a vector field on  $U$ . Show  $X$  is  $C^\infty$  on  $U$  iff the functions  $f_i$  are  $C^\infty$  on  $U$  for all  $i$ .
- (ii) If  $Y$  and  $Z$  are  $C^\infty$  fields on  $U$  show  $[Y, Z]$  is  $C^\infty$ , show that a coordinate field  $\frac{\partial}{\partial x_i}$  is  $C^\infty$  on its domain.
- (iii) If  $X_p$  is a given vector at  $p$  in  $M$  show there is a  $C^\infty$  field  $\overline{X}$  on a neighborhood of  $p$  with  $\overline{X} = X_p$ .
- (iv) Let  $x_1, \dots, x_n$  be a coordinate system with domain  $U$  and let  $A = \sum a_i \left( \frac{\partial}{\partial x_i} \right)$  and  $B = \sum b_j \left( \frac{\partial}{\partial x_j} \right)$  be  $C^\infty$  fields on  $U$  then find the representation of  $[A, B]$  in terms of the coordinate vector fields.
- (v) Show  $[fX, gY] = f(Xg)Y - g(Yf)X + fg[X, Y]$  where  $X$  and  $Y$  are  $C^\infty$  fields on  $U$  and  $f$  and  $g$  are in  $C^\infty(U, \mathbb{R})$ .
- (vi) Prove the **Jacobi Identity**.
11. (i) Let  $A, B$  and  $C$  be in  $C^\infty(\mathbb{R}^3, \mathbb{R})$  with  $B \neq 0$  anywhere. Let  $V = Ai + Bj + Ck$ ,  $X = -Bi + Aj$ , and  $Y = -Cj + Bk$  (advanced calculus notation). For  $p$  in  $\mathbb{R}^3$ , let  $P_p = \{Z \in (\mathbb{R}^3)_p : Z \cdot V_p = 0\}$ .



Show  $P_p$  is a two-dimensional space of vectors at each point by showing  $X_p$  and  $Y_p$  are a base for  $P_p$ .

- (ii) Show  $[X, Y]_p$  lies in  $P_p$  iff  $V_p \cdot (\text{curl } V)_p = 0$ .
  - (iii) Suppose there is a function  $f$  in  $C^\infty(\mathbb{R}^3, \mathbb{R})$  with  $\text{grad } f \neq 0$  such that  $P_p$  is the tangent plane to the constant surface of  $f$  thru  $p$ . show  $V_p \cdot (\text{curl } V)_p = 0$  (see section 9.1).
  - (iv) Instead of seeking surfaces that are orthogonal to  $V$  (as above), one could seek surfaces whose tangent plane contains  $V$  and then one has a “geometric quasi-linear partial differential equation of the first order”. Integral curves of  $V$  are called *characteristics* of the “equation”. One generates solution surfaces by taking a non-characteristic curve (an “initial value” curve) and considering the surface formed by characteristics thru the initial value curve. Show two solution surfaces must intersect along a characteristic. Show there are an infinite number of solution surfaces thru one characteristic. Can there be an initial value curve with no solution thru it?
12. (i) Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  by  $f(a, b) = (a^2 - 2b, 4a^3b^2)$  and let  $g : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  by  $g(u, v) = (u^2v + v^2, u - 2v^3, ve^u)$ . Compute a matrix for  $f_*$  at  $(1, 2)$  and  $g_*$  at any  $(u, v)$ .
- (ii) Find  $g_* \left( 4 \frac{\partial}{\partial x} - \frac{\partial}{\partial y} \right)_{(0,1)}$ .
- (iii) Find integral curves for the vector field  $X = yi + yj + 2k$  on  $\mathbb{R}^3$ .
- (iv) Find a coordinate system  $x_1, x_2, x_3$  on  $\mathbb{R}^3$  such that  $\frac{\partial}{\partial x_1} = 2i + 3j - k$  at all points.
13. Let  $X$  and  $Y$  be  $C^\infty$  fields about  $m$  in  $M$ . For small  $t \geq 0$  define the curve  $\sigma(t)$  as follows: go  $t$  parameter units on  $X$ 's integral curve thru  $m$  to  $p_1$ , go  $t$  units on  $Y$ 's integral curve thru  $p_1$  to  $p_2$ , go  $t$  units on  $(-X)$  curve thru  $p_2$  to  $p_3$ , go  $t$  units on  $(-Y)$

curve thru  $p_3$  to  $\sigma(t)$ . If  $\gamma(t) = \sigma(\sqrt{t})$  show  $T_\gamma(0) = [X, Y]_m$ . (Hint: use the lemma in section 9.1 and partial Taylor series.)

14. Let  $M$  and  $N$  be manifolds with  $M$  connected and let  $f$  and  $g$  be  $C^\infty$  maps of  $M$  into  $N$ .

(i) Show  $f_* \equiv 0$  iff  $f$  is a constant map.

(ii) If  $f(m) = g(m)$  at one  $m$  in  $M$  and  $f_* \equiv g_*$  at all points show  $f = g$ .

15. Let  $f$  be in  $C^\infty(M, \mathbb{R})$  and define the differential of  $f$ ,  $df$ , to be the linear map of  $\mathbf{T}_m M$  into  $\mathbb{R}$  where  $(df)_m(X_m) = X_m f$ . Show  $f_*(X_m) = [(df)_m(X)] \left( \frac{\partial}{\partial t} \right)$  where  $t$  is the identity coordinate function on  $\mathbb{R}$ . It is because of this case that in a general case the Jacobian  $f_*$  is often called the "differential of  $f$ ".

16. (i) Prove the Inverse Function Theorem (Theorem 1.3).

(ii) State and prove a version of the Implicit Function Theorem of advanced calculus in terms of the Jacobian map.

17. (i) Prove the last sentence in the third paragraph of section 1.6.

(ii) Show that the image of a regular ( $\sigma_* \neq 0$ ) univalent curve  $\sigma$  mapping an open interval into a manifold  $M$  is a one-dimensional submanifold of  $M$ .

(iii) Let  $X$  be a unit constant vector field on  $\mathbb{R}^2$  with irrational slope. Let  $T$  be the set of equivalence classes on  $\mathbb{R}^2$  where  $(a, b) \sim (c, d)$  iff  $a - c = n$  and  $b - d = m$  for integers  $m$  and  $n$ . Show  $T$  is a two-dimensional manifold (which is called the flat torus).

(iv) Show  $X$  induces a vector field on  $T$  such that the image of one integral curve of  $X$  defines a one-dimensional submanifold of  $T$  that is dense in  $T$ .

18. Let  $M_1$  and  $M_2$  be  $C^\infty$  manifolds. Let  $\pi_i : M_1 \times M_2 \rightarrow M_i$  by  $\pi_i(m_1, m_2) = m_i$  for  $i = 1, 2$ . Define a  $C^\infty$  structure on  $M_1 \times M_2$  so  $\pi_i$  are  $C^\infty$ . Show  $\mathbf{T}_{(m_1, m_2)} M_1 \times M_2$  is naturally isomorphic to  $\mathbf{T}_{m_1} M_1 \times \mathbf{T}_{m_2} M_2$ .
19. (i) Let  $M$  be a  $C^\infty$   $n$ -manifold. Let  $\mathbf{T}M = \{(m, X) : X \in \mathbf{T}_m M\}$ , and let  $\pi : \mathbf{T}M \rightarrow M$  by  $\pi(m, X) = m$ . If  $(\phi, U)$  is a coordinate pair on  $M$  with  $x_i = u_i \circ \phi$  let  $\bar{U} = \pi^{-1}(U)$ ,  $\bar{x}_i = x_i \circ \pi$ , and for  $(m, X)$  in  $\bar{U}$  let  $x_i(m, X) = a_i$  if  $X = \sum a_i \left( \frac{\partial}{\partial x_i} \right)$ . Let  $\bar{\phi} : \bar{U} \rightarrow \mathbb{R}^{2n}$  so  $u_i \circ \bar{\phi} = \bar{x}_i$  and  $u_{n+i} \circ \bar{\phi} = x_i$  for  $i = 1, \dots, n$ . Show the subatlas of pairs  $(\bar{\phi}, \bar{U})$  defines a  $C^\infty$  structure on  $\mathbf{T}M$  which is called the tangent bundle of  $M$ .
- (ii) If  $f$  is a  $C^\infty$  map of  $M$  into  $N$  show  $f_*$  induces a  $C^\infty$  map of  $\mathbf{T}M$  into  $\mathbf{T}N$ .
20. (i) Let  $G$  be a Lie group. If  $g \in G$  let  $L_g, R_g$ , and  $A_g$  denote the maps of  $G$  into  $G$  defined by  $L_g(h) = gh$ ,  $R_g(h) = hg$  and  $A_g(h) = ghg^{-1}$ . Show  $L_g, R_g$ , and  $A_g$  are  $C^\infty$ .
- (ii) A vector field  $X$  on  $G$  is left invariant if  $(L_g)_* X_g = X_{gh}$  for all  $g$  and  $h$ . Show a left invariant field is  $C^\infty$  and is completely determined by its value at the identity  $e$ .
- (iii) If  $X$  and  $Y$  are left invariant, show  $[X, Y]$  is left invariant.
- (iv) The set of left invariant vector fields on  $G$  forms an  $n$ -dimensional vector space called the Lie algebra of  $G$  which is denoted by  $\mathfrak{g}$ . Define a one-parameter subgroup of  $G$  to be the image of a  $C^\infty$  homomorphism of  $\mathbb{R}$  into  $G$ . Show there is a 1:1 correspondence between one-parameter subgroups and integral curves of left invariant vector fields thru  $e$ .
- (v) Show the map  $(g, h) \rightarrow gh^{-1}$  is  $C^\infty$  from  $G \times G$  into  $G$  iff the maps  $(g, h) \rightarrow gh$  and  $g \rightarrow g^{-1}$  are  $C^\infty$ .
21. (i) Let  $G = \mathbf{GL}(n, \mathbb{R})$  and for a matrix  $g \in G$  let  $u_{ij}(g) = g_{ij}$  (see example 3). Call  $u_{ij}$  the natural coordinate functions on

$G$ . Write  $u_{ij} \circ L_g$  as a linear combination of the natural coordinate functions.

- (ii) Let  $X_{ij}$  the unique left invariant field on  $G$  with  $X_{ij}(e) = \left(\frac{\partial}{\partial u_{ij}}\right)(e)$  where  $e$  is the identity element. Compute  $X_{ij}$  as a field on  $G$  in terms of the coordinate vector fields. Compute  $[X_{ij}, X_{rs}]$ .
- (iii) If  $A(t)$  is a  $C^\infty$  curve in  $G$  with  $A(0) = e$  and  $A(t)$  orthogonal for all  $t$  show  $\frac{dA}{dt} = \left(\frac{da_{ij}}{dt}\right)$  is a skew-symmetric matrix for  $t = 0$ .

22. (i) Let  $M$  be a  $C^\infty$   $n$ -manifold. Let

$$BM = \{(m; e_1, \dots, e_n) : m \in M \text{ and } e_1, \dots, e_n \text{ an ordered basis of } \mathbf{T}_m M\}.$$

Let  $\pi : BM \rightarrow M$  by  $\pi(m; e_1, \dots, e_n) = m$ . If  $(\phi, U)$  a coordinate pair on  $M$  with  $x_i = u_i \circ \phi$ , let  $(\bar{\phi}, \bar{U})$  be a coordinate pair on  $BM$  with  $\bar{U} = \pi^{-1}(U)$  and  $\bar{\phi} : \bar{U} \rightarrow \mathbb{R}^{n+n^2}$  by the coordinate functions  $\bar{x}_1, \dots, \bar{x}_n, x_{11}, x_{12}, \dots, x_{nn}$  where  $\bar{x}_i = x_i \circ \pi$  and if  $b = (m; e_1, \dots, e_n)$  then  $e_j = \sum_{i=1}^n x_{ij}(b) \left(\frac{\partial}{\partial x_i}\right)$ . Show the subatlas of pairs  $(\bar{\phi}, \bar{U})$  defines a  $C^\infty$  structure on  $BM$  which is called the bundle of bases over  $M$ .

(ii) For  $g$  in  $\mathbf{GL}(n, \mathbb{R})$  let  $R_g : BM \rightarrow BM$  by

$$R_g(b) \equiv bg \equiv \left(m; \sum_{i=1}^n g_{i1}e_i, \sum_{i=1}^n g_{i2}e_i, \dots, \sum_{i=1}^n g_{in}e_i\right)$$

if  $b = (m; e_1, \dots, e_n)$ . Show  $R_g$  is  $C^\infty$ .

- (iii) Let  $s_U : U \rightarrow BM$  by  $s_U(m) = \left(m; \left(\frac{\partial}{\partial x_1}\right)_m, \dots, \left(\frac{\partial}{\partial x_n}\right)_m\right)$  for  $m$  in  $U$ . Show  $s_U$  is  $C^\infty$  and  $\pi \circ s_U$  is the identity on  $U$ . The map  $s_U$  is called the coordinate section map over  $U$ .
- (iv) Let  $\hat{\phi} : U \times \mathbf{GL}(n, \mathbb{R}) \rightarrow \bar{U}$  by  $\hat{\phi}(m, g) = R_g \circ s_U(m) = s_U(m)g$ . Show  $\hat{\phi}$  is a diffeo. onto its image. The map  $\hat{\phi}$  is called a strip map.

- (v) If  $(\phi, U)$  and  $(\psi, V)$  are coordinate pairs on  $M$  define  $s_{UV} : U \cap V \rightarrow \mathbf{GL}(n, \mathbb{R})$  by  $s_{UV}(m) = g$  if  $s_U(m)g = s_V(m)$ . Show  $s_{UV}$  is  $C^\infty$ ; it is called a structural function for  $B(M)$ . Show  $(bg_1)g_2 = b(g_1g_2)$  which justifies the name right action for  $R_g$ .
- (vi) For fixed  $b$  in  $BM$  let  $f_b : \mathbf{GL}(n, \mathbb{R}) \rightarrow BM$  by  $f_b(g) = bg$ . Show  $f_b$  is  $C^\infty$ .
- (vii) Call the set  $F_m = \pi^{-1}(m)$  the (vertical) fiber over  $m$  in  $M$ . Show  $F_m$  is an  $n^2$ -submanifold of  $BM$  and  $f_b$  is a diffeo. of  $\mathbf{GL}(n, \mathbb{R})$  onto  $F_{\pi(b)}$ .
- (viii) If  $\pi(b) = \pi(c)$ , show  $f_c^{-1} \circ f_b$  is a left translation on  $\mathbf{GL}(n, \mathbb{R})$ .
- (ix) A tangent vector  $X$  on  $BM$  such that  $\pi_*(X) = 0$  is called a vertical vector. For  $b$  in  $BM$ , let  $E_{ij}(b) = (f_b)_* X_{ij}(e)$  define a vector  $E_{ij}(b)$  (see problem 21). Show  $E_{ij}$  is a global  $C^\infty$  vertical vector field on  $BM$ .
- (x) Compute  $[E_{ij}, E_{rs}]$ .



## 2. Hypersurfaces of $\mathbb{R}^n$

In a very real sense, this chapter and the next are too special, i.e., much of the theory belongs to an arbitrary submanifold of a “semi-Riemannian” manifold. We specialize because we can obtain many of the concepts and results of classical differential geometry quickly and easily. In so doing, we hope to develop the “geometric” intuition of the reader sufficiently to make later generalizations and definitions seem natural.

### 2.1 The Standard Connexion on $\mathbb{R}^n$

Recall in section 1.3 we shifted the classical notion of a vector from a “directed line segment” to an operator on functions, i.e., if  $X = a\vec{i} + b\vec{j} + c\vec{k}$  is a familiar vector on  $\mathbb{R}^3$  from advanced calculus, then we rewrite  $X = a\frac{\partial}{\partial x} + b\frac{\partial}{\partial y} + c\frac{\partial}{\partial z}$  so if  $f$  is a real valued  $C^\infty$  function on  $\mathbb{R}^3$ , then  $Xf$  is a derivative of  $f$  in the direction  $X$ ,

$$Xf = X \cdot \nabla f = a\frac{\partial f}{\partial x} + b\frac{\partial f}{\partial y} + c\frac{\partial f}{\partial z}.$$

Notice that  $X$  need not be a unit vector. When  $a, b$ , and  $c$  are  $C^\infty$  functions on  $\mathbb{R}^3$  themselves (possibly constant functions), then  $X$  is a  $C^\infty$  field and  $Xf$  is a  $C^\infty$  real valued function on  $\mathbb{R}^3$ ,

$$(Xf)(p) = X_p f = a(p)\frac{\partial f}{\partial x}(p) + b(p)\frac{\partial f}{\partial y}(p) + c(p)\frac{\partial f}{\partial z}(p)$$

Since both of the representations of a vector field  $X$  given above are awkward to write, let us simply write  $X = (a, b, c)$ , thus giving  $X$  by giving the coefficient functions (or constants)  $a, b$ , and  $c$  of the global base field  $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}$  on  $\mathbb{R}^3$ .

We now define the derivative of a vector field  $Y$  in a direction  $X$ . Let  $X$  be a vector at  $p$  in  $\mathbb{R}^n$  and let  $Y = (y_1, \dots, y_n)$  be a  $C^\infty$  field about  $p$ , thus each  $y_i$  is a  $C^\infty$  real valued function on the domain of  $Y$  which includes  $p$ . The covariant derivative of  $Y$  in the direction  $X$  is the vector  $\bar{D}_X Y = (X_p y_1, \dots, X_p y_n)$  as a vector at  $p$ . If  $X$  and  $Y$  are  $C^\infty$  fields with the same domain  $A$ , then  $\bar{D}_X Y$  is a  $C^\infty$  field with domain  $A$ .

For example take  $\mathbb{R}^3$ , let  $X = (a, b, c)$ , let  $Y = (xy^2 + 4z, y^2 - x, x + z^3)$ . and then

$$\begin{aligned}\bar{D}_X Y &= [X \cdot (y^2, 2xy, 4), X \cdot (-1, 2y, 0), X \cdot (1, 0, 3z^2)] \\ &= (ay^2 + 2xyb + 4c, -a + 2yb, a + 3z^2c)\end{aligned}$$

where  $a, b$ , and  $c$  may be functions or constants.

The properties of  $\bar{D}$  which we now list are one of the main analytic tools of these notes. Let  $X$  and  $W$  be vectors at  $p$  in  $\mathbb{R}^n$ , let  $Y$  and  $Z$  be  $C^\infty$  fields about  $p$ , and let  $f$  be a  $C^\infty$  real valued function about  $p$ . Then

- (1)  $\bar{D}_X(Y + Z) = \bar{D}_X Y + \bar{D}_X Z$
- (2)  $\bar{D}_{X+W}(Y) = \bar{D}_X Y + \bar{D}_W Y$
- (3)  $\bar{D}_{f(p)X} Y = f(p)\bar{D}_X Y$
- (4)  $\bar{D}_X(fY) = (Xf)Y_p + f(p)\bar{D}_X Y$

These follow directly from the definition of  $\bar{D}$ . It is important to notice  $\bar{D}_X Y$  can be computed once one knows  $Y$  along a curve  $\sigma$  that fits  $X$ , i.e., if  $\sigma(0) = p$  and  $T_\sigma(0) = T(0) = X_p$ . For let  $Y_{\sigma(t)} = (y_1(t), \dots, y_n(t))$  and then  $\bar{D}_X Y = \left( \frac{dy_1}{dt}(0), \dots, \frac{dy_n}{dt}(0) \right)$  since by



the chain rule,

$$\frac{dy_i}{dt}(0) = \sum_{j=1}^n \frac{\partial y_i}{\partial u_j}(p) \frac{du_j}{dt}(0) = X_p \cdot (\nabla y_i)_p$$

and  $T(0) = X_p$ . Thus if  $Y$  is an  $\mathbb{R}^n$  - vector field that is  $C^\infty$  on the curve  $\sigma$  with tangent  $T$ , then  $\bar{D}_T Y$  is a well-defined  $\mathbb{R}^n$ -vector field that is  $C^\infty$  on  $\sigma$ .

Using the operator  $\bar{D}$ , we can define parallel vector fields along a curve and geodesics. Let  $\sigma$  be a  $C^\infty$  curve (in  $\mathbb{R}^n$ ) with tangent  $T$  and let  $Y$  be an  $\mathbb{R}^n$ -vector field that is  $C^\infty$  on  $\sigma$ . The field  $Y$  is *parallel along  $\sigma$*  if  $\bar{D}_T Y = 0$  along  $\sigma$ . The curve  $\sigma$  is a *geodesic* if  $\bar{D}_T T = 0$ , i.e. if its tangent  $T$  is parallel along  $\sigma$ .

It is trivial to see these are the usual concepts of parallel fields and geodesics in  $\mathbb{R}^n$ ; for let  $\sigma(t) = (a_1(t), \dots, a_n(t))$  and  $Y_{\sigma(t)} = (y_1(t), \dots, y_n(t))$ . Then  $\bar{D}_T Y = \left( \frac{dy_1}{dt}, \dots, \frac{dy_n}{dt} \right) = 0$  iff each  $y_i(t)$  is a constant function of  $t$ , so  $Y$  is a "constant" vector field of  $\mathbb{R}^n$  evaluated on  $\sigma$ . The curve  $\sigma$  is a geodesic iff  $\bar{D}_T T = \left( \frac{d^2 a_1}{dt^2}, \dots, \frac{d^2 a_n}{dt^2} \right) = 0$ , and this implies  $a_i(t) = c_i t + d_i$  are linear functions of  $t$  so  $\sigma$  is a linear parameterization of a straight line.

Notice that the parameterization of a curve is important in the definition of a geodesic.

The generalization of the definition of covariant differentiation or a connexion on any  $C^\infty$  manifold  $M$  is clear, i.e. we merely demand the existence of an operator  $D$  which satisfies the above four properties (listed for  $\bar{D}$ ) and assigns to  $C^\infty$  vector fields  $X$  and  $Y$  with the domain  $A$ , a  $C^\infty$  field  $D_X Y$  on  $A$ . Notice there can be more than one connexion on a manifold. In the case of "semi-Riemannian" manifolds however there exists one connexion which fits the "semi-Riemannian" structure nicely, and in the case of  $\mathbb{R}^n$ ,  $\bar{D}$  is this nice connexion. In fact, we now explain how  $\bar{D}$  is "nice."

Henceforth, denote the usual dot product or inner product of vectors  $Y$  and  $Z$  tangent to  $\mathbb{R}^n$  by  $\langle Y, Z \rangle$ . Thus if  $Y = (y_1, \dots, y_n)$  and

$Z = (z_1, \dots, z_n)$ , then  $\langle Y, Z \rangle = \sum_{i=1}^n y_i z_i$ . If  $Y$  and  $Z$  are  $C^\infty$  fields with domain  $A$ , then  $\langle Y, Z \rangle$  is a  $C^\infty$  function with domain  $A$ . One checks easily that

$$(5) \quad \bar{D}_Y Z - \bar{D}_Z Y = [Y, Z] \text{ on } A, \text{ and}$$

$$(6) \quad X_p \langle Y, Z \rangle = \left\langle \bar{D}_X Y, Z \right\rangle_p + \left\langle Y, \bar{D}_X Z \right\rangle_p$$

for any vector  $X$  at  $p$  in  $A$ .

We now generalize and fix some terminology. A *Riemannian manifold* is a  $C^\infty$  manifold  $M$  on which one has singled out a  $C^\infty$  real valued, bilinear, symmetric, and positive definite function  $\langle -, - \rangle$  on ordered pairs of tangent vectors at each point. Thus if  $X, Y$  and  $Z$  are in  $\mathbf{T}_p M$ , then  $\langle X, Y \rangle$  is a real number and  $\langle -, - \rangle$  satisfies the following properties:

$$(a) \text{ (symmetric) } \langle X, Y \rangle = \langle Y, X \rangle,$$

$$(b) \text{ (bilinear)}$$

$$\langle X + Y, Z \rangle = \langle X, Z \rangle + \langle Y, Z \rangle$$

$$\langle aX, Y \rangle = a \langle X, Y \rangle \text{ for } a \text{ in } \mathbb{R}$$

$$(c) \quad \langle X, X \rangle > 0 \text{ for all } X \neq 0$$

$$(d) \quad (C^\infty) \text{ if } X \text{ and } Y \text{ are } C^\infty \text{ fields with domain } A \text{ then}$$

$$\langle X, Y \rangle_p = \langle X_p, Y_p \rangle \text{ is a } C^\infty \text{ function on } A$$

When (c) is replaced by

$$(c') \text{ (non-singular) } \langle X, Y \rangle = 0 \text{ for all } X \text{ implies } Y = 0,$$

then  $M$  is a *semi-Riemannian* (or *pseudo-Riemannian*) manifold. In either case, the functional  $\langle -, - \rangle$  is called the *inner product*, the

metric tensor, the Riemannian metric, or the infinitesimal metric of  $M$ . Notice the word “metric” in the preceding sentence is not referring to a metric function (distance function) in the topological sense. In Chapter 6, the connexion of the concepts is clarified. It is also customary to require a semi-Riemannian manifold to be Hausdorff; however, as far as the local differential geometry is concerned, this is irrelevant so the restriction is not enforced at this time.

If  $\mathbf{D}$  is a  $C^\infty$  connexion in a semi-Riemannian manifold  $M$ , then  $\mathbf{D}$  is a Riemannian connexion if it satisfies the above properties (5) and (6). In Chapter 6, the existence of Riemannian manifolds is discussed and the fundamental theorem asserting the existence and uniqueness of a Riemannian connexion is proved. In section 2.3 one sees that many hypersurfaces in  $\mathbb{R}^n$  ( $n \geq 3$ ) provide examples of Riemannian manifolds with a Riemannian connexion.

## 2.2 The Sphere Map and the Weingarten Map

An  $(n-1)$ -submanifold of an  $n$ -manifold is called a hypersurface. Throughout this section let  $M$  be a hypersurface of  $\mathbb{R}^n$ , let  $\bar{\mathbf{D}}$  be the natural connexion on  $\mathbb{R}^n$ , and assume  $N$  is a unit normal vector field that is  $C^\infty$  on  $M$ . Thus  $\langle N_p, N_p \rangle = 1$  and  $\langle N_p, X \rangle = 0$  for all  $p$  in  $M$  and  $X$  in  $\mathbf{T}_p M$ . Such an  $N$  always exists locally.

For any  $p$  in  $M$  and any vector  $X$  in  $\mathbf{T}_p M$ , define the linear map  $L: \mathbf{T}_p M \rightarrow \mathbf{T}_p M$  by

$$L(X) = \bar{\mathbf{D}}_X N. \tag{7}$$

The vector  $L(X)$  lies in  $\mathbf{T}_p M$ , since  $0 = X \langle N, N \rangle = 2 \langle L(X), N \rangle$  by property (6) for  $\bar{\mathbf{D}}$ . The map  $L$  is linear by properties (2) and (3). The map  $L$  is called the Weingarten map, and in the case of  $\mathbb{R}^n$ , it has a geometric interpretation as the Jacobian of the sphere map (Gauss map) which we now explain.

Let  $N = (a_1, \dots, a_n)$ , so the  $a_i$  are real valued  $C^\infty$  functions on  $M$  and  $\sum_i (a_i)^2 = 1$ . Then the map  $\eta: M \rightarrow S^{n-1}$  defined by  $\eta(p) =$

$(a_1(p), \dots, a_n(p))$  in  $\mathbb{R}^n$ , is a  $C^\infty$  map of  $M$  into the unit  $(n-1)$ -sphere  $S^{n-1}$ , and  $\eta$  is called the *sphere map* (or *Gauss map*). If  $X \in \mathbf{T}_p M$  and  $\sigma(t)$  is a curve fitting  $X$  (so  $\sigma(0) = p$  and  $T_\sigma(0) = X$ ), then  $\eta \circ \sigma(t) = (a_1 \circ \sigma(t), \dots, a_n \circ \sigma(t))$  and

$$\begin{aligned} \eta_*(X) &= T_{\eta \circ \sigma}(0) = \left( \frac{d(a_1 \circ \sigma)}{dt}(0), \dots, \frac{d(a_n \circ \sigma)}{dt}(0) \right) \\ &= (Xa_1, \dots, Xa_n) = \bar{D}_X N = L(X). \end{aligned}$$

The map  $L$  is  $C^\infty$  on  $M$  in the sense that if  $X$  is  $C^\infty$  on the subset  $A$  of  $M$  then  $L(X) = (Xa_1, \dots, Xa_n)$  is also  $C^\infty$  on  $A$  since each  $a_1$  is  $C^\infty$  on  $M$ .




Figure 2.1: The Weingarten Map (derivative of normal)

Our next objective is to show  $L$  is *self-adjoint* or *symmetric*, i.e., if  $X, Y$  are in  $\mathbf{T}_p M$  then  $\langle L(X), Y \rangle = \langle X, L(Y) \rangle$ .

To do this, let  $Z$  be a  $C^\infty$  field defined on a special coordi-

---

## Section 2.2: The Sphere Map and the Weingarten Map

nate neighbourhood  $U$  of  $p$  and let  $\bar{U}$  be the associated coordinate neighbourhood of  $p$  in  $\mathbb{R}^n$  with coordinate functions  $\bar{x}_1, \dots, \bar{x}_n$ . Then

$Z = \sum_{i=1}^{n-1} g_i \left( \frac{\partial}{\partial \bar{x}_i} \right)$ , where  $g_i$  are  $C^\infty$  real valued functions on  $U$ . We

want to extend  $Z$  to a  $C^\infty$  field  $\bar{Z}$  on  $\bar{U}$ , i.e., we want  $\bar{Z}$  so that  $\bar{Z}_p = Z_p$  for  $p$  in  $U$ . Let us assume the coordinate map  $\bar{\phi}$  maps  $\bar{U}$

onto a ball,  $B$ , about the origin in  $\mathbb{R}^n$ , i.e.,  $\bar{x}_i(p) = 0 = u_i \circ \bar{\phi}(p)$  for all  $i$ . Then if  $(t_1, \dots, t_n)$  is in  $B$ , let  $\pi: (t_1, \dots, t_n) \rightarrow (t_1, \dots, t_{n-1}, 0)$ . This

map  $\pi$  (which is  $C^\infty$ ) induces a  $C^\infty$  map  $\sigma: \bar{U} \rightarrow U$  by  $\sigma = \bar{\phi}^{-1} \circ \pi \circ \bar{\phi}$ .

Letting  $\bar{Z} = \sum_{i=1}^{n-1} (g_i \circ \sigma) \left( \frac{\partial}{\partial \bar{x}_i} \right)$ , the field  $\bar{Z}$  is a  $C^\infty$  extension of  $Z$  to  $\bar{U}$ .

Actually the above process allows us to extend an  $\mathbb{R}^n$ -field  $Z$  that is  $C^\infty$  on  $U$  to a  $C^\infty$  field  $\bar{Z}$  on  $\bar{U}$ .

Having the existence of such extensions we prove a proposition.

**Proposition 2.1.** Let  $\bar{U}$  and  $U$  be special neighborhoods of  $p$  as above and let  $\bar{Z}$  and  $Z$  be  $C^\infty$  fields on  $\bar{U}$  and  $U$ , respectively. Then  $\bar{Z}$  is an extension of  $Z$  (i.e.,  $\bar{Z}_p = i_*(Z_p)$  for  $p$  in  $U$ ) iff  $(\bar{Z}f)|_U = Z(f|_U)$  for all  $f$  in  $C^\infty(\bar{U}, \mathbb{R})$ . If  $\bar{X}$  and  $\bar{Y}$  are  $C^\infty$  extensions to  $\bar{U}$  of  $C^\infty$  fields  $X$  and  $Y$  on  $U$ , then  $[\bar{X}, \bar{Y}]$  is a  $C^\infty$  extension of  $[X, Y]$ .

*Proof.* If  $\bar{Z}_p = i_*(Z_p)$  for  $p$  in  $U$ , where  $i: M \rightarrow \mathbb{R}^n$  is the inclusion, then for  $f$  in  $C^\infty(\bar{U}, \mathbb{R})$ ,  $(\bar{Z}f)(p) = \bar{Z}_p f = (i_*(Z_p))f = Z_p(f \circ i) = Z(f|_U)(p)$ . Conversely, if the two extreme terms are equal, then the second equality follows.

For the rest of the proposition consider for  $p$  in  $U$

$$\begin{aligned} [\bar{X}, \bar{Y}]_p f &= \bar{X}_p(\bar{Y}f) - \bar{Y}_p(\bar{X}f) = X_p((\bar{Y}f)|_U) - Y_p((\bar{X}f)|_U) \\ &= X_p(Y(f|_U)) - Y_p(X(f|_U)) = [X, Y]_p(f|_U), \end{aligned}$$

thus  $[\bar{X}, \bar{Y}]$  is an extension of  $[X, Y]$ . □

**Theorem 2.2.** The Weingarten map is self-adjoint.

*Proof.* Take  $X$  and  $Y$  in  $\mathbf{T}_p M$ , imbed  $X$  and  $Y$  in  $C^\infty$  fields on a special neighborhood  $U$  of  $p$ , and extend  $X$  and  $Y$  to  $C^\infty$  fields  $\bar{X}$

and  $\bar{Y}$  on  $\bar{U}$  as above. Then

$$\begin{aligned}
 \langle LX, Y \rangle - \langle X, LY \rangle &= \langle \bar{D}_X N, Y \rangle - \langle X, \bar{D}_Y N \rangle \\
 &= \langle \bar{D}_{\bar{X}} \bar{N}, \bar{Y} \rangle_p - \langle \bar{X}, \bar{D}_{\bar{Y}} \bar{N} \rangle_p \\
 &= \bar{X}_p \langle \bar{N}, \bar{Y} \rangle - \langle \bar{N}, \bar{D}_{\bar{X}} \bar{Y} \rangle_p - \bar{Y}_p \langle \bar{N}, \bar{X} \rangle + \langle \bar{N}, \bar{D}_{\bar{Y}} \bar{X} \rangle_p \\
 &= \langle \bar{D}_{\bar{Y}} \bar{X} - \bar{D}_{\bar{X}} \bar{Y}, \bar{N} \rangle_p \\
 &= \langle [\bar{Y}, \bar{X}], \bar{N} \rangle_p = \langle [Y, X]_p, N_p \rangle = 0,
 \end{aligned}$$

since  $\bar{X}_p \langle \bar{N}, \bar{Y} \rangle = X_p \langle N, Y \rangle = 0 = Y_p \langle N, X \rangle$ . □

The *fundamental forms on  $M$*  can now be defined in terms of  $L$  and the inner product. If  $X$  and  $Y$  are in  $\mathbf{T}_p M$ , then

$$\text{I}(X, Y) = \langle X, Y \rangle$$

$$\text{II}(X, Y) = \langle L(X), Y \rangle$$

$$\text{III}(X, Y) = \langle L^2(X), Y \rangle$$

$$\text{IV}(X, Y) = \langle L^3(X), Y \rangle$$

etc., and these forms are called the *first, second, third, etc. fundamental forms on  $M$* . Notice  $M$  is a Riemannian manifold with metric tensor defined by the first fundamental form. Since the inner product is symmetric and  $L$  is self-adjoint, the fundamental forms are all symmetric bilinear functions on  $\mathbf{T}_p M \times \mathbf{T}_p M$  for all  $p$  in  $M$ . These forms are  $C^\infty$  in the sense that if  $X$  and  $Y$  are  $C^\infty$  fields with domain  $A$ , then  $\langle L^k(X), Y \rangle_p = \langle L^k(X_p), Y_p \rangle$  is a  $C^\infty$  real valued function on  $A$ . The first three forms have a direct interpretation geometrically since  $L$  represents the Jacobian of the sphere map.

The algebraic invariants of the linear map  $L$  at each point now define the imbedded geometric invariants of the submanifold  $M$  at each point. Thus the determinant of  $L$  at  $p$  is the total curvature (Gauss curvature)  $K(p)$  of  $M$  at  $p$ , the trace of  $L$  at  $p$  is the mean

curvature  $H(p)$ , etc. The eigenvalues of  $L$  are the *principal curvatures* and the eigenvectors of  $L$  are the *directions of curvature* or *principal vectors*. Since  $L$  is self-adjoint there are always  $(n - 1)$  independent directions of curvature. If  $L$  is a multiple of the identity map on  $\mathbf{T}_pM$ , then  $p$  is an *umbilic point* of  $M$ . If  $L = 0$  at  $p$  we call  $p$  a *flat point* of  $M$ . Non-zero vectors  $X$  and  $Y$  in  $\mathbf{T}_pM$  are *conjugate* if  $\langle LX, Y \rangle = 0$ . A vector  $X$  (not zero) is *asymptotic* if it is self-conjugate, i.e., if  $\langle LX, X \rangle = 0$ . A curve in  $M$  is a *line of curvature* if its tangent is a principal vector at each of its points.

The following facts come immediately from these definitions. An asymptotic direction  $X$  is a direction of curvature iff  $LX = 0$  iff  $X$  is conjugate to all vectors. Conjugate directions always exist since if  $LX \neq 0$  then there exists a  $Y$  which is orthogonal to  $LX$ . If the second fundamental form  $\langle LX, Y \rangle$  is positive or negative definite no asymptote directions exist. If  $X$  and  $Y$  are two directions of curvature belonging to unequal eigenvalues, then  $X$  is orthogonal to  $Y$ . The proof of this is standard algebra, i.e.,

$$0 = \langle LX, Y \rangle - \langle X, LY \rangle = \langle k_1X, Y \rangle - \langle X, k_2Y \rangle = (k_1 - k_2) \langle X, Y \rangle,$$

so  $k_1 \neq k_2$  implies  $\langle X, Y \rangle = 0$ . If  $X$  and  $Y$  are non-zero independent vectors with  $LX = kX$  and  $LY = -kY$ , then the vectors  $X + Y$  and  $X - Y$  are orthogonal asymptotic directions spanning the same subspace as  $X$  and  $Y$ . Finally one notices that  $L$  must satisfy its characteristic polynomial, which will also give a relation between the fundamental forms, i.e., if  $n = 3$ , then  $L^2 - HL + K(\text{identity}) = 0$  and  $\text{III} - H\text{II} + K\text{I} = 0$ .

When  $X$  is a principal vector, the Weingarten map says  $\bar{D}_X N = kX$ , where  $k$  is a principal curvature, and this equality is classically called the *formula of Rodrigues*.

Another classical concept is the *Dupin indicatrix* at each  $p$  in  $M$  which is the subset of  $\mathbf{T}_pM$  consisting of all vectors in  $X$  such that  $\langle L(X), X \rangle = \pm 1$ .

Let  $n = 3$  and let  $X$  and  $Y$  be unit orthogonal principal vectors in  $\mathbf{T}_pM$  with  $LX = kX$  and  $LY = hY$ . If  $Z = aX + bY$ , then  $\langle LZ, Z \rangle = ka^2 + hb^2$ . Thus the indicatrix is the curve (or curves) in  $\mathbf{T}_pM$  such that  $ka^2 + hb^2 = \pm 1$ . Consider the three cases:

1. If  $K(p) > 0$ , then  $h$  and  $k$  have the same sign (for  $K = hk = \det L$ ) so suppose they are positive. The indicatrix is then an ellipse determined by  $ka^2 + hb^2 = 1$ , and  $p$  is an *elliptic point*.
2. If  $K(p) < 0$ , then  $h$  and  $k$  have opposite signs, the indicatrix is two hyperbolas, and  $p$  is a *hyperbolic point*.
3. If  $K(p) = 0$ , say  $k = 0, h > 0$ , then  $b = \pm 1/\sqrt{h}$  gives two straight lines parallel to the  $X$  vector, and  $p$  is a *parabolic point*. (When  $k = h = 0$ ,  $p$  is an *umbilic* and a *flat point*.)

There is a geometric interpretation of the indicatrix as an approximation to the intersection of the surface with a plane which is parallel and close to the tangent plane; for details see [Str61] (p.84).

### 2.3 The Gauss Equation

As in the last section, let  $M$  be a hypersurface of  $\mathbb{R}^n$ , let  $\bar{D}$  be the natural connexion on  $\mathbb{R}^n$ , let  $N$  be a unit normal field that is  $C^\infty$  on  $M$ , and let  $L(X) = \bar{D}_X N$  for  $X$  tangent to  $M$ . Let  $U$  and  $\bar{U}$  be special coordinate neighborhoods of a point  $p$  in  $M$  and  $\mathbb{R}^n$  respectively, and let  $\bar{Z}$  be a  $C^\infty$  extension to  $\bar{U}$  of a  $C^\infty$  field  $Z$  on  $U$  as usual.

If  $Y$  is a  $C^\infty$  field about  $p$  in  $M$ , and  $X$  in  $\mathbf{T}_pM$ , define  $\mathbf{D}_X Y$  by

$$\mathbf{D}_X Y = \bar{D}_X Y - \langle LX, Y \rangle N. \quad (8)$$

This is the Gauss equation. First notice  $\mathbf{D}_X Y$  is in  $\mathbf{T}_pM$  for

$$\langle \mathbf{D}_X Y, N \rangle = \langle \bar{D}_X Y, N \rangle + \langle \bar{D}_X N, Y \rangle = X \langle Y, N \rangle = 0.$$



since  $\langle Y, N \rangle = 0$  in a neighborhood of  $p$ . Next notice if  $X, Y$  are  $C^\infty$  on  $U$ , then  $\bar{\mathbf{D}}_X Y = \bar{\mathbf{D}}_{\bar{X}} \bar{Y} \Big|_U$  and  $\langle LX, Y \rangle N$  are both  $C^\infty$  on  $U$ , so  $\mathbf{D}_X Y$  is  $C^\infty$  on  $U$ ; because of this, we say  $\mathbf{D}$  is  $C^\infty$ .

Thus  $\mathbf{D}$  becomes a candidate to define a covariant differentiation or a connexion on the submanifold  $M$  which is defined very simply from the natural connexion on  $\mathbb{R}^n$  by decomposing  $\bar{\mathbf{D}}_X Y$  into its unique tangent and normal components relative to the tangent space of  $M$ . One must now check if the properties (1), (2), (3) and (4) are satisfied for  $\mathbf{D}$ , and indeed they are, since they are satisfied for  $\bar{\mathbf{D}}$  and the second fundamental form is bilinear. The properties (5) and (6) are also valid for  $\mathbf{D}$ , so  $\mathbf{D}$  is the natural Riemannian connection associated with the induced metric (first fundamental form) on  $M$  (see Chapter 6). The proof of the first four properties is left to the reader, but we now show (5) and (6). Let  $Y$  and  $Z$  be fields on a neighborhood  $U$  about  $p$ , let  $\bar{Y}$  and  $\bar{Z}$  be extensions to  $\bar{U}$ , and let  $X$  be in  $\mathbf{T}_p M$ . Then

$$\begin{aligned} (\mathbf{D}_Y Z - \mathbf{D}_Z Y)_p &= (\bar{\mathbf{D}}_Y Z - \bar{\mathbf{D}}_Z Y)_p = (\bar{\mathbf{D}}_{\bar{Y}} \bar{Z} - \bar{\mathbf{D}}_{\bar{Z}} \bar{Y})_p \\ &= [\bar{Y}, \bar{Z}]_p = [Y, Z]_p \end{aligned}$$

and

$$\begin{aligned} X \langle Y, Z \rangle &= X \langle \bar{Y}, \bar{Z} \rangle = \langle \bar{\mathbf{D}}_X \bar{Y}, \bar{Z} \rangle + \langle \bar{Y}, \bar{\mathbf{D}}_X \bar{Z} \rangle \\ &= \langle \mathbf{D}_X Y, Z_p \rangle + \langle Y_p, \mathbf{D}_X Z \rangle. \end{aligned}$$

Thus the natural metric tensor and connexion on  $\mathbb{R}^n$  induce a Riemannian metric and Riemannian connexion on the hypersurface  $M$ .

Since the Gauss equation induces a connexion  $\mathbf{D}$  on  $M$ , one can define parallel vector fields along a curve and geodesics exactly as in Section 2.1. If  $\sigma$  is a  $C^\infty$  curve in  $M$  with tangent  $T$  and  $Y$  is a  $C^\infty$  field along  $\sigma$ , then  $Y$  is parallel along  $\sigma$  if  $\mathbf{D}_T Y = 0$  along  $\sigma$ . The curve  $\sigma$  is a geodesic if  $\mathbf{D}_T T = 0$  along  $\sigma$ .

Application of the Gauss equation to the tangent field along a curve gives two results immediately.




Figure 2.2: The Decomposition of  $\bar{D}_X Y$

**Theorem 2.3.** Let  $M$  be a hypersurface in  $\mathbb{R}^n$ . A curve in  $M$  is a geodesic in  $\mathbb{R}^n$  iff it is an asymptotic geodesic in  $M$ . A curve in  $M$ , which is not a geodesic in  $\mathbb{R}^n$ , is a geodesic in  $M$  iff  $\bar{D}_T T$  is normal to  $M$  along the curve (whose tangent is  $T$ ).

*Proof.* Let  $g$  be a curve in  $M$  with tangent  $T$ . The Gauss equation implies  $\bar{D}_T T = D_T T - \langle LT, T \rangle N$ . Thus  $\bar{D}_T T = 0$  iff  $D_T T = 0$  and  $\langle LT, T \rangle = 0$ . And  $D_T T = 0$  iff  $\bar{D}_T T$  is normal to  $M$ .  $\square$

**Corollary 2.4.** If  $M_1$  and  $M_2$  are two hypersurfaces of  $\mathbb{R}^n$  and  $g$  is a geodesic on both hypersurfaces that is not a geodesic in  $\mathbb{R}^n$ , on any parameter interval, then  $M_1$  and  $M_2$  are tangent along  $g$  (i.e., their tangent spaces coincide along  $g$ ).

*Proof.* Let  $T$  be the tangent to  $g$ . Since  $\bar{D}_T T \neq 0$  on any parameter interval, the normals to  $M_1$  and  $M_2$  determine the same subspace

## Section 2.4: The Gauss Curvature and Codazzi-Mainardi Equations

on a dense set of the parameter domain. Hence  $M_1$  and  $M_2$  are tangent along  $g$ .  $\square$

### 2.4 The Gauss Curvature and Codazzi-Mainardi Equations

Let  $M, N, L, D$  and  $\bar{D}$  be as is in the previous two sections. Our current goal is the “*theorema egregium*” of Gauss. This will show that the “curvature” is independent of the embedding, and motivate the definition of Riemannian curvature and curvature of a general connection. Let  $X, Y$  and  $Z$  be  $C^\infty$  fields on an open set  $A \in M$ . Notice that

$$\begin{aligned} & \bar{D}_X(\bar{D}_Y Z) - \bar{D}_Y(\bar{D}_X Z) - \bar{D}_{[X,Y]} Z = \\ & (XYz_1, \dots, XYz_n) - (YXz_1, \dots, YXz_n) - ([X, Y]z_1, \dots, [X, Y]z_n) \end{aligned}$$

where  $Z = (z_1, \dots, z_n)$  and  $z_i$  are  $C^\infty$  real valued functions on  $A$ . This fact will later verify that the “curvature of  $\mathbb{R}^n$  is zero.” By applying the Gauss equation and decomposing the above expression into tangent and normal parts, one obtains the Gauss curvature [9](#) and the Codazzi-Mainardi equations [10](#), respectively. Thus,

$$\begin{aligned} 0 &= \bar{D}_X(\mathbf{D}_Y Z - \langle LY, Z \rangle N) - \bar{D}_Y(\mathbf{D}_X Z - \langle LX, Z \rangle N) - \bar{D}_{[X,Y]} Z \\ &= \mathbf{D}_X \mathbf{D}_Y Z - \langle LX, \mathbf{D}_Y Z \rangle N - X(\langle LY, Z \rangle) N - \langle LY, Z \rangle L(X) \\ &\quad - \mathbf{D}_Y \mathbf{D}_X Z + \langle LY, \mathbf{D}_X Z \rangle N + Y(\langle LX, Z \rangle) N + \langle LX, Z \rangle L(Y) \\ &\quad - \mathbf{D}_{[X,Y]} Z + \langle L([X, Y]), Z \rangle N. \end{aligned}$$

Equating tangent and normal parts to zero gives

$$\mathbf{D}_X \mathbf{D}_Y Z - \mathbf{D}_Y \mathbf{D}_X Z - \mathbf{D}_{[X,Y]} Z = \langle LY, Z \rangle L(X) - \langle LX, Z \rangle L(Y) \quad (9)$$

and

$$\langle \mathbf{D}_X L(Y) - \mathbf{D}_Y L(X) - L([X, Y]), Z \rangle = 0$$

for all  $Z$ , so

$$\mathbf{D}_X L(Y) - \mathbf{D}_Y L(X) - L([X, Y]) = 0 \quad (10)$$

Define

$$R(X, Y)Z = \mathbf{D}_X \mathbf{D}_Y Z - \mathbf{D}_Y \mathbf{D}_X Z - \mathbf{D}_{[X, Y]} Z,$$

and notice 9 implies  $R(X, Y)Z$  does not depend on the field nature of  $X, Y$ , and  $Z$ . Thus  $R(X, Y)Z$  is a vector at  $p$  in  $A$  which depends only on  $X_p, Y_p$  and  $Z_p$  since these vectors are all that is needed to compute the left side of 9. Thus  $R(X_p, Y_p)$  define a linear transformation on  $\mathbf{T}_p M$  called the curvature of  $X_p$  and  $Y_p$ . The justification of this definition is the following theorem, Gauss' "theorema egregium."

**Theorem 2.5.** Let  $n = 3$  and let  $X$  and  $Y$  be an orthonormal base of  $\mathbf{T}_p M$ . Then the total curvature  $K(p) = \det L_p = \langle R(X, Y)Y, X \rangle$

*Proof.* Using the Gauss curvature equation 9,

$$\langle R(X, Y)Y, X \rangle = \langle LY, Y \rangle \langle LX, X \rangle - \langle LX, Y \rangle \langle LY, X \rangle = \det L = K(p).$$

□

The above theorem is significant because the term  $\langle R(X, Y)Y, X \rangle$  depends only on the metric  $\langle -, - \rangle$  and the connexion  $\mathbf{D}$ , and it is completely independent of the normal  $N$  or the map  $L$ . Thus the total curvature  $K(p) = \langle R(X, Y)Y, X \rangle$  is an "intrinsic" invariant that is independent of the "imbedding" (i.e., of  $N$  and  $L$ ). The theorem is generalized in Chapter 6.

## 2.5 Examples

See Figure 2.3 for sketches of (1), (2), (3).

1. Let  $M$  be an  $(n - 1)$ -dimensional hyperplane in  $\mathbb{R}^n$ , i.e., let  $N = (a_1, \dots, a_n)$  determine a constant unit normal field on  $M$ . Then  $L(X) = \mathbf{D}_X N = (Xa_1, \dots, Xa_n) = 0$  for all  $X$  at all points

of  $M$ , i.e.,  $L \equiv 0$  on all of  $M$ . Thus  $M$  consists entirely of flat (umbilic) points, the total curvature  $K$  and mean curvature  $H$  (and all others) are identically zero. All the fundamental forms, except the first, are completely singular. Every vector is asymptotic and a direction of curvature, and all principal curvatures are zero.

2. Let  $M$  be  $S$ , the unit sphere about the origin in  $\mathbb{R}^n$ , and let  $N$  be the outer normal on  $S$ , i.e., if  $p = (a_1, \dots, a_n)$  then  $N(p) = (a_1, \dots, a_n)$ . Thus the sphere map  $\eta$  is the identity map,  $\eta_*$  is also the identity map, and hence  $L(X) = X$  for all  $X$ . Thus  $K \equiv 1$ ,  $H \equiv (n - 1)$  on  $S$ . All the fundamental forms are equal to the first fundamental form, all points are umbilic, and all principal curvatures are unity. Every vector is a direction of curvature and there are no asymptotic directions.
3. Let  $M$  be the cylinder  $C = \{(t_1, \dots, t_n) \in \mathbb{R}^n : \sum_1^{n-1} (t_i)^2 = 1\}$  with  $N =$  the "outer" normal. For  $X = e_n = (0, 0, \dots, 0, 1)$  we have  $LX = 0$ , and for  $X$  orthogonal to  $e_n$  and tangent to  $C$  we have  $LX = X$ . Hence  $K \equiv 0$ ,  $H \equiv (n - 1)$ , all principal curvature are unity except one which is zero, etc.




Figure 2.3: Pieces of Examples (1), (2), (3)

4. Next let  $M$  be an open piece of a surface of revolution about the  $z = e_3$ -axis in  $\mathbb{R}^3$  (vaguely:  $M$  is obtained by revolving a

$C^\infty$  plane curve about an axis in the plane). Let  $P$  be a plane containing the  $z$ -axis and take  $m$  in  $M \cap P$  (and let us consider the point  $m$  not on the  $z$ -axis at first).

Since the normal  $N$  lines in  $P$ , the vector  $\bar{D}_X N = L(X)$  lies in  $P$  and is tangent to  $M$  so  $L(X) = kX$  and  $X$  is a direction of curvature, where  $X$  is the unit tangent to a meridian curve. From the remarks preceding the examples there is a direction of curvature orthogonal to  $X$ , so the unit vector  $Y$  tangent to the parallel curves is a direction of curvature. The vector field  $\bar{D}_X X$  is zero or orthogonal to  $X$  and must lie in the plane  $P$ , hence  $\bar{D}_X X = \pm \bar{k}_1 N$ , so  $D_X X = 0$ , and we see the meridians are geodesics. If the parallel curve through  $m$  is a geodesic, then  $\bar{D}_Y Y$  is normal to  $M$  and not zero, since these curves are not geodesics in  $\mathbb{R}^3$ . But  $\bar{D}_Y Y$  is orthogonal to  $e_3$ , the  $z$  direction, hence a parallel curve is a geodesic on  $M$  iff the normal  $N$  along the parallel curve is horizontal (i.e., orthogonal to the  $z$ -axis). If  $m$  is a point on the  $z$ -axis, then every direction  $X$  is tangent to a meridian and hence is a direction of curvature, so  $m$  is umbilic and  $K(m) \geq 0$ .

5. Let us apply the analysis of example 4 to a torus, i.e., let  $M$  be obtained by rotating a circle  $C$  in the  $x, z$ -plane about the  $z$ -axis here we assume the circle does not intersect the  $z$ -axis.

Then the meridians generated by  $C$  are geodesic, as is the minimum length parallel  $A$  and maximum length parallel  $B$ . Along  $B$ ,  $M$  has positive curvature, along  $A$  the curvature is negative, and the curvature is zero on the extreme top and bottom curves  $E$  and  $F$  where  $N$  is constant. Indeed, if  $r_1$  is the radius of  $A$  and  $r_2$  is the radius of  $B$ , then  $a = \frac{r_2 - r_1}{2}$  is the




Figure 2.4: The Torus

radius of  $C$  and

$$K = \frac{1}{ar_2} = \frac{2}{r_2(r_2 - r_1)} \quad \text{on } B,$$

$$K = -\frac{1}{ar_1} = \frac{-2}{r_1(r_2 - r_1)} \quad \text{on } A,$$

These expressions can be derived as follows. Let  $X$  be the unit tangent field to a circle of radius  $r$  about the origin in  $\mathbb{R}^2$  (see Figure 2.4) so  $f(t) = (r \cos(t/r), r \sin(t/r))$  parameterizes the circle to fit  $x$ . Then evaluating a unit outer normal  $N$  on  $f(t)$  gives  $N \circ f(t) = (\cos(t/r), \sin(t/r))$ . Hence,

$$\frac{d}{dt}(N \circ f(t)) = \frac{1}{r}X,$$

or if the circle lies on the surface then we see  $\bar{D}_X N = L(X) = \frac{1}{r}X$ . Now apply this to the circles on the torus.

6. We discuss ruled surfaces and developable surfaces briefly. A ruled surface is a two-dimensional submanifold  $M$  of  $\mathbb{R}^3$  such that through each point  $p$  in  $M$  there passes a segment of a straight line (the generator through  $p$ ) which lies in  $M$ . When

the normal field is a parallel field in  $\mathbb{R}^3$  along the generators, (thus the tangent plane is constant along generators), then the ruled surface is a *developable surface*. Notice we only consider the  $C^\infty$  case, although the above definitions can be generalized.

Let  $M$  be a ruled surface, and let  $X$  be a  $C^\infty$  unit vector tangent to the generator at each point of  $M$ . The generators are geodesics in  $\mathbb{R}^3$ , so  $\bar{D}_X X = 0$ , and hence, from the Gauss equation,  $D_X X = 0$  and  $\langle LX, X \rangle = 0$  (so generators are asymptotic lines). Let  $Y$  be a unit vector field orthogonal to  $X$  in the neighborhood of a point  $p$ , then  $K = \langle LX, X \rangle \langle LY, Y \rangle - \langle LX, Y \rangle^2 = -\langle LX, Y \rangle^2 \leq 0$  in this neighborhood. Thus a ruled surface has non-positive curvature. For a developable surface,  $0 = \bar{D}_X N = LX$  so  $K \equiv 0$ . A theorem due to Massey (see Chapter 3) states a closed connected surface is developable iff  $K \equiv 0$ .

We study the neighborhood of a point  $p$  in a ruled surface  $M$ . Let  $f(t)$  be the  $C^\infty$  curve through  $p$  which is parameterized by arc-length and is orthogonal to the generators at each point. Let  $T$  be the tangent to  $f$  (say  $T = Y$  along  $f$ ), and let  $f(0) = p$ . Then the map  $(t, s) \mapsto f(t) + sX(t)$  gives a coordinate system from a neighborhood of  $(0, 0)$  in  $\mathbb{R}^2$  to a neighborhood of  $p$  in  $M$ .

Let  $N$  be a local unit normal for this coordinate neighborhood. The unit fields  $X, T, N$  give an orthonormal frame along  $f$ , and we next obtain the Frenet formulas for this frame. On  $f$  we have

$$1 = \langle X, X \rangle = \langle T, T \rangle = \langle N, N \rangle \text{ so } 0 = T \langle X, X \rangle = 2 \langle \bar{D}_T X, X \rangle$$

implies  $\bar{D}_T X$  normal to  $X$ . Similarly,  $\bar{D}_T N$  normal to  $N$  and  $\bar{D}_T T$  normal to  $T$ . Thus we define functions  $a(t), b(t), c(t)$  by

$$\begin{aligned} \bar{D}_T T &= aX + bN \\ \bar{D}_T X &= -aT + cN \\ \bar{D}_T N &= -bT - cX \end{aligned}$$






Figure 2.5: Ruled Surface

where  $a = \langle \bar{D}_T T, X \rangle = T \langle T, X \rangle - \langle T, \bar{D}_T X \rangle = -\langle T, \bar{D}_T X \rangle$ , etc. Holding  $s$  constant, we get a curve  $f_s(t) = f(t) + sX(t)$  on  $M$  with tangent

$$A = T + s\bar{D}_T X = (1 - as)T + scN$$

(note that  $T(t)$  and  $N(t)$  are vectors at  $f(t)$  which are rigidly translated in  $\mathbb{R}^3$  to this  $f_s(t)$  to give  $A(t)$ ). The tangent space along a generator is spanned by  $A$  and  $X$  (and  $A$  is orthogonal to  $X$ ), hence this tangent space is constant along a generator iff  $c = 0$ . The function  $c/(c^2 + a^2)$  is called the *distribution parameter* and it is independent of the particular orthogonal trajectory  $f$  (which we show later). Thus (a)  $M$  is developable, (b)  $K = 0$ , (c)  $c = 0$ , (d)  $LX = 0$ , (note  $\langle LX, T \rangle = \langle LT, X \rangle - c$ ), and (e)  $\bar{D}_T X$  is tangent to  $M$ , are all equivalent for  $M$  closed and connected (assuming Massey's theorem).

Assuming  $M$  is closed (and ruled with  $c \neq 0$ ), on each generator there exists a distinguished point called the *central point*, and these points determine the curve of striction on the surface. Fixing two generators, say for  $t_1 < t_2$ , we compute the length  $J(s)$  of

an orthogonal trajectory between these two generators by

$$\begin{aligned} J(s) &= \int_{t_1}^{t_2} \sqrt{\langle A, A \rangle} dt \\ &= \int_{t_1}^{t_2} \sqrt{1 - 2as + a^2s^2 + c^2s^2} dt \end{aligned}$$

Let us find the value of  $s$  which minimizes  $J(s)$ , and we get  $J'(s) = 0$  if

$$-2a + 2(a^2 + c^2)s = 0$$

or  $s = a/(a^2 + c^2)$  at  $t_1$  as  $t_2 \rightarrow t_1$ . Hence the curve of stricture is the curve

$$f + \frac{a}{a^2 + c^2}X$$

as a function of  $t$ . This is precisely the point on each generator where the tangent plane is normal to  $\bar{D}_T X(t)$  since  $\bar{D}_T X$  is orthogonal to  $X$  we know, and

$0 = \langle \bar{D}_T X, A \rangle = -a + a^2s + c^2s$  again gives  $s = \frac{a}{a^2 + c^2}$ . As a problem we leave the formula for the curvature,

$$K(t, s) = \frac{-c^2}{(1 - 2as + a^2s^2 + c^2s^2)^2},$$

and hence the central point on each generator is also characterized as the point where  $K$  is a maximum ( $|K|$  a minimum). At the central point,

$K = -(a^2 + c^2)^2/c^2$ , which shows the distribution parameter  $\frac{c}{a^2 + c^2}$  depends only on the generator.

If  $s = 0$  gives the central point on a particular generator, i.e., we take our orthogonal curve  $f$  from this central point, then  $\bar{D}_T X$  is normal to  $m$  at  $s = 0$  and  $a = 0$ . Thus the distribution parameter  $p = 1/c$  and

$$K(t, s) = -\frac{c^2}{(1 + c^2s^2)^2} = -\frac{p^2}{(p^2 + s^2)^2}$$

. Along this generator  $A = T + csN$  where  $T$  and  $N$  are vectors

---

at the central point, hence the normal  $N(s)$  along the generator is given by

$$\begin{aligned} N(s) &= \frac{-scT + N}{\sqrt{1 + s^2c^2}} \\ &= \frac{-sT + pN}{\sqrt{p^2 + s^2}} \end{aligned}$$

Thus if  $\phi$  is the angle between the normal  $N(s)$  and the normal  $N$  at the central point, we have  $\tan \phi = s/p$ , i.e., the tangent of  $\phi$  is directly proportional to the distance from the central point. This is Chasles theorem (1839). This also shows the tangent plane turns even through  $180^\circ$  along a generator (turning  $90^\circ$  on either side of the central point). For references, see [Str61, p. 189] and [Wil59, p. 107].

We point out we could have viewed the ruled surface discussed above as being generated by the curve  $f(t)$  and the field  $X(t)$  along the curve. To generate surfaces in this way  $X$  need not be orthogonal to  $T$ . Indeed, in case  $\bar{D}_T T \neq 0$ , then we generate a surface via  $(t, s) \mapsto f(t) + sT(t)$ , for small  $s > 0$  (or small  $s < 0$ ), which we call the *tangential developable* of the curve  $f$ , which is the edge of regression of these two surfaces. It is a surface, since  $A = T + s\bar{D}_T T$  is independent of  $X = T$  (for  $s \neq 0$ ), and the tangent space along a generator will be determined by  $T$  and  $\bar{D}_T T$  for all  $s$ ; hence the surface is developable. It is, of course, not a closed surface in general (see. [Str61, p. 66]).

## 2.6 Some Applications

Let  $M$  be a hypersurface of  $\mathbb{R}^n$  with unit normal  $N = (a_1, \dots, a_n)$  where each  $a_i - I$  is a  $C^\infty$  function on  $M$  and  $\sum_1^n a_i^2 = 1$ . For any  $r$  in  $R$ , let

$M_r = \{p + rN_p : p \in M\}$ . Thus if  $p = (p_1, \dots, p_n)$  is in  $M$ , then

$$f(p) = p + rN_p = (p_1 + ra_1(p), \dots, p_n + ra_n(p))$$




Figure 2.6: Hyperboloid of Revolution

is in  $M_r$ . The map  $f$  is called the *natural map* of  $M$  into  $M_r$ , and if  $f$  is univalent, then  $M_r$  is a *parallel hypersurface* of  $M$  with unit normal  $N$ , i.e.,  $N_{f(p)} = N_p$  for all  $p$  in  $M$ . Let  $L_r$  be the Weingarten map on  $N_r$ .

**Theorem 2.6.** Let  $f: M \rightarrow M_r$  as just described. Then for  $X \in \mathbf{T}_p M$ ,  $f_*(X) = X + rL(X)$ ,  $L_r(f_*X) = L(X)$ , and  $f$  preserves principal directions of curvature, umbilics, and the third fundamental form. Also

$$\langle f_*X, f_*Y \rangle = I(X, Y) + 2r \text{ II}(X, Y) + r^2 \text{ III}(X, Y),$$

where  $I$ ,  $\text{II}$ ,  $\text{III}$  are the first, second, and third fundamental forms on  $M$ . If  $k$  is a principal curvature of  $M$  at  $m$  in direction  $X$ , then  $k/(1 + rk)$  is the corresponding principal curvature of  $M_r$  at  $f(m)$  in direction  $f_*X$ .

*Proof.* To compute  $f_*X$ , take a curve  $\sigma(t) = (b_1(t), \dots, b_n(t))$  with  $X = (b'_1(0), \dots, b'_n(0))$ , and compute the tangent to  $f \circ \sigma$  at  $t = 0$ . Let  $N(\sigma(t)) = (a_1(t), \dots, a_n(t))$ ; then  $f \circ \sigma(t) = (\dots, b_i(t) + ra_i(t), \dots)$ , and its tangent at  $t = 0$  is indeed  $X + rL(X)$ . Also  $N(\sigma(t)) = N(f \circ \sigma(t))$  from the definition of  $f$  and  $M_r$ . Thus  $L(X) = \bar{\mathbf{D}}_X N = (a'_1(0), \dots, a'_n(0)) =$

$\bar{D}_{f_*X}N = L_r(f_*X)$ . This shows

$$\text{III}_r(f_*X, f_*Y) = \langle L_r f_*X, L_r f_*Y \rangle = \langle LX, LY \rangle = \text{III}(X, Y)$$

Now let  $X$  be a unit vector at  $m \in M$  with  $LX = kX$ , so  $L_r(f_*X) = LX = kX$  and  $f_*X = (1 + rk)X$ . If  $1 + rk = 0$ , then  $f_*X = 0$  and  $L_r(f_*X) = kX = 0$ , so  $k = 0$  and  $1 = 0$ , thus  $1 + rk \neq 0$  if  $M_r$  is a hypersurface. Hence  $L_r(f_*X) = (k/(1 + rk))f_*X$ , which shows  $f$  preserves directions of curvature and umbilics. Finally, one can verify the expression for  $\langle f_*X, f_*Y \rangle$  by direction computation using  $f_*X = X + rLX$ .  $\square$

**Corollary 2.7.** In the hypothesis of the above theorem let  $n = 3$ , and let the total curvature and mean curvature of  $M$  (and  $M_r$ ) be denoted by  $K$  (and  $K_r$ ) and  $H$  (and  $H_r$ ). Then

$$K_r = \frac{K}{1 + rH + r^2K} \quad \text{and} \quad H_r = \frac{H + 2rK}{1 + rH + r^2K}.$$

**Theorem 2.8.** Let  $M$  be a connected hypersurface in  $\mathbb{R}^n$  consisting entirely of umbilics. Then  $M$  is either an open subset of a hyperplane or a sphere. If  $M$  is closed, then  $M$  is a hyperplane or a sphere.

*Proof.* Take  $p$  in  $M$  and  $X_p$  in  $\mathbf{T}_pM$ ,  $X_p \neq 0$ . Imbed  $X_p$  in a  $C^\infty$  field  $X$  about  $p$  and let  $Y$  be any other  $C^\infty$  field about  $p$  with  $X_p$  and  $Y_p$  independent. Let  $L = fI$  be the Ewingarten map where  $f$  is a  $C^\infty$  real valued function on  $M$  and,  $I$  is the identity of each tangent space. By the Codazzi-Mainardi equation 10,

$$0 = \mathbf{D}_X(fY) - \mathbf{D}_Y(fX) - f[X, Y] = (X_p f)Y_p - (Y_p f)X_p,$$

since  $\mathbf{D}_X Y - \mathbf{D}_Y X = [X, Y]$ . The independence of  $X_p$  and  $Y_p$  implies  $X_p f = 0$ . Since  $M$  is connected,  $f$  must be a constant function on  $M$  (problem 14).

Suppose  $L = kI$ ,  $k$  is constant on  $M$ . If  $k = 0$ , then  $L \equiv 0$  on  $M$ , so  $N$  is constant on  $M$  ( $\bar{D}_X N = 0$  for all  $X \in \mathbf{T}_p M$ ) and  $M$  must be an open subset of a hyperplane.

If  $k \neq 0$ , then we may assume  $k > 0$  by changing the sign of  $N$  if necessary. Let  $r = -1/k$  and let  $f: M \rightarrow \mathbb{R}^n$  by  $f(p) = p + rN_p$ . As in the preceding theorem, for all  $X$  in  $\mathbf{T}_p M$ ,  $f_*(X) = X + rL(X) = X - (1/k)kX = 0$ . Thus  $f_* = 0$ , and since  $M$  is connected,  $f$  is a constant map. Let  $c = p - (1/k)N_p$  for any  $p$  in  $M$ . Then all points of  $M$  are  $1/k$  units from  $c$ . Thus  $M$  is an open subset of a sphere about  $c$  of radius  $1/k$ .  $\square$

## Problems

23. (i) Let  $f$  be in  $C^\infty(\mathbb{R}^2, \mathbb{R})$ . Let  $M$  be the graph of  $f$ ; thus  $M = \{(x, y, f(x, y)) : (x, y) \in \mathbb{R}^2\}$ . Let  $W = \sqrt{(f_x)^2 + (f_y)^2 + 1}$  and let  $N = W^{-1}(-f_x, -f_y, 1)$ . Show  $X = (1, 0, f_x)$  and  $Y = (0, 1, f_y)$  span  $M_m$  at all  $m$  and  $N$  is a unit normal that is  $C^\infty$  on  $M$ .

- (ii) Let  $E = \langle X, X \rangle$ ,  $F = \langle X, Y \rangle$ , and  $G = \langle Y, Y \rangle$ . Show

$$L(X) = -\frac{f_{xx}G - f_{xy}F}{W^3}X + \frac{(f_{xy}E - f_{xx}F)Y}{W^3}$$

$$L(Y) = -\frac{(f_{xy}G - f_{yy}F)X + (f_{yy} - f_{xy}F)Y}{W^3}$$

$$K = \frac{f_{xx}f_{yy} - f_{xy}^2}{W^4}$$

$$H = -\frac{1}{W^3}(f_{xx}G + f_{yy}E - 2f_{xy}F).$$

- (iii) Compute  $b_{11} = \langle LX, X \rangle$ ,  $b_{12} = \langle LX, Y \rangle$ ,  $b_{22} = \langle LY, Y \rangle$ ,  $c_{11} = \langle LX, LX \rangle$ ,  $c_{12} = \langle LX, LY \rangle$ , and  $c_{22} = \langle LY, LY \rangle$ .

- (iv) Show

$$K = \frac{\det b_{ij}}{EG - F^2} = \left[ \frac{\det c_{ij}}{EG - F^2} \right]^{1/2} = \frac{2F_{xy} - E_{yy} - G_{xx}}{2W^4}.$$

Show  $L^2 - HL + KI = 0$  where  $I$  is the identity map. (Com-

pare with section 3.4).

24. Compute the invariants for the *right helicoid* (a ruled surface) which is the image of the map  $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $\phi(u, v) = (u \cos v, u \sin v, av)$  for  $a > 0$ .
25. Show a curve on a surface is a line of curvature iff the surface normals along the curve form a developable surface (Result is due to Monge).





### 3. Surfaces in $\mathbb{R}^3$

Throughout this chapter,  $M$  will denote a surface in  $\mathbb{R}^3$ , i.e.,  $M$  is a two-dimensional  $C^\infty$  submanifold of  $\mathbb{R}^3$ . Let  $N$  be a  $C^\infty$  unit normal field on  $M$  (such an  $N$  always exists locally). Let  $\bar{D}$  and  $D$  be the natural connexions on  $\mathbb{R}^3$  and  $M$ , respectively. Let  $L(X) = \bar{D}_X N$  be the Weingarten map for  $X$  tangent to  $M$ . Let  $U$  be the set of umbilics on  $M$  and let  $V = M - U$ . Let  $K$  and  $H$  denote the Gauss curvature (total curvature) and mean curvature functions on  $M$ , respectively.

Let  $h$  and  $k$  be the principal curvature functions on  $M$  where  $h(p) \geq k(p)$  for all  $p$  in  $M$ . Thus  $K(p) = \det L_p = h(p)k(p)$  and  $H(p) = \text{tr} L_p = h(p) + k(p)$  for  $p$  in  $M$ .

#### 3.1 Smoothness and the Neighborhood of a non-Umbilic Point

The first theorem establishes the smoothness of the invariants of  $M$  and the local existence of  $C^\infty$  orthonormal principal vectors on  $V$ .

**Theorem 3.1.** The set of umbilics  $U$  is closed in  $M$ , so its complement  $V$  is open in  $M$ . The functions  $K$  and  $H$  are  $C^\infty$  on  $M$ . The functions

$$h = \frac{H + \sqrt{H^2 - 4K}}{2} \quad \text{and} \quad k = \frac{H - \sqrt{H^2 - 4K}}{2}$$

are  $C^0$  on  $M$  and  $C^\infty$  on  $V$ . For any  $p \in V$  there is a neighborhood

$A$  of  $p$  with  $A \subset V$  and an orthonormal  $C^\infty$  base field of principal vectors on  $A$ .

*Proof.* For any  $m \in M$ , let  $B$  be the domain of a local coordinate system. By applying the Gram-Schmidt process to the coordinate vector fields on  $B$ , we obtain an orthonormal  $C^\infty$  base field  $Z, W$  on  $B$ . Since  $L$  is  $C^\infty$ , the vectors  $L(Z) = aZ + bW$  and  $L(W) = bZ + cW$  are  $C^\infty$  on  $B$ , and hence the functions  $a, b$ , and  $c$  are  $C^\infty$  on  $B$ . Thus  $K = ac - b^2$  and  $H = a + c$  are  $C^\infty$  on  $B$ , and hence, on  $M$ .

The eigenvalues  $h$  and  $k$  must satisfy the algebraic equation  $\lambda^2 - H\lambda + K = 0$  associated with the characteristic equation of  $L$ . Hence we get explicit global expressions for  $h$  and  $k$  by the quadratic formula and they are clearly continuous, since they are the composite of continuous functions. The set  $U$  is precisely the set where  $h = k$  or  $H^2 - 4K = 0$ , so by continuity,  $U$  is closed and  $V$  is open. Since  $H^2 - 4K > 0$  on  $V$ , the functions  $h$  and  $k$  are  $C^\infty$  on  $V$ .

For  $p$  in  $V$ , let  $B, Z$ , and  $W$  be as in the first paragraph, with  $B \subset V$ . We distinguish two cases:

1. If  $b(p) \neq 0$ : choose the neighborhood  $A \subset B$  such that  $b \neq 0$  on  $A$  and let  $Y' = bZ + (h - a)W$  and  $X' = (a - h)Z + bW$ . Then  $X', Y'$  are  $C^\infty$  orthogonal non-vanishing fields on  $A$  with  $LY' = hY'$  and  $LX' = k'$ . Let  $X$  and  $Y$  be unit fields in directions  $X'$  and  $Y'$ , respectively.
2. If  $b(p) = 0$ : suppose  $a(p) > c(p)$ , choose  $A \subset B$  so  $a > c$  on  $A$ , and let  $Y' = (h - c)Z + bW$  and  $X' = bZ + (c - h)W$ , etc.

□

In the next theorem we derive basic expressions for studying the neighborhood of a non-umbilic point.

**Theorem 3.2.** Let  $m$  be a non-umbilic point on  $M$  and let  $X$  and  $Y$  be an orthonormal  $C^\infty$  base field of principal vectors on the

Section 3.1: Smoothness and the Neighborhood of a non-Umbilic Point

---

neighborhood  $A$  of  $m$  with  $A \subset V$  and  $LX = kX, LY = hY$  on  $A$ . Defining the  $C^\infty$  functions  $a$  and  $b$  on  $A$  by

$$a = \frac{Yk}{h-k} \text{ and } b = -\frac{Xh}{h-k}$$

then  $\mathbf{D}_X Y = aX, \mathbf{D}_Y X = bY, \mathbf{D}_X X = -bY, \mathbf{D}_Y Y = -aX$

$$[X, Y] = aX - bY, \text{ and}$$

$$K = kh = \frac{(X^2h - Y^2k)(h-k) - (Xh)(2Xh - Xk) + (Yk)(Yh - 2Yk)}{(h-k)^2} \text{ on } A.$$

*Proof.* Since  $\langle X, X \rangle = 1, \langle Y, Y \rangle = 1,$  and  $\langle X, Y \rangle = 0$  on  $A,$   
 $0 = X \langle Y, Y \rangle = 2 \langle \mathbf{D}_X Y, Y \rangle$  so  $\mathbf{D}_X Y = aX$  for some  $C^\infty$  function  $a,$  which we compute below. Similarly,  $\mathbf{D}_X Y = bY$  for some  $b.$  Also  
 $0 = X \langle X, X \rangle = 2 \langle \mathbf{D}_X X, X \rangle$  and  $0 = X \langle X, Y \rangle = \langle \mathbf{D}_X X, Y \rangle + \langle X, \mathbf{D}_X Y \rangle,$  so  $\mathbf{D}_X X = -aY,$  and similarly,  $\mathbf{D}_Y Y = -bX.$  Then  $[X, Y] = \mathbf{D}_X Y - \mathbf{D}_Y X = aX - bY.$

To compute the expressions for  $a$  and  $b$  in terms of  $X, Y, h$  and  $k,$  we apply the Codazzi-Mainardi equation 10. Thus

$$\mathbf{D}_X LY - \mathbf{D}_Y LX = (Xh)Y + haX - (Yk)X - kbY = L([X, Y]) = akX - bhY.$$

Equating coefficients of  $X$  and  $Y$  leads to the expressions for  $a$  and  $b.$

To compute  $K,$  first notice

$$R(X, Y)Y = \mathbf{D}_X(-bX) - \mathbf{D}_Y(aX) - \mathbf{D}_{aX-bY}Y = -(Xb)X - (Ya)X - a^2X - b^2X.$$

By the Gauss curvature equations,  $K = \langle R(X, Y)Y, X \rangle = -(Xb) - (Ya) - a^2 - b^2,$  and the final expression for  $K$  follows by inserting the formulas for  $a$  and  $b$  and computing.

□

**Corollary 3.3.** If  $m$  is a non-umbilic critical point of both principal curvatures, then  $K(m) = \frac{X^2h - Y^2k}{h - k}$ . If  $H$  has no umbilics and  $K$  and  $H$  are constant (or the principal curvatures are constant) then  $K = 0$

### 3.2 Surfaces of Constant Curvature

Let  $M$  be a closed connected surface in  $\mathbb{R}^3$  with constant Gauss curvature  $K$ . Then  $M$  is a sphere, a developable surface, or doesn't exist, according as  $K > 0$ ,  $K = 0$ , or  $K < 0$ , respectively. The cases when  $K > 0$  (due to Liebmann) and  $K < 0$  (due to Hilbert) were solved around 1900. It is amazing that the case  $K = 0$  (due to Massey) was not completely solved until 1962.

Consider the case  $K > 0$ . The result of Liebmann follows from a lemma due to Hilbert.

**Lemma 3.4.** If  $K$  is a positive constant on  $M$ , then  $h$  cannot have a relative maximum (and  $k$  cannot have a relative minimum) at any non-umbilic point.

*Proof.* Suppose  $m \in V$  and  $m$  is a relative maximum for  $h$  and a relative minimum for  $k$  (since  $K = hk = \text{constant}$ ). With the notation Theorem 3.2,

$X^2h \leq 0$  and  $Y^2k \geq 0$  at  $m$ . Thus by the above corollary,  $K(m) \leq 0$ , which is a contradiction.  $\square$

A theorem of Bonnet, proved in Chapter 10, shows the “compact” assumption in the following theorem can be replaced by “closed”.

**Theorem 3.5.** A compact connected surface in  $\mathbb{R}^3$  of constant positive Gauss curvature is a sphere.

*Proof.* At all points, the principal curvature  $h \geq \sqrt{K}$ , since  $h^2 \geq hk = K$ . Since  $M$  is compact,  $h$  must have an absolute maximum  $m \in M$ , and  $m$  must be umbilic by Hilbert's lemma 3.4. Thus  $h(m) = k(m) =$

$\sqrt{K}$ , and hence  $h \leq \sqrt{K}$  on  $M$ . Thus  $h = \sqrt{K}$ , all points are umbilic, and  $M$  must be a sphere.  $\square$

The preceding theorem can be paraphrased by saying “a sphere cannot be bent”. For a precise interpretation of this phrase, see Chapter 8, where a generalization, the rigidity theorem for convex bodies, is proved.

A proof of Hilbert’s theorem stating that a closed connected surface with constant  $K < 0$  cannot exist in  $\mathbb{R}^3$  is in [Wil59]. Here again, the compact case is easily disposed of by the first corollary of the following theorem; indeed, no compact  $M$  exists with variable  $K \leq 0$  on  $M$ .

**Theorem 3.6.** On a compact surface in  $\mathbb{R}^3$  there is a point  $m$  with  $K(m) > 0$ .

*Proof.* Let  $r(p) = |p|$  give the distance from a point  $p \in \mathbb{R}^3$  to the origin. Then  $r \circ i$  is a continuous function on the compact surface  $M$  so it takes on a maximum at a point  $m \in M$ . By a rotation (orthogonal transformation) of  $\mathbb{R}^3$ , we may assume  $m$  lies on the  $z$ -axis (or  $u_3$ -axis). Let  $N$  be a  $C^\infty$  unit normal to  $M$  on a neighborhood of  $m$  with  $N_m = (0, 0, 1)$ . Let  $X$  be any unit principal vector at  $m$  with  $L(X) = \bar{D}_X N = kX$ .

Let  $\sigma(t) = (f(t), g(t), h(t))$  be a  $C^\infty$  curve on  $M$  with unit tangent vector  $X$  at  $t = 0$ ; thus  $X = (f'(0), g'(0), h'(0))$ . Since  $m$  is an absolute maximum of  $u_3 \circ i = z \circ i$  on  $M$ ,  $h''(0) < 0$ . Letting  $X$  be the tangent to  $\sigma$ , we have at  $m$ ,  $\bar{D}_X X = (f''(0), g''(0), h''(0))$ . Decomposing this vector into tangent and normal components, we get, by the Gauss equation,

$$-\langle LX, X \rangle N = (0, 0, -k) = (0, 0, h''(0)),$$

so  $k = -h''(0) > 0$ .

Since all principal curvatures are greater than zero at  $m$ ,  $K(m) > 0$ .  $\square$

Notice the theorem is true for any compact hypersurface in  $\mathbb{R}^n$  with a trivial modification of the proof.

**Corollary 3.7.** There is no compact hypersurface in  $\mathbb{R}^n$  with non-positive Gauss curvature at all points.

**Corollary 3.8.** There is no compact minimal ( $H = 0$ ) surface in  $\mathbb{R}^3$ .

*Proof.* If  $H = 0$ , then  $k = -h$  and  $K = -h^2 \leq 0$ . □

Before considering the case  $K = 0$ , recall that a generator on a surface  $M$  is a straight line in  $\mathbb{R}^3$  that lies on  $M$  with the normal to  $M$  constant along the line. A *developable surface* is a ruled surface with the normal constant along the ruling lines in the surface. If a developable surface is closed, then it has a generator through each point.

**Theorem 3.9.** Let  $M$  be a closed connected surface in  $\mathbb{R}^3$  with  $K = 0$  on  $M$ . Then either  $M$  is a plane, or through each point of  $M$  passes a unique generator and all generators are parallel in  $\mathbb{R}^3$ . Moreover, the mean curvature is constant along generators, and hence the boundary of the umbilic set is a union of these generators.

*Proof.* Supposing  $M$  is not a plane; then the set  $V$  is non-empty. Let  $A$  be a connected neighbourhood in  $V$  as described in theorems 3.1 and 3.2. Since  $H$  does not vanish on  $V$  and  $A$  is connected, we may assume  $H = h > 0$  while  $k = 0$  on  $A$ . Theorem 3.1 gives an orthonormal pair of  $C^\infty$  fields  $X$  and  $Y$  on  $A$ , with  $LX = 0$  and  $LY = HY$  on  $A$ . Since  $Yk = 0$  on  $M$ , referring to theorem 3.2 we have  $a = 0$  on  $A$ , so  $D_X Y = 0$  and  $D_X X = 0$  on  $A$ . By the Gauss equation,  $\bar{D}_X X = D_X X - \langle LX, X \rangle N = 0$  on  $A$ . Thus the integral curves of  $X$  in  $A$  are straight line segments in  $\mathbb{R}^3$ . Since  $M$  is closed, the continuation of these line segments must lie in  $M$ . Hence for  $p$  in  $V$  there is a unique line  $G_p$  through  $p$  with  $G_p \subset M$ . We next show  $G_p \subset V$ .

On the neighborhood  $A$  of  $p$ , by theorem 3.2,

$$K = 0 = \frac{X^2 H}{H} - \frac{2(XH)^2}{H^2} = -HX^2 \left( \frac{1}{H} \right).$$

Hence if  $s$  is the arc length on  $G_p$  in the direction  $X$  with  $s = 0$  at  $p$ , then

$$\frac{1}{H} = cs + d \implies H = \frac{1}{cs + d} \text{ for points in } G_p \cap A.$$

If there was an umbilic point at  $s'$  on  $G_p$  then  $H(s') = 0$ . At  $s'' = \inf\{s' : s' \text{ is umbilic}\}$ ,  $H(s'') = 1/(cs'' + d) \neq 0$ , since  $H$  is continuous. Hence there are no umbilics on  $G_p$ ,  $G_p \subset V$ , and to avoid an impossible singularity in  $H$  at  $s = -c/d$ , it follows  $H$  is constant on  $G_p$ .

After extending  $X$  and  $Y$  along  $G_p$  by letting  $X$  be the unit tangent to  $G_p$ , an overlapping neighborhood argument will show  $X$  and  $Y$  remain principal vectors; hence  $L(X) = 0$  and  $L(Y) = HY$  on  $G_p$ . Then  $\bar{D}_X N = L(X) = 0$  implies  $N$  is constant on  $G_p$ , so  $G_p$  is a generator.

In the neighborhood  $A$ , since  $H$  is constant in the  $X$  direction, by theorem 3.2,  $D_Y X = 0$ , and so  $\bar{D}_Y X = D_Y X - \langle LX, Y \rangle N = 0$ . Thus  $X$  is parallel in  $\mathbb{R}^3$  along an integral curve of  $Y$ , which implies all generators through points in  $A$  are parallel. This implies all generators in one connected component of  $V$  must be parallel by another overlapping neighborhood argument. Hence the boundary of one connected component of  $V$  consists of two (or just one) lines parallel to the generators in that component. Consider now a connected component  $U_1$  of the umbilic set. If  $U_1$  has a non-empty interior in  $M$ , then this interior is an open surface of umbilics with  $K = 0$ , and hence it is an open subset of a plane in  $\mathbb{R}^3$ . This open plane subset is bounded by two generator lines in the boundary of  $V$ , and these generator lines cannot intersect (by the uniqueness of the generators through points in  $V$  and its boundary), and hence they are parallel. Thus parallel generators are defined through all points of  $M$ . □

**Corollary 3.10.** A closed connected surface is a developable surface iff its Gauss curvature is identically zero.

Problem 26 provides additional theorems leading to surfaces with constant  $K$  and  $H$ , and it is hoped that by now their proofs would provide little difficulty. Another “classic” type of argument is provided by the following theorem and some of the theorems in the next section.

**Theorem 3.11.** Let  $M$  be a closed connected surface whose sphere map (Gauss map) is strictly conformal. Then  $M$  is a sphere or a minimal surface with negative curvature. If  $M$  is compact, it must be a sphere.

*Proof.* Let  $\eta : M \rightarrow S$  be the sphere map. Since  $\eta$  is strictly conformal, there is a  $C^\infty$  positive real valued scale function  $F$  on  $M$  with

$$\langle \eta_* X, \eta_* Y \rangle = \langle LX, LY \rangle = F(m) \langle X, Y \rangle$$

for all  $X, Y$  in  $T_m M$  for all  $m$  in  $M$ . Hence  $\langle L^2(X) - F(m)X, Y \rangle = 0$  for all  $Y$  so  $L^2(X) = FX$  for all  $X$ . One always has  $L^2 - HL + KI = 0$ , where  $I$  is the identity map; hence  $HL = (K+F)I$ . If  $H(m) \neq 0$ , then  $m$  is an umbilic and  $K(m) = H^2(m)/4 > 0$ . If  $m$  is umbilic and  $H(m) = 0$ , then  $K(m) = -F(m) < 0$ , but an umbilic  $K(m) = k^2(m) \geq 0$  always. Thus the umbilic set  $U$  is exactly the set of  $m$  where  $H(m) \neq 0$ , and hence  $U$  is open and closed. Since  $M$  is connected, either  $M = U$  and  $M$  is a sphere ( $F > 0$  rules out a plane) or  $M = V$ ,  $H = 0$ , and  $K = -F < 0$ .

The last assertion of the theorem follows from corollary 3.7.  $\square$

### 3.3 Parallel Surfaces (Normal Maps)

Let us state a standard hypothesis for some theorems (and problems on “parallel surfaces’):  $M$  is a closed connected surface in  $\mathbb{R}^3$  with  $C^\infty$  unit normal  $N$ ,  $r$  is a non-zero real number, and  $f$  is a map  $f : M \rightarrow \mathbb{R}^3$  defined by  $f(p) = p + N_p$  (see section 2.6).



---

### Section 3.3: Parallel Surfaces (Normal Maps)

**Theorem 3.12.** With the standard hypothesis, if  $f$  is strictly conformal, then  $M$  is a sphere, plane, or has constant mean curvature  $H = -\frac{2}{r}$  with no umbilics.

**Proof.** From section 2.6, if  $X \in \mathbf{T}_m M$ , then  $f_*(X) = X + rL(X)$ . Since  $f$  is strictly conformal, there is a  $C^\infty$  real-valued function  $F$  on  $M$  with

$$\langle f_*X, f_*Y \rangle = F(m) \langle X, Y \rangle = \langle X + 2rLX + r^2L^2X, Y \rangle$$

for all  $X, Y$  in  $\mathbf{T}_m M$  for all  $m$  in  $M$ . Hence  $r^2L^2 + 2rL + (1 - F)I = 0$  and, as always,  $L^2 - HL + KI = 0$ , so

$$\left(H + \frac{2}{r}\right)L = \left[K - \frac{1 - F}{r^2}\right]I.$$

If  $H(m) + 2/r \neq 0$ , then  $m$  is an umbilic, and, indeed,  $U = \{m \in M : H(m) \neq -2/r\}$ . For if  $m$  umbilic and  $H(m) = -2/r = 2k$ , then

$$k = -\frac{1}{r}, K = \frac{1}{r^2}, K - \frac{1 - F}{r^2} = \frac{F}{r^2} = 0$$

, and so  $F(m) = 0$ , which is impossible. Thus  $M = U$  or  $M = V$ , and the only possibilities give the conclusion of the theorem.  $\square$

**Theorem 3.13.** With the standard hypothesis, if  $f$  preserves the second fundamental form, then  $M$  is a plane.

**Proof.** From section 2.6, for all  $X$  and  $Y$  in  $\mathbf{T}_m M$ ,

$$\langle LX, Y \rangle = \langle L_r f_*X, f_*Y \rangle = \langle LX, Y + rLY \rangle$$

thus  $\langle LX, rLY \rangle = \langle X, rL^2Y \rangle = 0$  for all  $X$  and  $Y$ , and hence  $L^2 = 0$ . Thus the principal curvatures are zero,  $L = 0$ , and  $M$  is a plane.  $\square$

Similar results are given as problems. The following theorem is due to Bonnet, and the examples in the next section show the hypothesis is not vacuous.

**Theorem 3.14.** Let  $M$  be a surface of constant positive Gauss curvature  $K$  with no umbilics. Let  $r_1 = 1/\sqrt{K}$  and  $r_2 = -1/\sqrt{K}$  define parallel sets  $M_1$  and  $M_2$  respectively. Then  $M_1$  and  $M_2$  are immersions of  $M$  which have constant mean curvature  $\sqrt{K}$  and  $-\sqrt{K}$ , respectively. If  $M'$  is a surface with constant mean curvature  $H$  (non zero) and non-zero Gauss curvature, letting  $r = -1/H$  yields a parallel set that is an immersion of  $M'$  with constant positive Gauss curvature  $H^2$ .

*Proof.* The proof is a corollary to the formulas for  $H_r$  and  $K_r$  in section 2.6. The special assumptions avoid trivial cases (sphere or cylinder) and singularities.

For the first part,  $f_*$  is non-singular, since for principal vectors

$$f_*X = (1 + rk)X \text{ and } 1 + rk = 1 \pm \frac{k}{\sqrt{K}} \neq 0,$$

since there are no umbilics. Then

$$H_1 = \frac{H + 2\sqrt{K}}{2 + H/\sqrt{K}} = \sqrt{K},$$

and similarly,  $H_2 = -\sqrt{K}$ .

For the second part,  $f_*$  is non-singular, since  $1 + rk = 1 - k/H = 0$  would imply  $k = H$ , so the other principal curvature is zero and  $K = 0$  contrary to the hypothesis. Then

$$K_r = \frac{K}{1 - 1 + K/H^2} = H^2$$

.

□

### 3.4 Examples (Surfaces of Revolution)

Some general methods for computations with “parameterized” surfaces are introduced in this section. Let  $A$  be an open set in  $\mathbb{R}^2$  and let  $\phi: A \rightarrow \mathbb{R}^3$  be defined by the three real-valued slot functions

---

### Section 3.4: Examples (Surfaces of Revolution)

$f, g$  and  $h$ , so

$\phi(u, v) = (f(u, v), g(u, v), h(u, v))$  for  $(u, v) \in A$ . Write  $T_u = (f_u, g_u, h_u)$ , where  $f_u = \frac{\partial f}{\partial u}$ ,  $T_{uv} = (f_{uv}, g_{uv}, h_{uv})$ , etc. Notice  $T_u = \phi_*\left(\frac{\partial}{\partial u}\right)$  is the tangent to the  $u$ -parameter curves on  $\phi(A)$ . Let us assume  $T_u \times T_v \neq 0$ , where “ $\times$ ” is the cross-product of advanced calculus; thus  $\phi$  is an immersion of  $A$  into  $\mathbb{R}^3$ . Let  $N = (T_u \times T_v)/W$  with  $W = |T_u \times T_v| \neq 0$  on  $A$ .

To compute the Weingarten map  $L$  associated with  $N$ , notice  $L(T_u) = \bar{D}_{T_u}(N) = N_u$ . Notice

$$\begin{aligned} \langle L(T_u), T_u \rangle &= \langle N_u, T_u \rangle \\ &= W^{-1} \langle (T_{uu} \times T_v) + (T_u \times T_{uv}) - W_u N, T_u \rangle \\ &= -\langle T_{uu}, N \rangle. \end{aligned}$$

Similarly,  $\langle L(T_i), T_j \rangle = -\langle T_{ij}, N \rangle$  with obvious values of  $i$  and  $j$ . In case  $T_u$  and  $T_v$  are orthogonal,

$$L(T_u) = -\frac{\langle T_{uu}, N \rangle}{\langle T_u, T_u \rangle} T_u - \frac{\langle T_{uv}, N \rangle}{\langle T_v, T_v \rangle} T_v$$

and similarly for  $T_v$ ; hence,

$$H = -\frac{\langle T_{uu}, N \rangle}{\langle T_u, T_u \rangle} = -\frac{\langle T_{vv}, N \rangle}{\langle T_v, T_v \rangle}$$

and

$$K = \frac{\langle T_{uu}, N \rangle \langle T_{vv}, N \rangle - \langle T_{uv}, N \rangle^2}{\langle T_u, T_u \rangle \langle T_v, T_v \rangle}.$$

A little more computation is necessary to determine the matrix for  $L$  in terms of  $T_u$  and  $T_v$  when they are not orthogonal.

Specializing further, let  $f$  be a positive (at least  $C^2$ ) function, and for  $u > 0$  let

$$\phi(u, v) = (u \cos v, u \sin v, f(v))$$


---

define a “surface of revolution”. Applying the above analysis, one sees directly that  $T_u$  and  $T_v$  are principal vectors and  $K = \frac{f' f''}{u[1 + (f')^2]}$  where  $f'(u) = \frac{df}{du}$ . To find surfaces of constant curvature one must solve the differential equation  $f' f'' = uK[1 + (f')^2]$ , a task that is left to reader via several problems. For more details and pictures, see [Str61].

### 3.5 Lines of Curvature

In this section we place some results involving lines of curvature, i.e., curves whose tangent vectors are principal directions of curvature.

*Definition.* A triply orthogonal system of surfaces in a neighborhood  $U$  of  $\mathbb{R}^3$  is a family of surfaces such that through each point of  $U$  there passes exactly three members of the family whose normals are mutually perpendicular.

*Theorem 3.15 (Dupin).* Intersecting surfaces from a triply orthogonal system intersect along a line of curvature.

*Proof.* Let  $S_1, S_2$  and  $S_3$  be mutually orthogonal families of surfaces with unit normals  $N_i$ , respectively. Let  $L_i X = \bar{D}_X N_i$ , as usual. The field  $N_3$  is a tangent to the intersection of  $S_1$  and  $S_2$ , so one must show  $N_3$  is a principal direction on  $S_1$  and  $S_2$  or  $L_i(N_3) = a_i N_3$  for  $i = 1, 2$ . This is equivalent to showing  $L_i(N_3)$  is orthogonal to  $N_1$  and  $N_2$  for  $i = 1, 2$ . To be specific, consider  $L_1(N_3)$ . Since  $L_1(N_3)$  is tangent to  $S_1$ ,  $\langle L_1 N_3, N_1 \rangle = 0$ . While

$$\langle L_1 N_3, N_2 \rangle = \langle \bar{D}_{N_3} N_1, N_2 \rangle = - \langle -N_1, \bar{D}_{N_3} N_2 \rangle = - \langle N_1, L_2 N_3 \rangle = - \langle L_2 N_1, N_3 \rangle,$$

since  $L_2$  is self-adjoint. Thus by symmetry, as one cyclically permutes the indices,

$$\langle L_1 N_3, N_2 \rangle = - \langle L_2 N_1, N_3 \rangle = + \langle L_3 N_2, N_1 \rangle = - \langle L_1 N_3, N_2 \rangle$$

Hence  $\langle L_1 N_3, N_2 \rangle = 0$ . □

Examples of triply orthogonal coordinate systems are given by the coordinate surfaces in rectangular coordinates, cylindrical coordinates, and spherical coordinates. Another example is provided by a system of confocal quadrics, i.e., the surfaces

$$\sum_{i=1}^3 \frac{(x_i)^2}{a_i - \lambda} = 1 \text{ with } a_1 < a_2 < a_3 \text{ fixed,}$$

are orthogonal for unequal values of  $\lambda$  ([Str61, p. 100]). The classic work in this area is by Darboux.

**Theorem 3.16 (Liouville).** A conformal diffeomorphism of  $\mathbb{R}^3$  onto  $\mathbb{R}^3$  maps spheres into spheres.

*Proof.* Let  $S$  be a sphere. For  $p \in S$ , take an orthogonal family of curves on  $S$  and use the normal direction to  $S$  to generate an orthogonal family of surfaces. Adding in the “parallel” surfaces to  $S$ , one obtains a triply orthogonal system about  $p$ . Let  $f$  be a map in question, so  $f$  maps a neighborhood of  $p$  into a triply orthogonal system of surfaces about  $f(p)$  on  $f(S)$ . By Dupin’s theorem 3.15, the images of our original family of curves on  $S$  must be lines of curvature on  $f(S)$ . But we may choose an orthogonal family of curves on  $S$  to pass through any orthonormal pair of vectors  $X$  and  $Y$  at  $p$ . Hence all vectors tangent to  $f(S)$  and  $f(p)$  are principal, and  $f(p)$  is an umbilic of  $f(S)$ . Thus  $f(S)$  is completely umbilic, and since it is compact and connected it must be a sphere. □

The differentiability hypothesis in the above theorem is much too strong. The theorem can be used to show a conformal map of  $\mathbb{R}^3$  onto  $\mathbb{R}^3$  is a combination of similarities and isometries (also due to Liouville). For more details see [Gug12, p.225].

We next discuss the behaviour of the normal lines (in  $\mathbb{R}^3$ ) to a surface  $M$  along a line of curvature  $C$ . Let  $k$  be the principal curvature of  $M$  along  $C$  with respect to the unit normal field  $N$ ,

and let  $X$  be a unit tangent to  $C$ . If  $k \equiv 0$  on  $C$ , then  $\bar{D}_X N = LX = kX = 0$  implies  $N$  is a constant field (in  $\mathbb{R}^3$ ) along  $C$ , and  $C$  is a plane curve (see section 6.3); thus the normal lines form a “cylinder”, a developable surface. If  $k$  is a constant ( $\neq 0$ ) along  $C$ , let  $C(t)$  be the parameterization of  $C$  by arc length in the direction  $X$ , so  $X(t) = C'(t) = \frac{d}{dt}C$ . Then  $kX = kC'$ ; thus all the normal lines along the curve pass through a single point (and thus form a “cone”). If  $k \neq 0$  and  $k' \neq 0$  along  $C$ , then let  $B(t) = C(t) + f(t)N(t)$ , so  $B' = X + fkX + f'N$ , and choosing  $1 + fk = 0$  or  $f(t) = -1/k(t)$ , we obtain a curve  $B$  whose tangent developable gives the normal lines along  $C$ .

When both principal curvatures  $k$  and  $h$  are non-zero and non-constant in a neighborhood of  $p$ , then the points  $p - (1/k)N$  and  $p - (1/h)N$  are called the centers of principal curvature of  $p$  on  $M$ . The loci of the centres of principal curvature are called center surfaces (see [Str61, p. 95]).

## Problems

All surfaces are in  $\mathbb{R}^3$ .

26. (i) If  $M$  is a closed connected surface with  $K = 0$  and  $H$  constant, show  $M$  is a plane or a right circular cylinder.
- (ii) If  $M$  has no umbilics and  $K$  and  $H$  are constant, show  $M$  is a right circular cylinder.
- (iii) If  $I = II$  or if  $I = III$ , show  $M$  is a sphere of radius one, and vice versa.
- (iv) If  $II = III$ , show  $M$  is a sphere of radius one, a plane, or a right circular cylinder of radius one.
27. For  $u < \left| \frac{1}{b} \right|$  and  $b > 0$ , let

$$f(u) = \int_0^u \frac{bt}{\sqrt{1 - b^2t^2}} dt$$

and let  $\phi : (u, v) \mapsto (u \cos v, u \sin v, f(v))$ . Show that the surface of revolution determined by  $\phi$  for  $0 < u < \frac{1}{b}$  is an open subset of a sphere with curvature  $b^2$ .

28. For  $0 < u < \frac{1}{b}$  and  $b > 0$ , let

$$f(u) = \int_0^{b^{-1} \log bu} \sqrt{1 - e^{2bt}} dt.$$

Show that the surface of revolution induced by  $f$  has constant curvature  $-b^2$  and draw its graph (tractrix).

29. Find a surface of constant positive curvature that is not an open subset of a sphere.
30. Show a surface is minimal ( $H = 0$ ) iff there are orthogonal asymptotic vectors at each point.
31. Let  $f(u) = \cosh^{-1} u$  for  $u > 1$ , and show the surface of revolution induced by  $f$  (catenoid) is a minimal surface ( $H = 0$ ).





## 4. Tensors and Forms

### 4.1 Tensors and Forms

The material in the first three chapters was based on a minimum amount of structure, i.e., manifolds, functions, and vector fields; moreover, there was a strong bias on hypersurfaces in Euclidian space. By the time the reader should be at home with these concepts, and before discussion general connexions on manifolds, it is convenient to define tensors and forms. They are there, and they are useful. At times in the past, one notices a strong compulsion to seek out and label tensors *ad nauseum*, and objects that were not tensors were eyed with suspicion. In a sense, this chapter is the “7th” section of Chapter 1; it is just more structure that a  $C^\infty$  manifold has automatically, and Chapter 7 continues the theme. It is hoped by breaking the definitions up they become more digestible.

Let  $M$  be a  $C^\infty$   $n$ -manifold throughout this chapter, and let  $m$  be a point in  $M$ . Since the tangent space  $\mathbf{T}_m M$  at  $m$  is an  $n$ -dimensional vector space, the theory of linear algebra can be applied to define tensors and forms. A  $p$ -covariant tensor at  $m$  (for  $p > 0$ ) or a  $p$ -co tensor at  $m$  is a real valued  $p$ -linear (i.e., linear in each slot) function on  $\mathbf{T}_m M \times \mathbf{T}_m M \times \cdots \times \mathbf{T}_m M$  ( $p$  copies). Thus  $\alpha$  is a 2-co

tensor at  $m$  if

$$\alpha(X + Y, Z) = \alpha(X, Z) + \alpha(Y, Z)$$

$$\alpha(X, Y + Z) = \alpha(X, Y) + \alpha(X, Z)$$

$$\alpha(rX, Y) = \alpha(X, rY) = r\alpha(X, Y)$$

for all  $X, Y$  and  $Z$  in  $\mathbf{T}_m M$  and  $r \in \mathbb{R}$ . In a similar way, one defines a  $V$ -valued  $p$ -co tensor at  $m$ , where  $V$  is any vector space over  $\mathbb{R}$ ; indeed,  $V$  could be  $\mathbf{T}_m M$  itself.

Let  $\mathbf{T}_m^* M$  be the dual space of  $\mathbf{T}_m M$ . Thus  $\mathbf{T}_m^* M$  is the set of real valued 1-co tensors at  $m$ , or the set of linear functionals from  $\mathbf{T}_m M$  into  $\mathbb{R}$ , and  $\mathbf{T}_m^* M$  is endowed with its natural vector space structure (i.e., one adds functions by adding their values and multiplies by a constant in an obvious way). Similarly, the set of  $p$ -co tensors at  $m$ , denoted by  $\mathbf{T}_m^{0,p} M$ <sup>1</sup>, is a vector space over  $\mathbb{R}$ . A  $p$ -contravariant or  $p$ -contra tensor at  $m$  (for  $p > 0$ ) is a real valued  $p$ -linear function on  $(\mathbf{T}_m^* M)^p$ , the cross product of  $p$  copies of  $\mathbf{T}_m^* M$ , and the natural vector space formed by  $p$ -contra tensors at  $m$  is denoted by  $\mathbf{T}_m^{p,0} M$ . Define  $\mathbf{T}_m^{0,0} M = \mathbb{R}$ . (The sets of  $p$ -co tensor and  $p$ -contra tensors on any vector space  $W$  are denoted by  $\mathbf{T}^{0,p} W$  and  $\mathbf{T}^{p,0} W$ , respectively.) Again,  $V$ -valued  $p$ -contra tensors are defined analogously. Finally, a  $p$ -co and  $q$ -contra tensor at  $m$  is a  $(p+q)$ -linear real valued function on  $(\mathbf{T}_m M)^p \times (\mathbf{T}_m^* M)^q$ , and the vector space of these tensors is denoted by  $\mathbf{T}_m^{q,p} M$ . If  $p$  and  $q$  are greater than zero, elements of  $\mathbf{T}^{p,q}$  are called *mixed tensors*. Notice that a vector at  $m$  is a 1-contra tensor at  $m$ . Similarly, there is a special name for a 1-co tensor at  $m$ , for it is called a *1-form* at  $m$ .

A tensor is *symmetric* iff its value remains the same for all possible permutations of its arguments (thus only  $\mathbf{T}^{p,0}$  or  $\mathbf{T}^{0,p}$  tensors can be symmetric). A tensor is *skew-symmetric* or *alternating* iff its value after any permutation of its arguments is the product of

---

<sup>1</sup>In the original book, the tangent space was notated  $M_m$ , and  $\mathbf{T}^{p,q} V$  denoted the  $(p, q)$  tensors over  $V$ . So the original book had  $\mathbf{T}^{p,q}(M_m)$  to denote this. However as we decided to change notations for tangent space, and since  $\mathbf{T}^{p,q}(\mathbf{T}_m M)$  does not look good, we changed the notation here. So whenever we say  $\mathbf{T}_m^{p,q} M$ , what is really meant is  $(p, q)$  tensors over  $\mathbf{T}_m M$ .

its value before the permutation and the sign of the permutation. For example, let  $\alpha$  be a 3-co tensor at  $m$  and let  $\pi$  be a permutation of the set  $\{1, 2, 3\}$ . Then  $\alpha$  is symmetric iff

$$\alpha^\pi(X_1, X_2, X_3) = \alpha(X_{\pi 1}, X_{\pi 2}, X_{\pi 3}) = \alpha(X_1, X_2, X_3)$$

for all permutations  $\pi$  and all vectors  $X_i$  in  $\mathbf{T}_m M$ . When  $\alpha^\pi$  is defined by the first equality in the above line,  $\alpha$  is alternating iff  $\alpha^\pi = (-1)^\pi \alpha$ , where  $(-1)^\pi$  is the sign of the permutation  $\pi$ . Then a  $p$ -form at  $m$  (for  $p > 0$ ) is an alternating  $p$ -co tensor at  $m$ , and the set of  $p$ -forms at  $m$  is denoted by  $\Omega_m^p(M)$ . A 0-form at  $m$  is a real number; thus  $\Omega^0(W) = \mathbb{R}$  for any vector space  $W$  over  $\mathbb{R}$ . A  $p$ -form is said to be of degree  $p$ .

Tensor fields and  $C^\infty$  tensor fields are now defined in a way that is analogous to the definition of a vector field, once a vector was defined. For example, a  $p$ -co tensor field on a set  $U$  is a mapping that assigns to each  $m$  in  $U$  a  $p$ -co tensor at  $m$ . A  $p$ -co tensor field  $\alpha$  on a set  $U$  is  $C^\infty$  iff  $U$  is open and for all sets of  $C^\infty$  vector fields  $X_1, \dots, X_p$  on  $U$ , the function

$$[\alpha(X_1, \dots, X_p)](m) = \alpha_m(X_1(m), \dots, X_p(m))$$

is a  $C^\infty$  function on  $U$ . A  $C^\infty$   $p$ -form field on an open set  $U$  is called a differential  $p$ -form on  $U$ .

The tensor product of covariant tensors is defined as follows: if  $\alpha$  in  $\mathbf{T}^{0,p}W$  and  $\beta$  in  $\mathbf{T}^{0,q}W$ , then  $\alpha \otimes \beta$  is the element in  $\mathbf{T}^{0,p+q}W$  defined by

$$(\alpha \otimes \beta)(X_1, \dots, X_{p+q}) = \alpha(X_1, \dots, X_p)\beta(X_{p+1}, \dots, X_{p+q})$$

for all  $X_i$  in  $W$ . Notice that

- $(\alpha_1 + \alpha_2) \otimes \beta = (\alpha_1 \otimes \beta) + (\alpha_2 \otimes \beta)$ ,
- $\alpha \otimes (\beta_1 + \beta_2) = (\alpha \otimes \beta_1) + (\alpha \otimes \beta_2)$ ,

## Chapter 4: Tensors and Forms

---

- $(r\alpha) \otimes \beta = \alpha \otimes (r\beta) = r(\alpha \otimes \beta)$  for  $r$  in  $\mathbb{R}$ ,
- However, in general,  $\alpha \otimes \beta \neq \beta \otimes \alpha$
- $(\alpha \otimes \beta) \otimes \gamma = \alpha \otimes (\beta \otimes \gamma)$ .

Thus the tensor product is bilinear and associative but not symmetric. The tensor product of contravariant tensors or mixed tensors is defined analogously, but the details are omitted since these products are rarely used in this study.

If  $\alpha$  and  $\beta$  are forms of degree  $p$  and  $q$ , respectively, then the exterior, wedge, or Grassman product  $\alpha \wedge \beta$  is defined to be the  $(p+q)$ -form

$$\alpha \wedge \beta = \left( \frac{1}{p!q!} \right) \sum (-1)^\pi (\alpha \otimes \beta)^\pi,$$

where the sum is taken over all permutations  $\pi$  of the set  $\{1, 2, \dots, p+q\}$ . In problem 35 there is an expression for  $\alpha \wedge \beta$  that avoids division. Notice that

- $\alpha \wedge \beta = (-1)^{pq} \beta \wedge \alpha$ ,
- $\alpha \wedge (\beta_1 + \beta_2) = \alpha \wedge \beta_1 + \alpha \wedge \beta_2$ , where  $\beta_i$  are forms of the same degree,
- $(\alpha \wedge \beta) \wedge \gamma = \alpha \wedge (\beta \wedge \gamma)$  which is proved by using problem 35.

To continue the definitions in terms of the abstract vector space  $W$  over  $\mathbb{R}$ , the tensor algebra  $\mathbf{T}W$  over  $W$  and the Grassman algebra (exterior algebra)  $\Omega(W)$  over  $W$  are defined as the weak direct sums

$$\mathbf{T}W = \sum_{p,q \geq 0} \mathbf{T}^{p,q}W \quad \text{and} \quad \Omega(W) = \sum_{p \geq 0} \Omega^p(W).$$

By a weak direct sum,  $\sum_I M_i$  of modules over an index set  $I$ , one means the set of formal finite linear combinations of elements  $m_1 + m_2 + \dots + m_k$  where each  $m_i$  in  $M_i$ ; or more precisely,

$$\sum_I M_i = \left\{ f \in \prod_I M_i : f(i) = 0 \text{ for all but a finite number of elements } i \in I \right\},$$

and the one writes  $f = m_1 + m_2 + \dots + m_k$  if  $f(i) = m_i$  for  $i = 1, \dots, k$  and  $f(j) = 0$  for  $j \neq 1, \dots, k$ , (see [Che56] and [Jac13] for more details). The tensor multiplication and the exterior product can be extended distributively to  $\mathbf{T}W$  and  $\Omega(W)$ , respectively, thus making them algebras over  $\mathbb{R}$ .

If  $U$  is an open set in the manifold  $M$ , let  $\mathbf{T}^{p,q}U$  be the set of  $C^\infty$   $p$ -contra and  $q$ -co tensor fields on  $U$ , and let  $\mathbf{T}U$  and  $\Omega(U)$  be defined analogously. On the other hand, let  $\mathcal{F}$  be the ring of  $C^\infty$  real valued functions on  $U$  and let  $\mathcal{X}_u$  be the  $\mathcal{F}_u$ -module of  $C^\infty$  vector fields on  $U$ . Then the above definitions can be extended to define the  $\mathcal{F}_u$ -modules  $\mathbf{T}^{p,q}\mathcal{X}_u$  and  $\Omega^p(\mathcal{X}_u)$  for  $p, q \geq 0$ , where  $\mathbf{T}^{0,0}\mathcal{X}_u = \Omega^0(\mathcal{X}_u) = \mathcal{F}_u$ . The next theorem and its corollary are designed to illuminate the relation between  $\mathbf{T}U$  and  $\mathbf{T}\mathcal{X}_u$ . To accomplish this, let us define an open set  $V$  in  $M$  to be *framed* if there exists a  $C^\infty$  base field on  $V$ , i.e., a set of  $n$   $C^\infty$  vector fields  $e_1, \dots, e_n$  on  $V$  that are independent at each point of  $V$ .

**Theorem 4.1** (characterization of  $C^\infty$  tensors). If  $U$  is a framed open set in  $M$ , then  $\mathbf{T}^{p,q}U$  is isomorphic to  $\mathbf{T}^{p,q}\mathcal{X}_u$  in a natural way.

*Proof.* Let  $e_1, \dots, e_n$  be a  $C^\infty$  base field on  $U$ , and let  $w_1, \dots, w_n$  be the dual  $C^\infty$  1-forms on  $U$  (see problem 32). It is sufficient to illustrate the proof for  $\mathbf{T}^{0,p}\mathcal{X}_U$  where  $p > 0$ , since the other cases are analogous. Consider  $\alpha$  in  $\mathbf{T}^{0,p}$ , and let

$$\bar{\alpha} = \sum_{1 \leq i_j \leq n} \alpha(e_{i_1}, e_{i_2}, \dots, e_{i_p}) [w_{i_1} \otimes w_{i_2} \otimes \dots \otimes w_{i_p}],$$

be an element in  $\mathbf{T}^{0,p}\mathcal{X}_U$  defined by

$$\begin{aligned} & [\bar{\alpha}(X_1, \dots, X_p)](m) \\ &= \sum_{1 \leq i_j \leq n} [\alpha(e_{i_1}, \dots, e_{i_p})](m) [w_{i_1}(X_1(m)) w_{i_2}(X_2(m)) \dots w_{i_p}(X_p(m))]. \end{aligned}$$

where  $X_i$  are  $C^\infty$  fields on  $U$ . Then  $\alpha = \bar{\alpha}$  as elements of  $\mathbf{T}^{0,p}\mathcal{X}_U$ ,

for if  $X_i$  are in  $\mathcal{X}_U$ , then the function

$$\begin{aligned}\alpha(X_1, \dots, X_p) &= \alpha\left(\sum_{i_1=1}^n w_{i_1}(X_1)e_{i_1}, \sum_{i_2=2}^n w_{i_2}(X_2)e_{i_2}, \dots, \sum_{i_p=p}^n w_{i_p}(X_p)e_{i_p}\right) \\ &= \sum_{1 \leq i_j \leq n} w_{i_1}(X_1)w_{i_2}(X_2) \cdots w_{i_p}(X_p)\alpha(e_{i_1}, \dots, e_{i_p}),\end{aligned}$$

since  $\alpha$  is multilinear over  $\mathcal{F}_U$  and each  $w_j(X_i)$  is a function in  $\mathcal{F}_U$ .

But  $\bar{\alpha}$  is only an element of  $\mathbf{T}^{0,p}U$ , and notice  $[\bar{\alpha}(X_1, \dots, X_p)]$  depends only on the vectors  $X_1(m), \dots, X_p(m)$  and not on the fields  $X_1, \dots, X_p$ . Thus the map  $\alpha \mapsto \bar{\alpha}$  defines an isomorphism of  $\mathbf{T}^{0,p}\mathcal{X}_U$  onto  $\mathbf{T}^{0,p}U$ .  $\square$

One can “roughly” paraphrase the above theorem by saying that an  $\mathcal{F}_U$ -multilinear function on vector fields on  $U$  is actually a smooth piecing together of  $\mathbb{R}$  multilinear functions on  $\mathbf{T}_m M$  for each  $m$  in  $U$ .

**Corollary 4.2.** Let  $U$  be open in  $M$ . Let  $\alpha$  be a map that assigns to each framed open set  $V \subset U$  an element  $\alpha_V$  in  $\mathbf{T}^{q,p}\mathcal{X}_V$  with  $\bar{\alpha}_V = \bar{\alpha}_W$  in  $\mathbf{T}^{q,p}(V \cap W)$  for all open framed  $V$  and  $W$  contained in  $U$ . Then there is a unique tensor  $\alpha$  in  $\mathbf{T}^{p,q}U$  such that  $\alpha|_V = \bar{\alpha}_V$  for each framed open  $V \subset U$ . Moreover, if  $m$  in  $U$  and  $X_1, \dots, X_p$  are in  $\mathbf{T}_m M$  while  $z_1, \dots, z_q$  are in  $\mathbf{T}_m^* M$ , then

$$\alpha_m(X_1, \dots, X_p, z_1, \dots, z_q) = [a_V(\bar{X}_1, \dots, \bar{X}_p, \bar{z}_1, \dots, \bar{z}_q)](m), \quad (*)$$

*Proof.* Use  $(*)$  to define  $\alpha_m$  at any  $m$  in  $U$ . If  $W$  is any other framed open neighborhood of  $m$ , then  $\alpha_m = (\bar{\alpha}_W)_m = (\bar{\alpha}_W)_m$ , and one need only know the values of fields and forms at  $m$  in order to evaluate both of the tensors on the right.  $\square$

If the reader will become familiar with tensors and computations involving their linearity via some of the problems, then the above theorem and corollary should become more natural.

To close this chapter we study the maps on tensors induced by a  $C^\infty$  map  $f: M \rightarrow M'$ , where  $M$  is a  $C^\infty$   $n$ -manifold and  $M'$  is a  $C^\infty$   $n'$ -manifold. Because the Jacobian  $f_*$  maps vectors on  $M$  into vectors on  $M'$ , it induces a map  $f^*$  of covariant tensors (and forms) on  $M'$  into covariant tensors (and forms) on  $M$ . If  $g$  is in  $\mathbf{T}^{0,0}U' = \mathcal{F}_{U'}$ , for open  $U'$  on  $M'$ , then  $f^*(g) = g \circ f$  is a  $C^\infty$  real valued function in  $\mathcal{F}_U$  where  $U = \Omega^{-1}(U')$ . If  $\alpha$  is a  $p$ -co tensor at  $f(m)$  in  $M'$ , then  $(f^*\alpha)_m$  is the  $p$ -co tensor at  $m$  defined on  $X_1, \dots, X_p$  in  $\mathbf{T}_m M$  by

$$(f^*\alpha)_m(X_1, \dots, X_p) = \alpha_{f(m)}(f_*X_1, \dots, f_*X_p).$$

If  $\alpha$  is  $C^\infty$  on the open set  $U' \subset M'$ , then  $f^*\alpha$  is  $C^\infty$  on the open set  $f^{-1}(U') \subset M$ . In the next paragraph we prove this for a 1-form  $\alpha$  and leave the other cases to the problems.

Let  $\alpha$  be a  $C^\infty$  1-form on  $U'$ , let  $X$  be any  $C^\infty$  vector field on  $U$ , and we show that  $(f^*\alpha)(X)$  is a  $C^\infty$  function on  $U$ . Take  $m \in U$ , let  $x_1, \dots, x_n$  be a coordinate system about  $m$  with domain  $V \subset U$ , and let  $y_1, \dots, y_{n'}$  be a coordinate system about  $f(m)$  with domain  $V' \subset U'$ . Define  $C^\infty$  functions  $a_i$  on  $V$  and  $b_j$  on  $V'$  by

$$X = \sum_1^n a_i \left( \frac{\partial}{\partial x_i} \right) \text{ and } \alpha = \sum_1^{n'} b_j (\mathbf{d}y_j), \text{ where } \mathbf{d}y_r \left( \frac{\partial}{\partial y_s} \right) = \delta_{rs} = 0 \text{ or } 1,$$

according as  $r \neq s$  or  $r = s$ , respectively (see problem 32). Then on  $V$ ,

$$(f^*\alpha)(X) = \sum a_i (b_j \circ f) \frac{\partial (y_j \circ f)}{\partial x_i}$$

for  $i = 1, \dots, n$  and  $j = 1, \dots, n'$ , and since the right side is a  $C^\infty$  function on  $V$ ,  $(f^*\alpha)(X)$  is  $C^\infty$  on  $V$ , and hence  $f^*\alpha$  is  $C^\infty$  on  $U$ .

Finally, one checks that

- $f^*(\alpha_1 + \alpha_2) = f^*\alpha_1 + f^*\alpha_2$ , where  $\alpha_i$  are tensors of the same degree.
- $f^*(\gamma_1 \otimes \gamma_2) = f^*(\gamma_1) \otimes f^*(\gamma_2)$ , where  $\gamma_i$  are any covariant tensors.

- $f^*(\beta_1 \wedge \beta_2) = (f^*\beta_1) \wedge (f^*\beta_2)$ , where  $\beta_i$  are alternating covariant tensors.

Thus  $f^* : \Omega(M') \rightarrow \Omega(M)$  is a degree preserving exterior-algebra map of the  $C^\infty$  forms on  $M'$  into the  $C^\infty$  forms on  $M$ .

There are certain natural tensors on every manifold called *universal tensors*. These are mixed tensors that let the arguments “work on each other.” For example, let  $I$  be the 1, 1-tensor  $I(w, X) = w(X)$  for  $X$  in  $\mathbf{T}_m M$  and  $w$  in  $\mathbf{T}_m^* M$ . Another is the 2, 2-tensor  $E(w_1, w_2, X_1, X_2) = w_1(X_1)w_2(X_2)$ , etc.

The 1,1-tensors,  $\mathbf{T}^{1,1}W$ , over a vector space  $W$  have a natural interpretation, for there is a natural isomorphism of  $\mathbf{T}^{1,1}W$  with the group,  $\text{Hom}_{\mathbb{R}}(W, W)$ , of linear transformations of  $W$  into itself. If  $B$  is in  $\mathbf{T}^{1,1}W$ , then let  $\bar{B}$  be the linear map

$$\bar{B}(Z_i) = \sum_{j=1}^n B(w_j, Z_i)Z_j,$$

where  $Z_1, \dots, Z_n$  is a base of  $W$  with the dual base  $w_1, \dots, w_n$  of  $W^*$  (see problem 36).

## Problems

In these problems,  $W$  is an  $n$ -dim real vector space and  $M$  is a  $C^\infty$   $n$ -manifold.

32. Let  $e_1, \dots, e_n$  be a base of  $W$ . For  $i = 1, \dots, n$ , let  $w_i(e_j) = \delta_{ij}$ , where  $\delta_{ij} = 0$  if  $i \neq j$  and  $\delta_{ii} = 1$ . Show  $w_1, \dots, w_n$  is a base of  $W^*$ , and

$$\Theta = \sum_{i=1}^n \Theta(e_i)w_i \text{ for } \Theta \in W^*$$

33. (i) Let  $e_1, \dots, e_n$  be a base of  $W$ , and let  $w_1, \dots, w_n$  be the dual



base of  $W$ , ( $w_i(e_j) = \delta_{ij}$ ). If  $\alpha \in \mathbf{T}^{0,2}W$ , show

$$\alpha = \sum_{i,j=1}^n \alpha(e_i, e_j) w_i \otimes w_j;$$

thus  $\alpha$  is determined by its values on a basis.

(ii) If  $f_1, \dots, f_n$  another base, let  $\alpha(e_i, e_j) = a_{ij}$ ,  $\alpha(f_i, f_j) = b_{ij}$  and

$$f_j = \sum_{i=1}^n c_{ij} e_i. \text{ Show}$$

$$b_{ij} = \sum_{r,s=1}^n c_{ri} c_{sj} a_{rs}$$

.

34. (i) Show  $\mathbf{T}^{1,0}W$  is isomorphic to  $W$ .

(ii) Show  $\mathbf{T}^{p,1}W$  has dimension  $(p+q)n$ .

(iii) Show  $\Omega^p(W)$  has dimension  $\binom{n}{p} = \frac{n!}{p!(n-p)!}$ .

35. Let  $\alpha$  in  $\Omega^p(W)$  and  $\beta$  in  $\Omega^q(W)$ . If  $X_1, \dots, X_{p+q}$  in  $W$ , show

$$\alpha \wedge \beta(X_1, \dots, X_{p+q}) = \sum (-1)^\pi \alpha(X_{\pi_1}, \dots, X_{\pi_p}) \beta(X_{\pi_{p+1}}, \dots, X_{\pi_{p+q}})$$

where the sum is over all shuffle permutations  $\pi$  for  $p$  and  $q$ , i.e., if

$1 \leq i < j \leq p$  or  $p+1 \leq i < j \leq p+q$ , then  $\pi_i < \pi_j$ .

36. (i) Show  $\mathbf{T}^{1,1}W$  is isomorphic to  $\text{Hom}_{\mathbb{R}}(W, W)$ , the set of all  $\mathbb{R}$ -linear maps of  $W$  into  $W$ , via the above map  $B \mapsto \bar{B}$ , and show this map is independent of the base  $Z_i$ .

(ii) Show the universal tensor  $I$  in  $\mathbf{T}^{1,1}W$  corresponds to the identity map on  $W$ .

37. If  $e_1, \dots, e_n$  is a  $C^\infty$  base field on  $U$  in  $M$ , and  $w_1, \dots, w_n$  is the set of dual 1-forms on  $U$ , show each  $w_i$  is  $C^\infty$  on  $U$ .

38. Let  $f$  be in  $C^\infty(M, \mathbb{R})$ . For  $p$  in  $M$  and  $X$  in  $M_p$ , let  $(df)_p X = Xf$ .<sup>2</sup>

(i) Show  $(df)_p$  is a 1-form at  $p$ .

(ii) If  $x_1, \dots, x_n$  is a coordinate system with domain  $U$ , show  $dx_1, \dots, dx_n$  is the dual base to  $\frac{\partial}{\partial x_1} \dots \frac{\partial}{\partial x_n}$  and

$$df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i \text{ on } U.$$

(iii) Show  $df$  is  $C^\infty$  on  $M$ .

(iv) Show  $d(f+g) = df+dg$  and  $d(fg) = f dg + g df$  for  $g$  in  $C^\infty(M, \mathbb{R})$ .

(v) If  $y_1, \dots, y_n$  is also a coordinate system on  $U$  and

$$w = \sum_1^n a_i dx_i = \sum_1^n b_j dy_j,$$

show

$$b_j = \sum_{i=1}^n \frac{\partial x_i}{\partial y_j}.$$

(vi) If  $\alpha = \sum a_{ij} dx_i \wedge dx_j = \sum b_{ij} dy_i \wedge dy_j$ , find expressions for  $a_{ij}$  in terms of  $b_{ij}$

(vii) If  $g$  in  $C^\infty(M, M')$ , show  $g^* \circ d = d \circ g^*$  on  $\Omega^0(M')$ .

39. If  $w_1, \dots, w_n$  is a base of  $\mathbf{T}_p^*M$ , show there is a coordinate system  $x_1, \dots, x_n$  about  $p$  with  $(dx_i)_p = w_i$  for all  $i$ .

40. Let  $\mathbf{T}^*M$  be the set of all ordered pairs  $(m, w)$  for  $m \in M$  and  $w \in \mathbf{T}_m^*M$ . Let  $\pi: \mathbf{T}^*M \rightarrow M$  by  $\pi(m, w) = m$ . Let  $x_1, \dots, x_n$  be a coordinate system on  $M$  with domain  $U$ , and define functions  $q_1, \dots, q_n, p_1, \dots, p_n$  on  $\pi^{-1}(U)$  by  $q_j = x_j \circ \pi$  and  $w = \sum p_j(m, w) dx_j$ . With these coordinate functions  $\mathbf{T}^*M$  becomes a  $C^\infty$   $2n$ -manifold called the cotangent bundle of  $M$ . The fundamental 1-form  $W$  on  $M$  is defined by  $W_{(m,w)} = \pi^*w$ .

---

<sup>2</sup>This defines a map  $d: \Omega^p(A) \rightarrow \Omega^{p+1}(A)$  called the exterior derivative. See sections 5.2 and 7.1.

- (a) Show  $W = \sum p_i dq_i$  on  $\pi^{-1}(U)$ .
- (b) Show  $d$  is a non-degenerate 2-form; i.e.,  $dW(X, Y) = 0$  for all  $Y$  implies  $X = 0$ . The forms  $W$  and  $dW$  are fundamental in classical mechanics (see [Mac04]).
41. Let  $X$  be tangent to  $B(M)$  at  $b = (m: e_1, \dots, e_n)$ . Let  $\bar{w}_i(X)$  be real numbers such that  $X = \sum \bar{w}_i(X) e_i$ . Show  $w_i$  are  $C^\infty$  1-forms on  $B(M)$  (see 22).



## 5. Connexions

This chapter is a study of a general connexion on a  $C^\infty$  manifold, the concepts belonging to the connexion, and the different ways of defining the connexion. These connexions are historically called affine or linear connexions on a manifold. The generalization to connexions in principal fiber bundles is sketched in section 5.5, but these generalizations will not be focused upon in these notes.

### 5.1 Invariant Viewpoint

The approach to connexions that follows is due to Koszul and is found in [KN63] and the first chapter of [Hel12]. The definition was motivated in 2.1.

Let  $M$  be a  $C^\infty$   $n$ -manifold. A *connexion, infinitesimal connexion or covariant differentiation* on  $M$  is an operator  $D$  that assigns to each pair of  $C^\infty$  fields  $X$  and  $Y$ , with domain  $A$ , a  $C^\infty$  field  $\mathbf{D}_X Y$  with domain  $A$ ; and if  $Z$  is a  $C^\infty$  field on  $A$  while  $f$  is a  $C^\infty$  real valued function on  $A$ , then  $D$  satisfies the following four properties:

$$(1) \mathbf{D}_X(Y + Z) = \mathbf{D}_X Y + \mathbf{D}_X Z$$

$$(2) \mathbf{D}_{(X+Y)}(Z) = \mathbf{D}_X Z + \mathbf{D}_Y Z$$

$$(3) \mathbf{D}_{(fX)} Y = f \mathbf{D}_X Y$$

$$(4) \mathbf{D}_X(fY) = (Xf)Y + f \mathbf{D}_X Y.$$

These properties imply the vector  $(\mathbf{D}_X Y)_m$ , at a point  $m \in M$ , depends only on  $X_m$  and the values of  $Y$  on some curve that fits  $X_m$ .

For, let  $e_1, \dots, e_n$  be a  $C^\infty$  base field about  $m$ , let  $X_m = \sum_1^n a_i(m)(e_i)_m$

and  $Y = \sum_1^n b_j e_j$  on the domain of the base field (intersected with the domain of  $Y$ ). Then

$$\begin{aligned} (\mathbf{D}_X Y)_m &= \left[ \mathbf{D}_X \left( \sum_j b_j e_j \right) \right]_m \\ &= \sum_j \left[ (X_m b_j)(e_j)_m + b_j(m) \sum_i a_i(m) (\mathbf{D}_{e_i} e_j)_m \right]. \end{aligned}$$

Thus  $a_i(m)$ ,  $b_j(m)$  and  $X_m b_j$  determine  $\mathbf{D}_X Y$  completely if the fields  $\mathbf{D}_{e_i} e_j$  are known (see section 5.2)

The existence of many manifolds with connexions has been illustrated by the natural induced connexions on hypersurfaces of  $\mathbb{R}^n$ .

Let  $\sigma$  be a curve in  $M$  with tangent field  $T$ . A  $C^\infty$  vector field  $Y$  on  $\sigma$  is *parallel along*  $\sigma$  iff  $\mathbf{D}_T Y = 0$  on  $\sigma$ . The curve  $\sigma$  is a *geodesic* iff  $\mathbf{D}_T T = 0$  on  $\sigma$ . Thus a curve is a geodesic iff its tangent field is a parallel field along the curve. The following two theorems give the existence of parallel fields and geodesics. The domain of an index of summation is always  $1, \dots, n$  unless otherwise specified.

**Theorem 5.1.** Let  $\sigma$  be a curve on  $[a, b]$  with tangent  $T$ . For each vector  $Y$  in  $\mathbf{T}_{\sigma(a)}M$  there is a unique  $C^\infty$  field  $Y(t)$  on  $\sigma$  such that  $Y(a) = Y$  and the field  $Y(t)$  is parallel along  $\sigma$ . The mapping  $P_{a,t}: \mathbf{T}_{\sigma(a)}M \rightarrow \mathbf{T}_{\sigma(t)}M$  by  $P_{a,t}(Y) = Y(t)$  is a linear isomorphism which is called *parallel translation along*  $\sigma$  from  $\sigma(a)$  to  $\sigma(t)$ .

**Proof.** Let  $x_1, \dots, x_n$  be a coordinate system about  $\sigma(a)$  with domain  $U$ , and let  $X_1, \dots, X_n$  be the associated vector fields. We define  $C^\infty$  functions  $\Gamma_{jk}^i$  on  $U$  by  $\mathbf{D}_{X_k} X_j = \sum_i \Gamma_{jk}^i X_i$ . Let  $\sigma$  map the domain  $[a, b_1]$  into  $U$ . If  $Y(t)$  is a field on  $\sigma$  with domain  $[a, b_1]$  then define functions

$a_i(t)$  on this domain by  $Y(t) = \sum a_i(t)X_i(\sigma(t))$ . Let  $g_i(t) = x_i \circ \sigma(t)$  on  $[a, b_1]$ , so  $T(t) = \sum g'_j(t)X_j(\sigma(t))$ , where  $g'_j(t) = \frac{dg_j}{dt}$ . If  $Y(t)$  is parallel along  $\sigma$ , then

$$0 = \mathbf{D}_T Y = \sum_i \left[ a'_i X_i + a_i \sum_j g'_j \Gamma_{ij}^k X_k \right].$$

Thus  $Y(t)$  parallel along  $\sigma$  iff

$$\frac{da_k}{dt} + \sum_{i,j} a_i \frac{dg_j}{dt} \Gamma_{ij}^k = 0 \tag{5}$$

for  $k = 1, \dots, n$  and for  $t \in [a, b_1]$ . The condition  $Y(a) = Y$  defines  $n$  initial values  $a_i(a)$ , and the theory of ordinary differential equations then gives a unique set of  $C^\infty$  functions  $a_i(t)$ , satisfying the above equations on the whole domain  $[a, b_1]$ , since the equations are linear. This defines the parallel field  $Y(t)$ .

For  $t \in [a, b_1]$ , the map  $P_{a,t}$  is linear because of the linearity of the equations 5.

If  $t$  is any number in  $[a, b_1]$ , we obtain  $P_{a,t}$  by covering the compact set  $\sigma([a, t])$  by a finite number of coordinate neighbourhoods and parallel translating through each neighbourhood via solutions of the systems 5. □

**Theorem 5.2.** Let  $m \in M$ ,  $X \in \mathbf{T}_m M$ . Then for any real number  $b$  there exists a real number  $r > 0$  and a unique curve  $\sigma$ , defined on  $[b - r, b + r]$  such that  $\sigma(b) = m$ ,  $T_\sigma(b) = X$ , and  $\sigma$  a geodesic.

*Proof.* Using the notation of the above proof, we must find  $C^\infty$  functions  $g_i(t)$  that satisfy the second order differential system,

$$\frac{d^2 g_k}{dt^2} + \sum_{i,j} \Gamma_{ij}^k \frac{dg_i}{dt} \frac{dg_j}{dt} = 0 \tag{6}$$

with initial conditions  $g_i(b) = x_i(m)$  and  $X = \sum g'_i(b)X_i$ . The theory of ordinary differential equations provides us with the  $r > 0$  and the

functions  $g_i(t)$ . □

The existence and uniqueness theory of ordinary differential equations will actually give us more than the conclusion of the above theorem. In particular, if we let  $\sigma(t; m, X, b)$  be the curve provided by the theorem, then the mapping  $\sigma$  is actually  $C^\infty$  with respect to all its parameters  $t, m, X$  and  $b$ .

The *torsion tensor* of a connexion  $\mathbf{D}$  is a vector valued tensor  $\text{Tor}$  that assigns to each pair of  $C^\infty$  vectors  $X$  and  $Y$ , with domain  $A$ , a  $C^\infty$  vector field  $\text{Tor}(X, Y)$ , with domain  $A$ , by

$$\text{Tor}(X, Y) = \mathbf{D}_X Y - \mathbf{D}_Y X - [X, Y]. \quad (7)$$

One easily checks that for  $Z \in \mathcal{X}_A$  and  $f \in \mathcal{F}_A$

- $\text{Tor}(X, Y) = -\text{Tor}(Y, X)$ ,
- $\text{Tor}(X + Y, Z) = \text{Tor}(X, Z) + \text{Tor}(Y, Z)$ ,
- $\text{Tor}(fX, Y) = f\text{Tor}(X, Y)$ .

Thus the value of  $\text{Tor}(X, Y)$  at a point  $m$  depends only on  $\mathbf{T}_m X$  and  $\mathbf{T}_m Y$ , and not on the fields  $X$  and  $Y$ , by the theorem at the end of Chapter 4. If more than one connexion enters the discussion, we write  $\text{Tor}_\mathbf{D}$  for the torsion of the connexion  $\mathbf{D}$ . If  $\text{Tor}_\mathbf{D} \equiv 0$ , then we say that  $\mathbf{D}$  is *symmetric*, or *torsion free*.

As far as we know, there is no nice motivation for the word “torsion” to describe the above tensor. In particular, it has nothing to do with the “torsion of a space curve”.

The following definition of curvature has been motivated in section 2.4.

The *curvature tensor* of a connexion  $\mathbf{D}$  is a linear transformation valued tensor  $R$  that assigns to each pair of vectors  $X$  and  $Y$  at  $m$  a linear transformation  $R(X, Y)$  of  $\mathbf{T}_m M$  into itself, we define  $R(X, Y)Z$  by imbedding  $X, Y$ , and  $Z$  in  $C^\infty$  fields about  $m$  and setting

$$R(X, Y)Z = (\mathbf{D}_X \mathbf{D}_Y Z - \mathbf{D}_Y \mathbf{D}_X Z - \mathbf{D}_{[X, Y]} Z)_m. \quad (8)$$



Again we check linearity over the ring of  $C^\infty$  functions as coefficients on the right to determine the tensor character of  $R$ . Here,  $R(X, Y)Z = -R(Y, X)Z$ , and if  $f$  is  $C^\infty$ , then

$$\begin{aligned} R(fX, Y)Z &= f\mathbf{D}_X\mathbf{D}_Y Z - (Yf)\mathbf{D}_X Z - f\mathbf{D}_Y\mathbf{D}_X Z + (Yf)\mathbf{D}_X Z - f\mathbf{D}_{[X, Y]}Z \\ &= fR(X, Y)Z. \end{aligned}$$

Also

$$\begin{aligned} R(X, Y)(fZ) &= \mathbf{D}_X[(Yf)Z + f\mathbf{D}_Y Z] - \mathbf{D}_Y[(Xf)Z + f\mathbf{D}_X Z] - ([X, Y]f)Z - f\mathbf{D}_{[X, Y]}Z \\ &= (XYf)Z + (Yf)\mathbf{D}_X Z + (Xf)\mathbf{D}_Y Z + f\mathbf{D}_X\mathbf{D}_Y Z - (YXf)Z \\ &\quad - (Xf)\mathbf{D}_Y Z - (Yf)\mathbf{D}_X Z - f\mathbf{D}_Y\mathbf{D}_X Z - ([X, Y]f)Z - f\mathbf{D}_{[X, Y]}Z \\ &= fR(X, Y)Z. \end{aligned}$$

The linearity of  $R(X, Y)Z$  with respect to addition (in each of its variables) is trivial to check.

The tensor nature of the torsion and curvature will again be verified in section 5.3 with exhibition of the classical coordinate representations of these tensors.

The concept of a “connexion-preserving” map follows naturally. Let  $M$  and  $M'$  be  $C^\infty$  manifolds with connexions  $D$  and  $D'$ , respectively. A  $C^\infty$  map

$f: M \rightarrow M'$  is *connexion preserving* if  $f_*(\mathbf{D}_X Y) = \mathbf{D}'_{f_*X}(f_*Y)$  for all vectors  $X$  and fields  $Y$ . Note the right side is well-defined since  $f_*Y$  is a well-defined field on some curve that fits  $f_*X$ . A  $C^\infty$  map  $f: M \rightarrow M'$  is *geodesic preserving* if  $f \circ g$  is a geodesic in  $M'$  for each geodesic  $g$  in  $M$ . Trivially, a connexion-preserving map is geodesic preserving.

**Theorem 5.3.** Let  $f$  be a diffeomorphism of  $M$  onto  $M'$ , and let  $D'$  be a connexion on  $M'$ . Then there is a unique connexion  $D$  on  $M$  for which  $f$  is connexion preserving.

*Proof.* Take  $X$  in  $\mathbf{T}_m M$  and let  $Y$  be a field around  $m$ . Since  $f$  is a

diffeo,  $f_*Y$  is a field around  $f(m)$ . Define  $\mathbf{D}_X Y = f_*^{-1}(D'_{f_*X} f_*Y)$ . The verification that  $\mathbf{D}$  is a connexion is left as an exercise.  $\square$

If every geodesic  $g(t)$  can be extended so it is a geodesic for all  $t \in \mathbb{R}$ , then the connexion  $\mathbf{D}$  is complete.

## 5.2 Cartan Viewpoint

For local problems concerning a connexion, one can transform the properties of  $\mathbf{D}$  to certain properties of differential forms. By using fiber bundles associated with a manifold, one can also study global problems via differential forms. We develop the local viewpoint here.

Let  $\mathbf{D}$  be a connexion on an  $n$ -manifold  $M$ , and fix  $\mathbf{D}$  and  $M$  throughout this section. Let  $U$  be a fixed open set (perhaps a coordinate domain) in  $M$ , and let  $e_1, \dots, e_n$  be a fixed base field of independent  $C^\infty$  vectors on  $U$ . Let  $w_1, \dots, w_n$  be the  $C^\infty$  1-forms on  $U$  which are the dual base to  $e_1, \dots, e_n$  at each point of  $U$ . Define  $n^2$  connexion 1-forms  $w_{ij}$  on  $U$  which are associated with  $\mathbf{D}$  and the base field by

$$\mathbf{D}_X e_j = \sum_{i=1}^n w_{ij}(X) e_i. \quad (9)$$

The  $w_{ij}$  are linear by property (2) of the connexion  $\mathbf{D}$ , and  $w_{ij}$  are  $C^\infty$ , since if  $X$  a  $C^\infty$  field on  $U$ , then  $\mathbf{D}_X e_j$  is a  $C^\infty$  field, so  $w_{ij}(X) = w_i(\mathbf{D}_X e_j)$  is a  $C^\infty$  function.

The torsion and curvature tensors may also be expressed via differential forms associated with the base field. Define 2-forms  $T_i$  and  $R_{ij}$  on  $U$  by

$$T(X, Y) = \sum_{i=1}^n T_i(X, Y) e_i \quad (10)$$

$$R(X, Y) e_j = \sum_{i=1}^n R_{ij}(X, Y) e_i \quad (11)$$

where the properties of an alternating tensor sum are checked for  $T_i$  and  $R_{ij}$  via the properties of  $T$  and  $R$ .

The forms  $w_i, w_{ij}, T_i$  and  $R_{ij}$  are related by the Cartan structural equations which are equivalent to the definition of the torsion and curvature tensors. We merely express everything in terms of the base field. Let  $X$  and  $Y$  be  $C^\infty$  fields on  $U$ . Then,

$$\begin{aligned} T_i(X, Y)e_i &= \mathbf{D}_X Y - \mathbf{D}_Y X - [X, Y] \\ &= \mathbf{D}_X \left( \sum w_j(Y) e_j \right) - \mathbf{D}_Y \left( \sum w_j(X) e_j \right) - \sum w_j([X, Y]) e_j \\ &= \sum (Xw_j(Y) - Yw_j(X) - w_j[X, Y]) e_j + (w_j(Y)w_{ij}(X) - w_j(X)w_{ij}(Y)) e_i \end{aligned}$$

Equating components,

$$T_i(X, Y) - \left( \sum w_{ij} \wedge w_j \right)(X, Y) = Xw_i(Y) - Yw_i(X) - w_i[X, Y]$$

Since the expression on the left is a 2-form, so is the expression on the right (taken as a whole), and indeed, it is the exterior derivative  $\mathbf{d}w_i$  of  $w_i$  evaluated on  $X$  and  $Y$ . With this motivation we define the exterior derivative operator  $d$  on 1-forms and functions (0-forms) as follows.

For a  $C^\infty$  function  $f$  with domain  $A$ , let  $\mathbf{d}f(X) = Xf$ ; thus  $\mathbf{d}f$  is a  $C^\infty$  1-form on  $A$ . Let  $w$  be any  $C^\infty$  1-form with domain  $A$ . Then  $\mathbf{d}w$  is a  $C^\infty$  2-form with domain  $A$ , defined on  $C^\infty$  fields  $X, Y$  on  $A$  by

$$\mathbf{d}w(X, Y) = Xw(Y) - Yw(X) - w[X, Y] \tag{12}$$

We leave it to the reader to check that the right side is linear in each slot over the ring of  $C^\infty$  functions on  $A$ , and hence that  $\mathbf{d}w(X_m, Y_m)$  is defined for  $m$  in  $A$  independent of the fields  $X$  and  $Y$ .

If  $f$  is a  $C^\infty$  function on  $A$ , then  $\mathbf{d}^2 f - \mathbf{d}(\mathbf{d}f) = 0$ . To see this, let

$X$  and  $Y$  be  $C^\infty$  fields on  $A$ ; then,

$$\begin{aligned} \mathbf{d}^2 f(X, Y) &= X \mathbf{d}f(Y) - Y \mathbf{d}f(X) - \mathbf{d}f[X, Y] \\ &= XYf - YXf - [X, Y]f = 0 \end{aligned}$$

Also note that if  $x_1, \dots, x_n$  a coordinate system on  $A$ , then  $\mathbf{d}x_1, \dots, \mathbf{d}x_n$  is the dual base to  $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}$ , since  $dx_i \left( \frac{\partial}{\partial x_j} \right) = \frac{\partial x_i}{\partial x_j} = \delta_{ij}$  (the Kronecker delta).

Now we can write the first Cartan structural equation  
Cartan structural equations

$$\mathbf{d}w_i = - \sum_{j=1}^n w_{ij} \wedge w_j + T_i \quad (13)$$

By a comutation involving the definition of  $R(X, Y)$ , which is completely analogous to the above computation, one obtains the second Cartan structural equation,

$$\mathbf{d}w_{ij} = - \sum_{k=1}^n w_{ik} \wedge w_{kj} + R_{ij} \quad (14)$$

These equations provide an alternate proof of the tensor character of  $T$  and  $R$ , since they show that  $T_i$  and  $R_{ij}$  are 2-forms.

### 5.3 Coordinate Viewpoint

Let  $U$  be a coordinate neighborhood, and let  $X_1, \dots, X_n$  be the coordinate base field associated with the system  $x_1, \dots, x_n$  on  $U$ . Then  $w_i = \mathbf{d}x_i$  and the associated forms  $w_{ij}, T_i$ , and  $R_{ij}$  define functions

$\Gamma_{jk}^i$ ,  $T_{ijk}$ , and  $R_{jkh}^i$ , respectively, by

$$\begin{aligned} \Gamma_{jk}^i = w_{ij}(X_k) & \quad \text{so} & \quad w_{ij} = \sum_k \Gamma_{jk}^i \mathbf{d}x_k \\ T_{ijk} = T_i(X_j, X_k) & \quad \text{so} & \quad T_i = \sum_{jk} T_{ijk} \mathbf{d}x_j \otimes \mathbf{d}x_k, \\ R_{jkh}^i = R_{ij}(X_k, X_h) & \quad \text{so} & \quad R_{ij} = \sum_{kh} R_{jkh}^i \mathbf{d}x_k \otimes \mathbf{d}x_h \end{aligned}$$

From the structural equations, we have

$$T_{ijk} = \left( \mathbf{d}^2 x_i + \sum_{r,s} \Gamma_{rs}^i \mathbf{d}x_s \wedge \mathbf{d}x_t \right) (X_j, X_k) = \Gamma_{kj}^i - \Gamma_{jk}^i,$$

since  $\mathbf{d}^2 x_i = 0$ , and

$$\begin{aligned} R_{jkh}^i &= \left( \mathbf{d}w_{ij} + \sum_{r,s,t} (\Gamma_{rs}^i \mathbf{d}x_s) \wedge (\Gamma_{jt}^r \mathbf{d}x_t) \right) (X_k, X_h) \\ &= X_k w_{ij}(X_h) - X_h w_{ij}(X_k) + \sum_r (\Gamma_{rk}^i \Gamma_{jh}^r - \Gamma_{rh}^i \Gamma_{jk}^r) \\ &= \frac{\partial \Gamma_{jh}^i}{\partial x_k} - \frac{\partial \Gamma_{jk}^i}{\partial x_h} + \sum_r (\Gamma_{rk}^i \Gamma_{jh}^r - \Gamma_{rh}^i \Gamma_{jk}^r) \end{aligned}$$

which are the classical coordinate components of these tensors.

## 5.4 Difference Tensor of Two Connexions

The reference of this section is [WS60]. Let  $M$  be a  $C^\infty$  manifold, and let  $\mathbf{D}$  and  $\bar{\mathbf{D}}$  be connexions on  $M$ . For fields  $X$  and  $Y$  we define the *difference tensor*  $B(X, Y) = \bar{\mathbf{D}}_X Y - \mathbf{D}_X Y$ . The linearity of  $B$  in the first slot is trivial from properties of the connexions (namely, (2) and (3)). To check the second slot, let  $f$  be  $C^\infty$  on the domain of  $X$  and  $Y$ ; then

$$B(X, fY) = (Xf)Y + f\bar{\mathbf{D}}_X Y - (Xf)Y - f\mathbf{D}_X Y = fB(X, Y)$$

Let  $B(X, Y) = S(X, Y) + A(X, Y)$  be the stanard decomposition of a bilinear tensor into symmetric and skew-symmetric pieces; i.e,

$$S(X, Y) = \frac{1}{2}[B(X, Y) + B(Y, X)]$$

and

$$A(X, Y) = \frac{1}{2}[B(X, Y) - B(Y, X)]$$

Actually, we can express  $A$  in terms of the torsion tensors  $T$  and  $\bar{T}$  of  $\mathbf{D}$  and  $\bar{\mathbf{D}}$ , respectively, for

$$\begin{aligned} 2A(X, Y) &= \bar{\mathbf{D}}_X Y - \mathbf{D}_X Y - \bar{\mathbf{D}}_Y X + \mathbf{D}_Y X \\ &= \bar{T}(X, Y) + [X, Y] - T(X, Y) - [X, Y] \\ &= \bar{T}(X, Y) - T(X, Y) \end{aligned}$$

**Theorem 5.4.** The following statements are equivalent:

- (a) The connections  $\mathbf{D}$  and  $\bar{\mathbf{D}}$  have the same geodesics.
- (b)  $B(X, X) = 0$  for all vectors  $X$ .
- (c)  $S = 0$
- (d)  $B = A$

*Proof.*

- (a)  $\implies$  (b): Take  $X$  at  $m \in M$  and let  $g$  be the geodesic with initial vector  $X$ . Extend  $X$  along  $g$  by letting  $X$  be the tangent to  $g$ ; then

$$p = B(X, X) = \bar{\mathbf{D}}_X X - \mathbf{D}_X X = 0 - 0,$$

since  $g$  is a geodesic for both connections.

- (b)  $\implies$  (a): Let  $g$  be a geodesic for  $\mathbf{D}$  with tangent field  $X$ ; then  $\bar{\mathbf{D}}_X X = B(X, X) + \mathbf{D}_X X = 0$  on  $g$ ; hence  $g$  is a geodesic for  $\bar{\mathbf{D}}$

- (b)  $\iff$  (c): Since  $S$  is symmetric, it is determined by its diagonal values  $S(X, X)$ , and  $B(X, X) = 0 \iff S(X, X) = 0$

(c)  $\iff$  (d): For  $B = S + A$ . □

*Theorem 5.5.* The connexions  $\mathbf{D}$  and  $\bar{\mathbf{D}}$  are equal iff they have the same geodesics and the same torsion tensors.

*Proof.* That the first part implies the second is trivial. Conversely if the geodesics are the same, then  $S = 0$ , and if the torsion tensors are equal, then  $A = 0$ ; hence  $B = 0$  and  $\mathbf{D} = \bar{\mathbf{D}}$  □

*Theorem 5.6.* Given a connexion  $\bar{\mathbf{D}}$  on  $M$ , there is a unique connexion  $\mathbf{D}$  having the same geodesics as  $\bar{\mathbf{D}}$  and zero torsion.

*Proof.* Let  $\mathbf{D}_x Y = \bar{\mathbf{D}}_x Y - \frac{1}{2}\bar{T}(X, Y)$ . It is trivial to check that  $\mathbf{D}$  satisfies the required properties to define a connexion. Here  $B = \frac{1}{2}\bar{T} = A$ , since a torsion tensor is skew-symmetric; thus  $S = 0$ , so  $\mathbf{D}$  and  $\bar{\mathbf{D}}$  have the same geodesics. Moreover,  $T = \bar{T} - 2A = 0$ , so  $\mathbf{D}$  has zero torsion. The uniqueness follows from the preceding theorem. □

Thus if we partition connexions into equivalence classes by placing two connexions with the same geodesics in the same class, then in each class there exists a unique torsion-free (zero torsion) connexion. Moreover, given any connexion  $\mathbf{D}$  and any skew-symmetric vector-valued 2-covariant tensor  $\bar{T}$ , there exists a connexion with torsion tensor  $\bar{T}$  and the same geodesics as  $\mathbf{D}$ . From the above proof we have  $\bar{T}(X, Y) = 2(\bar{\mathbf{D}}_x Y - \mathbf{D}_x Y)$ , which provides a geometric interpretation of the torsion tensor of a connexion as measuring the difference between covariant differentiation in the given connexion and covariant differentiation in the torsion-free connexion with the same geodesics.

## 5.5 Bundle Viewpoint

In this section we define a connexion on the bundle of bases over a manifold and sketch a proof of the equivalence of such a definition with our previous viewpoints. This is the fourth (and

last) viewpoint we consider. The bundle viewpoint provides a natural “jumping off” for generalizations to connexions in all kinds of bundles, and much of the research in differential geometry at this time uses these concepts. For more details the reader is referred to the [BC11] or [KN63].

Throughout this section let  $M$  be a  $C^\infty n$ -manifold, let  $B = B(M)$  be the bundle of bases over  $M$  (see problem 22), and let  $\pi : B \rightarrow M$  be the natural projection map. If  $\mathbf{D}$  is a connexion on  $M$ , then by integrating ordinary differential equations (5 above), we can parallel translate the tangent space along curves in  $M$ . If  $b = (m; e_1, \dots, e_n)$  is in  $B$  and  $\sigma$  is a curve in  $M$  with  $\sigma(0) = m$ , then by parallel translation we define a  $C^\infty$  curve  $\bar{\sigma}(t) = (\sigma(t); e_1(t), \dots, e_n(t))$  in  $B$ , where  $e_i(t)$  is the parallel translate of  $e_i = e_i(0)$  along  $\sigma$  to  $\sigma(t)$ . Since  $\pi \circ \bar{\sigma} = \sigma$ , we say  $\bar{\sigma}$  is a “lift of  $\sigma$ ”, or  $\bar{\sigma}$  “lies over  $\sigma$ ” and since  $\bar{\sigma}$  reads off a parallel base, we say  $\bar{\sigma}$  is a “horizontal” curve in  $B$ . Thus a connexion  $\mathbf{D}$  on  $M$  yields unique “horizontal lifts” of  $C^\infty$  curves in  $M$ . The bundle definition of a connexion gives an independent method for defining “horizontal lifts” (of curves in  $M$ ) with the correct properties.

Recall at each point  $b \in B$  we defined the subspace of vertical vectors

$V_b = \{X \in B_b : \pi_*(X) = 0\}$ . A connexion on  $B$  is a mapping  $H$  that assigns to each  $b$  in  $B$ , a subspace  $H_b$  of  $B_b$  such that:

- (1)  $H_b \cap V_b = 0$  and  $\pi_*|_{H_b}$  is an isomorphism of  $H_b$  onto  $\mathbf{T}_{\pi(b)}M$  (hence  $H_b$  is  $n$ -dimensional).
- (2)  $(R_g)_*(H_b) = H_{bg}$  for all  $g$  in  $\mathbf{GL}(n, \mathbb{R})$ .
- (3)  $H$  is  $C^\infty$ ; i.e., for each  $b$  in  $B$  there is a neighborhood  $U$  and a set of  $n$  independent  $C^\infty$  vector fields  $E_1, \dots, E_n$  on  $U$  that give a base for  $H_{b'}$  for every  $b'$  in  $U$ .

If  $X$  is in  $H_b$ , we say  $X$  is a horizontal vector. Property (1) implies for each  $X$  in  $B_b$  there is a unique decomposition  $X = X_H + X_V$  with



$X_H \in H_b$  and  $X_V \in V_b$ , and property (3) implies if  $X$  is  $C^\infty$  then  $X_H$  and  $X_V$  are  $C^\infty$  fields. If  $X$  is a  $C^\infty$  field with domain  $U$  in  $M$ , then there is a unique  $C^\infty$  horizontal field  $\bar{X}$  on  $\bar{U} = \pi^{-1}(U)$  with  $\pi_*(\bar{X}) = X_{\pi(b)}$  for all  $b$  in  $\bar{U}$ .

Having the existence of “horizontal lifts” for vector fields, one can “horizontally” lift curves in a natural way. Thus if  $\sigma$  is a curve in  $M$  with tangent  $T$  (non-vanishing), extend  $T$  to a  $C^\infty$  field in a neighborhood  $U$  of a univalent part of  $\sigma$ , lift  $T$  to a horizontal field  $\bar{T}$  on  $\bar{U}$ , and take integral curves of  $\bar{T}$  to find horizontal lifts of  $\sigma$ . The parallel translation so defined will be independent of the base (the starting point for  $\bar{\sigma}$ ) by property (2); i.e., if  $\bar{\sigma}$  is horizontal (has a horizontal tangent), then  $R_g \circ \bar{\sigma}$  is also horizontal.

There is a dual viewpoint involving differential forms. To motivate it, let  $H$  be a connexion as described above and notice at each  $b = (m; e_1, \dots, e_n)$  in  $B$  we can define a unique horizontal field  $E_i(b)$  with  $\pi_*(E_i(b)) = e_i$  by (1). The fields  $E_1, \dots, E_n$  are global independent horizontal  $C^\infty$  fields on  $B$ . Together with the natural vertical fields  $E_{11}, \dots, E_{nn}$ , we get a global base field on  $B$ . Let  $\bar{w}_1, \dots, \bar{w}_n, \bar{w}_{11}, \dots, \bar{w}_{nn}$  be the dual 1-forms to this base (where  $\bar{w}_1, \dots, \bar{w}_n$  are the natural 1-forms of problem 41). Then if  $X \in B_b, X_V = \sum_{i,j=1}^n \bar{w}_{ij}(X)(E_{ij})_b$ . If one knows  $X_V$ , then, of course,  $X_H = X - X_V$ . Thus giving  $X_H$  (or giving  $H$ ) is equivalent to giving “vertical projections” at each point in  $B$ . Thus a set of connexion 1-forms  $\bar{w}_{ij}$  (for  $i, j = 1, \dots, n$ ) on  $B$  is a set of 1-forms such that

(1')  $\bar{w}_{ij}|_{V_b}$  form a dual base to  $E_{ij}$  at all  $b$  in  $B$ ,

(2')  $\bar{w}_{ij}((R_g)_*X) = \sum_{t_s=1}^n g_{it}^{-1} \bar{w}_{ts}(X) g_{sj}$  for all  $X$  in  $B_b$ ,

(3')  $\bar{w}_{ij}$  are  $C^\infty$  for all  $i$  and  $j$ .

That the definition of a connexion on  $B$  in terms of  $H$  or in terms of  $\bar{w}_{ij}$  is equivalent is left as a problem.

Notice that the  $\bar{w}_{ij}$  can be used to define a Lie algebra (of  $\text{GL}(n, \mathbb{R})$ ) valued 1-form  $\bar{w}$  by  $\bar{w}(X) = \sum_{i,j=1}^n \bar{w}_{ij}(X)X_{ij}$ , where the  $X_{ij}$  are the canonical left invariant fields on  $\text{GL}(n, \mathbb{R})$  (see problem 21).

Finally, we connect with the Cartan viewpoint. Let  $e_1, \dots, e_n$  be a base field on the open set  $U$  in  $M$ . Define a  $C^\infty$  map  $f: U \rightarrow B$  by  $f(m) = (m; (e_1)_m, \dots, (e_n)_m)$  for  $m$  in  $U$ . Since  $\pi \circ f$  is the identity of  $U$ , we call  $f$  a section over  $U$ . Let  $w_{ij}$  be the connexion forms defined in section 5.2, and let  $\bar{w}_{ij}$  be the global forms defined above. Then  $w_{ij} = \bar{w}_{ij} \circ f_*$  on  $U$ .

Thus the Cartan structural equations 13 and 14 (and the torsion and curvature 2-forms) can be carried up to global equations on  $B$ .

## Problems

Let  $M$  and  $M'$  be  $C^\infty$  manifolds.

42. Let  $x_1 = x$  and  $x_2 = y$  be the usual coordinates on  $\mathbb{R}^2$ . Define a connexion  $D$  on  $\mathbb{R}^2$  by letting  $\Gamma_{jk}^i = 0$  except for  $\Gamma_{12}^1 = \Gamma_{21}^1 = 1$ .

(i) Set up and solve the differential equations for the geodesics thru any point in  $\mathbb{R}^2$ .

(ii) Find the particular geodesic  $g$  with  $g(0) = (2, 1)$  and  $T_g(0) = \frac{\partial}{\partial x} + \frac{\partial}{\partial y}$ .

(iii) Is  $D$  complete?

(iv) Do the geodesics emanating from the origin pass thru all points of the plane?

(v) If  $\sigma$  and  $\gamma$  are geodesics with  $\gamma(0) = \sigma(0)$ , and  $T_\gamma(0) = bt_\sigma(0)$  for  $b \in \mathbb{R}$ , show  $\gamma(t) = \sigma(bt)$  for all possible  $t$

(vi) Investigate the connexion  $D'$  with all  $\Gamma_{jk}^i = 0$  except  $\Gamma_{12}^1 = 1$

43. Let  $D$  be a connexion on  $M$ . Let  $\sigma(t)$  be an integral curve of the  $C^\infty$  field  $X$ , let  $e_1(t), \dots, e_n(t)$  be a parallel base along  $\sigma$  and let  $Y(t) = \sum y_i(t)e_i(t)$  be a  $C^\infty$  field along  $\sigma$ .

(i) Show  $(\mathbf{D}_X Y)(t) = \sum \frac{dy_i}{dt} e_i(t)$  along  $\sigma$ .

(ii) Show

$$(\mathbf{D}_X Y)(0) = \lim_{t \rightarrow 0} \frac{P_{t,0} Y(\sigma(t)) - Y(0)}{t}$$

44. Let  $f$  be a connexion preserving  $C^\infty$  map of  $M$  into  $M'$ . Show

$$f_*(\text{Tor}(X, Y)) = \text{Tor}'(f_*X, f_*Y) \quad \text{and} \quad f_*(R(X, Y)Z) = R'(f_*X, f_*Y)(f_*Z).$$

45. A manifold  $M$  is *parallelizable* if there is a connexion  $\mathbf{D}$  on  $M$  in which parallel translation is independent of curves, and such a  $\mathbf{D}$  is called a *flat connexion*.

(i) Show  $M$  is parallelizable iff there is a global  $C^\infty$  base field on  $M$ .

(ii) If  $\mathbf{D}$  is a flat connexion, show its curvature tensor is zero (see problem 85).

46. Let  $G$  be a Lie group. Define the *left invariant connexion*  $\mathbf{D}$  on  $G$  by asserting all vector fields in the Lie algebra  $\mathfrak{g}$  are parallel fields.

(i) Show  $\mathbf{D}$  is flat,  $G$  is parallelizable, and if  $X$  and  $Y$  are in  $\mathfrak{g}$  then  $\text{Tor}(X, Y) = -[X, Y]$ .

(ii) Show that each geodesic  $g$  on  $G$  is the left translate of a one-parameter sub group  $\sigma$ ; i.e.,  $g(t) = L_{g(0)}(\sigma(t))$  for all  $t$ .

(iii) Show  $\mathbf{D}$  is complete.

47. Let  $\mathbf{D}$  be a connexion on  $M$ . For  $m$  in  $M$ , let  $H_m$  denote the set of linear maps of  $M$  into itself, obtained by parallel translation of  $M_m$  around broken  $C^\infty$  curves starting and ending at  $m$ .

(i) Show  $H_m$  is a group.

(ii) If  $M$  is connected, show  $H_m$  is isomorphic to  $H_{m'}$ , for  $m'$  in  $M$ .

The group  $H_m$  is called the *holonomy group* at  $m$ , and if  $M$  is connected then the *holonomy group of  $M$*  is the group  $H = H_m$  for any  $m$  in  $M$ . Restricting group of  $M$  is the group  $H = H_m$  for any  $m$  in  $M$ . Restricting the closed curves to be null-homotopic, one obtains the *restricted holonomy group*  $H_m^0$ .

(iii) If  $D$  is flat, show  $H_m = 0$ .

(iv) If  $M$  is the unit sphere in  $\mathbb{R}^3$  and  $D$  is the Riemannian connexion, show that  $H = \text{SO}(2, \mathbb{R})$ , where  $\text{SO}(2, \mathbb{R})$  is the *special orthogonal group*, or *rotation group*, consisting of orthogonal maps with determinant one.

48. (Continuing problem 13.) Let  $X = \frac{\partial}{\partial x_1}$  and  $Y = \frac{\partial}{\partial x_2}$  for a coordinate system  $x_1, \dots, x_n$  on  $M$  about  $m$ , and show

$$\lim_{t \rightarrow 0} \frac{[(P_{0,t} - I)(\partial/\partial x_j)]^n}{t} = R(Y, X) \left( \frac{\partial}{\partial x_j} \right) = \sum_k R_{j21}^k \left( \frac{\partial}{\partial x_k} \right)$$

where  $I$  is the identity map and  $P_{0,t}$  is parallel translation along  $\gamma$  from  $\gamma(0)$  to  $\gamma(t)$ . Because of this, one often says  $R(X, Y)$  is “infinitesimal parallel translation around an infinitesimal parallelogram spanned by  $X$  and  $Y$ .”

## 6. Riemannian Manifolds and Submanifolds

The definition of a Riemannian (and a semi-Riemannian) manifold was given in section 2.1. A manifold on which one has singled out a specific symmetric and positive definite (or non-singular) 2-covariant tensor field, called the *metric tensor*, is a Riemannian (or semi-Riemannian) manifold. In this chapter we generalize the theory of Chapters 2 and 3 in a natural way. Much of the theory applies to semi-Riemannian manifolds and submanifolds, but, in general, we phrase things only in Riemannian terms.

### 6.1 Length and Distance

The metric tensor allows us to define lengths, angles, and distances. Let  $M$  be a Riemannian manifold with metric tensor  $\langle -, - \rangle$ . Let  $X, Y \in \mathbf{T}_m M$ . Define the *length* of  $X$  by  $|X| = \sqrt{\langle X, X \rangle}$ . Define the *angle*  $\theta$  between  $X$  and  $Y$  (both non-zero) by letting  $\langle X, Y \rangle = |X||Y| \cos \theta$  where  $0 \leq \theta \leq \pi$ , and notice the *Schwartz inequality*  $|\langle X, Y \rangle| \leq |X||Y|$  makes this possible.

The length of a curve is now defined by integrating the length of its tangent vector field. Let  $\sigma$  be a  $C^\infty$  curve on  $[a, b]$  with tangent field  $T$  (or  $T_\sigma$  if necessary). The *length* of  $\sigma$  from  $a$  to  $b$ , denoted

by  $|\sigma|_a^b$ , is defined by

$$|\sigma|_a^b = \int_a^b \sqrt{\langle T(t), T(t) \rangle} dt \quad (1)$$

The integral exists, since the integrand is continuous. The length of a broken  $C^\infty$  curve is defined as the (finite) sum of the lengths of its  $C^\infty$  pieces. The number  $|\sigma|_a^b$  is independent of the parameterization of its image set in the following sense: let  $g$  be a  $C^1$  map of  $[c, d]$  into  $[a, b]$  with end points mapping to end points (assume  $g(c) = a$  and  $g(d) = b$ ); then

$$\begin{aligned} \int_a^b \sqrt{\langle T_\sigma(t), T_\sigma(t) \rangle} dt &= \int_c^d \sqrt{\langle T_\sigma(g(t)), T_\sigma(g(t)) \rangle} g'(t) dt \\ &= \int_c^d \sqrt{\langle T_{\sigma \circ g}(t), T_{\sigma \circ g}(t) \rangle} dt \end{aligned}$$

since  $T_{\sigma \circ g}(t) = g'(t)T_\sigma(g(t))$  by the chain rule. Thus we can write  $|\sigma|_q^p = |\sigma|_a^b$  where  $q = \sigma(a)$  and  $p = \sigma(b)$ .

Classically, the metric tensor is almost always expressed by the notation

“ $ds^2 = g_{ij} dx_i dx_j$ ”. This means one is giving the inner product on a coordinate domain  $U$  with coordinate functions  $x_1, \dots, x_n$  in terms of the coordinate bases; i.e., if  $X_i = \frac{\partial}{\partial x_i}$ , then  $g_{ij} = \langle X_i, X_j \rangle$  is a  $C^\infty$  function on  $U$ . If

$$Y = \sum_i y_i X_i, \quad Z = \sum_k z_k X_k$$

then

$$\langle Y, Z \rangle = \sum_{i,k=1}^n y_i z_k g_{ik}$$

Thus, giving the matrix of functions  $g_{ij}$  on  $U$  determines the inner product on  $U$ . The “ $ds$ ” only makes sense when one is discussing

a curve  $\sigma$  which maps into  $U$ , for then let  $s(t) = |\sigma|_a^t$  and

$$\left(\frac{ds}{dt}\right)^2 = \langle T, T \rangle = \sum_{ij} g_{ij} \frac{d(x_i \circ \sigma)}{dt} \frac{d(x_j \circ \sigma)}{dt}$$

If  $M$  is connected, a pseudo-metric is defined on  $M$  by

$$d(p, m) = \inf \{|\sigma| : \sigma \text{ a broken } C^\infty \text{ curve from } p \text{ to } m\} \quad (2)$$

Trivially,  $d(p, m) \geq 0$ ,  $d(p, p) = 0$ , and  $d(p, m) = d(m, p)$ . The triangle inequality is left as a problem.

**Theorem 6.1.** The pseudo-metric topology on  $M$  equals the manifold topology.

*Proof.* (After [ST51, p. 44]). Let  $m \in M$ , and let  $x_1, \dots, x_n$  be a coordinate system about  $m$  with domain  $U$ . For  $p \in U$  let  $d_m(p) = d(m, p)$  defined above, and let  $d'(p) = [\sum x_i(p)^2]^{1/2}$  where we assume  $x_i(m) = 0$ . Choose  $a > 0$  so  $A = \{p : d'(p) \leq a\} \subset U$ . On the compact set

$$B = \left\{ (p, X_p) : p \in A, 1 = \sum dx_i(X_p)^2 \right\}$$

the norm function

$$|X_p| = \sqrt{\left[ \sum_{ij} g_{ij}(p) dx_i(X_p) dx_j(X_p) \right]}$$

is a continuous function which takes on a maximum  $R$  and a minimum  $r > 0$ .

Let  $\sigma$  be any broken  $C^\infty$  curve in  $A$  with  $\sigma(0) = m$ ,  $\sigma(b) = p$  and  $(\sigma(t), T_\sigma(t)) \in B$  for all  $t$ . Then

$$|\sigma| = \int_0^b |T_\sigma(t)| dt \geq rb \geq rd'(p)$$

For a broken curve  $\sigma$  from  $m$  to  $p$  that leaves  $A$ , one has  $|\sigma| \geq ra \geq$

$rd'(p)$ . Hence, (1)  $d(p) \geq rd'(p)$ . But if  $\sigma$  is a curve with  $x_i \circ \sigma(t) = \frac{tp_i}{d'(p)}$ , where  $x_i(p) = p_i$ , then

$$|\sigma| = \int_0^{d'(p)} |T_\sigma(t)| dt \leq Rd'(p)$$

Hence, (2)  $d(p) \leq Rd'(p)$ . The inequalities (1) and (2) prove the theorem.  $\square$

**Corollary 6.2.** A connected Riemannian manifold  $M$  is Hausdorff iff the pseudo-metric  $d$  is a metric.

In Chapter 10 we show that geodesics are the curves that locally minimize arc length, i.e., the length of a small piece of a geodesic in  $M$  is precisely the distance between the end points of the piece.

Henceforth we assume all manifolds we mention are Hausdorff. A Riemannian manifold is complete if it is complete as a metric space, i.e., every Cauchy sequences must converge.

## 6.2 Riemannian Connexion and Curvature

A Riemannian connexion  $\mathbf{D}$  on a Riemannian manifold  $M$  is a connexion  $\mathbf{D}$  such that

$$\mathbf{D}_X Y - \mathbf{D}_Y X = [X, Y] \tag{3}$$

and

$$Z \langle X, Y \rangle = \langle \mathbf{D}_Z X, Y \rangle + \langle X, \mathbf{D}_Z Y \rangle \tag{4}$$

for all fields  $X, Y$ , and  $Z$  with a common domain. The fundamental theorem of (semi-) Riemannian manifolds is the following:

**Theorem 6.3.** There exists a unique Riemannian manifold connexion on a (semi-) Riemannian manifold.

*Proof.* We show a Riemannian connexion  $\mathbf{D}$  exists and is unique on every coordinate domain  $U$ . The uniqueness implies  $\mathbf{D}$  must agree on overlapping domains; hence  $\mathbf{D}$  exists and is unique on all of  $M$ .



---

## Section 6.2: Riemannian Connexion and Curvature

Let  $X_1, \dots, X_n$  be the coordinate fields on  $U$ , let  $g_{ij} = \langle X_i, X_j \rangle$  on  $U$ , and let  $(g^{-1})_{ij}$  be the  $ij$ th entry of the inverse matrix of  $g = (g_{ij})$  (which is non-singular). If (3) and (4) hold, then

$$X_i \langle X_r, X_j \rangle + X_j \langle X_r, X_i \rangle - X_r \langle X_i, X_j \rangle = 2 \langle \mathbf{D}_{X_i} X_j, X_r \rangle \quad (5)$$

since  $[X_k, X_s] = 0$  for all  $k, s$ . By section 5.2, giving  $\mathbf{D}$  on  $U$  is equivalent to giving functions  $\Gamma_{jk}^i$  with

$$\mathbf{D}_{X_k}(X_j) = \sum_{i=1}^n \Gamma_{jk}^i X_i$$

and demanding properties (1) through (4) of section 5.1 are valid. Thus (5) implies

$$2 \sum_k \Gamma_{ji}^k g_{kr} = X_i g_{rj} + X_j g_{ri} - X_r g_{ij}$$

hence

$$\Gamma_{ij}^k = \frac{1}{2} \sum_r (g^{-1})_{kr} \left( \frac{\partial g_{rj}}{\partial x_i} + \frac{\partial g_{ri}}{\partial x_j} - \frac{\partial g_{ij}}{\partial x_r} \right) \quad (6)$$

This is the classical expression for the Christoffel function  $\Gamma_{ij}^k$  in terms of the metric tensor. Use (6) to define  $\mathbf{D}$  on  $U$ . A direct check of (3) and (4) shows  $\mathbf{D}$  is Riemannian, and the explicit representation (6) shows  $\mathbf{D}$  is unique.  $\square$

The above theorem is a special case of a more general theorem (problem 70). For the rest of this section, let  $M$  be a (semi-) Riemannian manifold and let  $\mathbf{D}$  be the Riemannian connexion on  $M$ . The Riemann-Christoffel curvature tensor (of type  $\mathbf{0}, 4$ ) is the 4-covariant tensor

$$K(X, Y, Z, W) = \langle X, R(Z, W)Y \rangle$$

for  $X, Y, Z, W \in \mathbf{T}_m M$ .

**Theorem 6.4.** The following relations are true:

(a)  $R(X, Y)Z + R(Z, X)Y + R(Y, Z)X = 0$

(b)  $K(X, Y, Z, W) = -K(Y, X, Z, W)$

(c)  $K(X, Y, Z, W) = -K(X, Y, W, Z)$

(d)  $K(X, Y, Z, W) = K(Z, W, X, Y)$

The relation (a) is called the *first Bianchi identity* and it holds for any symmetric connexion. These relations are equivalent to the “symmetries” of the indices of the classical  $R_{ijkl}$  functions.

*Proof.* For (a), use the Jacobi identity, property (3) above, and compute. For (c), use  $R(Z, W) = -R(W, Z)$ . For (b), use property (4) to shift D from one slot to the other. For (d), notice (a) implies

$$K(X, Y, Z, W) + K(X, W, Y, Z) + K(X, Z, W, Y) = 0 \quad (\text{a}')$$

By writing (a') three more times, cyclically permuting the arguments of the first term one step from one line to the next, adding all four equations, and cancelling via (b) and (c), one obtains (d).  $\square$

For  $X, Y \in \mathbf{T}_m M$ , let

$$A(X, Y) = \langle X, X \rangle \langle Y, Y \rangle - \langle X, Y \rangle^2 \quad (7)$$

If  $A(X, Y) \neq 0$ , let

$$\bar{K}(X, Y) = \frac{K(X, Y, X, Y)}{A(X, Y)} \quad (8)$$

and by direct computations, using the above properties of  $K$ , one can show

$$\bar{K}(X, Y) = \bar{K}(Y, X) = \bar{K}(rX, sY) = \bar{K}(X + tY, Y)$$

for some  $r, s, t \neq 0$ . Thus if  $A(X, Y) \neq 0$  and  $ad - bc \neq 0$ , then

$$\bar{K}(X, Y) = \bar{K}(aX + bY, cX + dY)$$

and we define  $K(P)$ , the Riemannian curvature of the 2-dimensional subspace  $P$  of  $\mathbf{T}_m M$  spanned by  $X$  and  $Y$ , by

$$K(P) = \frac{\langle X, R(X, Y)Y \rangle}{A(X, Y)}$$

In section 2.4, we showed  $K(\mathbf{T}_m M) = K(m)$  is the Gauss curvature of a surface  $M \subset \mathbb{R}^3$ . In the Riemannian case,  $\sqrt{A(X, Y)}$  is the area of the parallelogram spanned by  $X$  and  $Y$ .

Let  $f : M \rightarrow M'$  be a  $C^\infty$  map between Riemannian manifolds. If there is a  $C^\infty$  real valued positive function  $F$  on  $M$  such that, for all  $m \in M$  and all  $X, Y \in \mathbf{T}_m M$ , we have  $\langle f_* X, f_* Y \rangle = F(m) \langle X, Y \rangle$ , then  $f$  is a conformal (or strictly conformal) map and  $F$  is called the scale function. If  $F$  exists but  $F \geq 0$  only, then  $f$  is weakly conformal. If  $F = 1$ , then  $f$  is an isometry. If  $f$  is an isometry and a diffeomorphism, then  $f$  is isometric and  $M$  is isometric to  $M'$ . If  $F$  is constant, then  $f$  is homothetic.

At this point, we explicitly call the reader's attention to problem 52, which is considered an integral part of the theory of Riemannian manifolds.

### 6.3 Curves in Riemannian Manifolds

This section parallels the standard treatment of curves in advanced calculus. Let  $M$  be a Riemannian manifold with Riemannian connexion  $\mathbf{D}$ . Let  $\sigma$  be a  $C^\infty$  curve in  $M$  with tangent field  $V = \sigma_* \left( \frac{\mathbf{d}}{\mathbf{d}t} \right)$ , which can legitimately be called the "velocity vector" of  $\sigma$  since "length" is defined. Assuming  $V$  does not vanish on the domain of  $\sigma$ , define the unit tangent vector and the speed function

$$T(t) = \frac{V(t)}{|V(t)|}, \quad s' = \frac{\mathbf{d}s}{\mathbf{d}t} = |V(t)|$$

respectively, so  $V(t) = s'(t)T(t)$  for  $t$  in the domain of  $\sigma$ . Define the geodesic curvature vector field of  $\sigma$  to be the field  $\mathbf{D}_T T$ , and its

length  $k_1$  is the *geodesic curvature* of  $\sigma$ . Notice that  $\mathbf{D}_T T$  and  $k_1$ , at a particular point on the curve, do not depend on the parameterization of the “point set of the curve” but only on the orientation (choice of “direction”) and the existence of a  $C^\infty$  parameterization with non-vanishing tangent at the point.

The curve  $\sigma$  is a *geodesic* ( $\mathbf{D}_V V = 0$ ) iff  $V$  has constant length and (a)  $\mathbf{D}_T T = 0$  or (b)  $k_1 = 0$ . This follows since

$$\mathbf{D}_V V = s' \mathbf{D}_T(s'T) = s' s'' T + (s')^2 \mathbf{D}_T T$$

and  $s' > 0$  while  $\mathbf{D}_T T$  is orthogonal to  $T$  ( $\langle T, T \rangle = 1$  so  $0 = T \langle T, T \rangle = 2 \langle \mathbf{D}_T T, T \rangle$ ).

When  $k_1(t) > 0$ , define the (first) *normal* to  $\sigma$  at  $\sigma(t)$  to be the unit vector  $N_1(t)$  such that  $\mathbf{D}_T T = k_1 N_1$  at  $t$ . If  $N_1$  is defined on an interval then

$$0 = T \langle N_1, T \rangle = \langle \mathbf{D}_T N_1, T \rangle + \langle N_1, \mathbf{D}_T T \rangle = \langle \mathbf{D}_T N_1, T \rangle + k_1$$

so  $\mathbf{D}_T N_1 \neq 0$  on the interval. The vector  $\mathbf{D}_T N_1 + k_1 T$  is orthogonal to both  $T$  and  $N_1$ ; hence, let its length be  $k_2$ , the *second curvature* or *torsion*. If  $k_2(t) > 0$ , define the *second normal* to  $\sigma$  at  $\sigma(t)$  to be the unit vector  $N_2(t)$  such that  $\mathbf{D}_T N_1 + k_1 T = k_2 N_2$ . If  $k_2 > 0$  on an interval, then the above process can be continued to define  $k_3$ , and where  $k_3 > 0$ , one gets  $N_3$ , etc. The vectors  $T, N_1, N_2, \dots$  are called *Frenet vectors*, and the equations that express the  $\mathbf{D}_T N_i$  in terms of the Frenet vectors are called the *Frenet formulae*.

When  $M$  is a 2-manifold and  $k_1 > 0$ , then the Frenet formulae become

$\mathbf{D}_T T = k_1 N_1$  and  $\mathbf{D}_T N_1 = -k_1 T$ . In this case it is possible to locally choose  $N_1$  along  $\sigma$  independently of  $\mathbf{D}_T T$  (on univalent pieces of  $\sigma$ ), and letting  $\mathbf{D}_T T = k_1 N_1$  would define  $k_1$ , which could take on negative values (see problem 72).

## 6.4 Submanifolds

The theory in sections 2.3 and 2.4 is now generalized. Throughout this section, let the  $k$ -manifold  $M$  be a (non-singular) submanifold of the (semi-)Riemannian manifold  $\bar{M}$ . In the semi-Riemannian case, the submanifold  $M$  is *non-singular* if the metric tensor is non-singular when restricted to  $\mathbf{T}_m M$  for all  $m \in M$  (thus  $M$  is a semi-Riemannian manifold under the induced metric tensor). The induced metric tensor on  $M$  is called the *first fundamental form* on  $M$ . Let  $\bar{D}$  be the Riemannian connexion on  $\bar{M}$ .

**Theorem 6.5.** For  $C^\infty$  fields  $X$  and  $Y$  with domain  $A$  on  $M$  (and tangent to  $M$ ), define  $\mathbf{D}_X Y$  and  $V(X, Y)$  on  $A$  by decomposing  $\bar{D}_X Y$  into its unique tangential and normal components, respectively; thus,

$$\bar{D}_X Y = \mathbf{D}_X Y + V(X, Y) \tag{9}$$

Then  $\mathbf{D}$  is the Riemannian connexion on  $M$  and  $V$  is a symmetric vector-valued 2-covariant  $C^\infty$  tensor called the *second fundamental tensor*. The decomposition equation (9) is called the *Gauss equation*.

*Proof.* We will establish the  $C^\infty$  nature of the decomposition. The rest of the proof will only be outlined, for it is a simple exercise. Use the properties of  $\bar{D}$  (since it is a connexion) to establish the properties of  $\mathbf{D}$  (making it a connexion) and the tensor character of  $V$  (its multilinearity). Zero torsion for  $\bar{D}$  implies zero torsion for  $\mathbf{D}$ , and  $V$  is symmetric (use the proposition in section 2.2, which generalizes trivially). Since  $\bar{D}$  satisfies condition (4) (section 6.2),  $\mathbf{D}$  does too. Hence  $\mathbf{D}$  is Riemannian, and by the uniqueness theorem,  $\mathbf{D}$  is the Riemannian connexion on  $M$ .

To show  $\mathbf{D}$  and  $V$  are  $C^\infty$  on  $A$ , choose  $p \in A$ . Let  $\bar{U}$  and  $U$  be special coordinate domains about  $p$  in  $\bar{M}$  and  $M$ , respectively, with  $U \subset A$ , and let  $\bar{Z}_1, \dots, \bar{Z}_n$  and  $Z_1 = \bar{Z}_1|_U, \dots, Z_k = \bar{Z}_k|_U$  be the coordinate vector fields on  $\bar{U}$  and  $U$ , respectively. Apply the Gram-

Schmidt process to  $\bar{Z}_1, \dots, \bar{Z}_n$  on  $\bar{U}$  to obtain  $C^\infty$  (the Gram-Schmidt process is algebraic) orthonormal fields  $W_1, \dots, W_n$  on  $\bar{U}$  such that  $W_1|_U, \dots, W_k|_U$  give a  $C^\infty$  orthonormal base of  $\mathbf{T}_m M$  for  $m$  in  $U$ , while  $W_{k+1}|_U, \dots, W_n|_U$  give  $\bar{M}$ -vector fields that are  $C^\infty$  on  $U$  and form a base of the orthogonal complement to  $\mathbf{T}_m M$ , for  $m$  in  $U$ . Let

$$X = \sum_{i=1}^k x_i W_i, \quad Y = \sum_{j=1}^k y_j W_j$$

define  $C^\infty$  functions  $x_i$  and  $y_i$  on  $U$  for  $1 \leq i \leq k$ , and let

$$\bar{\mathbf{D}}_{W_i} W_j = \sum_{r=1}^n B_{ji}^r W_r$$

define  $C^\infty$  functions  $B_{ij}^r$  on  $\bar{U}$ . Then

$$\bar{\mathbf{D}}_X Y = \sum (X Y_j) W_j + \sum y_j B_{ji}^r W_r$$

where  $1 \leq i, j \leq k$  and  $1 \leq r \leq n$ ; thus

$$\mathbf{D}_X Y = \sum_{r=1}^k \left[ (X y_r) + \sum_{i,j=1}^k y_j x_i B_{ji}^r \right] W_r$$

and

$$V(X, Y) = \sum_{r=k+1}^n \left[ \sum_{i,j=1}^k y_j x_i B_{ji}^r \right] W_r$$

are  $C^\infty$  on  $U$ . □

By decomposing the curvature  $\bar{R}$  into tangent and normal parts, we obtain the Gauss curvature equation (10), and the Codazzi-Mainardi equation (11), respectively. Let  $X, Y, Z$  be  $C^\infty$  fields tangent to  $M$  with a common domain. Writing the decomposition of a vector  $W$  as  $W = \tan W + \text{nor } W$ ,

$$\tan \bar{R}(X, Y)Z = R(X, Y)Z + \tan \left[ \bar{\mathbf{D}}_X V(Y, Z) - \bar{\mathbf{D}}_Y V(X, Z) \right] \quad (10)$$

and

$$\begin{aligned} \text{nor } \bar{R}(X, Y)Z &= V(X, \mathbf{D}_Y Z) - V(Y, \mathbf{D}_X Z) - V([X, Y], Z) \\ &+ \text{nor } [\bar{\mathbf{D}}_X V(Y, Z) - \bar{\mathbf{D}}_Y V(X, Z)] \end{aligned} \quad (11)$$

Since  $V$  is a tensor, i.e.,  $V(X_m, Y_m)$  is well-defined and independent of the fields  $X$  and  $Y$  used to compute it in the Gauss equation, we define  $X$  and  $Y$  to be conjugate vectors at  $m$  if  $V(X, Y) = 0$ . A vector  $X \in \mathbf{T}_m M$  is an asymptotic vector if  $V(X, X) = 0$ , and in any case define the asymptotic (or normal) curvature of  $X$ ,  $k_X$ , by  $k_X = |V(X, X)|$ . If  $V_m = 0$  then  $m$  is a flat point on  $M$ .

If  $\sigma$  is a curve in  $M$  with  $C^\infty$  unit tangent  $T$ , then  $V(T, T)$  is the normal curvature vector field of  $\sigma$  and  $k_T = |V(T, T)|$  is the normal curvature of  $\sigma$ .

**Theorem 6.6. (Meusnier).** All curves on  $M$  with the same unit tangent  $T$  at a point have the same normal curvature at that point. If  $\sigma$  is a curve on  $M$  with  $C^\infty$  unit tangent  $T$ , then  $(\bar{k}_1)^2 = (k_1)^2 + (k_T)^2$  relates the geodesic curvatures  $\bar{k}_1$  and  $k_1$  of  $\sigma$  in  $\bar{M}$  and  $M$  with its normal curvature  $k_T$ . Moreover,

$k_T = \bar{k}_1 \cos \phi$  determines the angle  $\phi$  between the normal  $\bar{N}_1$  of  $\sigma$  in  $\bar{M}$  and the normal curvature vector  $V(T, T)$  if  $\phi$  is defined.

*Proof.* The first sentence follows since  $V$  is a tensor. The second sentence follows from the Gauss equation  $\bar{\mathbf{D}}_T T = \mathbf{D}_T T + V(T, T)$  since the vectors on the right are orthogonal. For the third sentence, if  $\bar{k}_1 = 0$  then  $k_1 = k_T = 0$  and  $\phi$  is undefined. If  $\bar{k}_1 > 0$  and  $k_T = 0$  then  $V(T, T) = 0$ ,  $\bar{N}_1$  is tangent to  $M$ , and  $\phi = \frac{\pi}{2}$  (if anything). If  $k_T \neq 0$ , let  $N$  be the unit normal in the direction of  $V(T, T)$  and

$$k_T = \langle V(T, T), N \rangle = \langle \bar{\mathbf{D}}_T T, N \rangle = \bar{k}_1 \cos \phi$$

□

The theorem and corollary at the end of section 2.3 can now be generalized by replacing  $\mathbb{R}^n$  by  $\bar{M}$ .

## 6.5 Hypersurfaces

In this section, let  $M$  be a hypersurface in the Riemannian manifold  $\bar{M}$  and let  $N$  be a  $C^\infty$  unit normal on  $M$ . Define the Weingarten map  $L(X) = \bar{\mathbf{D}}_X N$  for  $X \in \mathbf{T}_m M$  (as in section 2.2). The Gauss equation for  $M$  now becomes

$$\bar{\mathbf{D}}_X Y = \mathbf{D}_X Y - \langle LX, Y \rangle N \quad (12)$$

since

$$\langle V(X, Y), N \rangle = \langle \bar{\mathbf{D}}_X Y, N \rangle = X \langle N, Y \rangle - \langle Y, LX \rangle$$

and  $\langle N, Y \rangle \equiv 0$ . Thus  $V(X, Y) = -\langle LX, Y \rangle N$ .

The fundamental forms and the imbedded geometric variants of  $M$  in  $\bar{M}$  are defined in terms of  $L$  exactly as in section 2.2. Notice in this case  $V$  being symmetric is equivalent to  $L$  being self-adjoint.

The Gauss curvature equation (10) and Codazzi-Mainardi equation (11) now become

$$\tan \bar{R}(X, Y)Z = R(X, Y)Z - [\langle LY, Z \rangle LX - \langle LX, Z \rangle LY] \quad (13)$$

and

$$\text{nor } \bar{R}(X, Y)Z = -\langle \mathbf{D}_X LY - \mathbf{D}_Y LX - L[X, Y], Z \rangle N \quad (14)$$

respectively.

The torsion tensor is generalized by defining for any  $C^\infty$  linear transformation valued tensor  $W_p : \mathbf{T}_p M \rightarrow \mathbf{T}_p M$ , on a  $C^\infty$  manifold  $M$ , the torsion of  $W$ ,  $\text{Tor}_W$ , by

$$\text{Tor}_W(X, Y) = \mathbf{D}_X W(Y) - \mathbf{D}_Y W(X) - W[X, Y] \quad (15)$$

The Codazzi-Mainardi equation (14) on a hypersurface becomes

$$\text{nor } \bar{R}(X, Y)Z = -\langle \text{Tor}_L(X, Y), Z \rangle N$$



Thus  $\text{Tor}_L = 0$  on  $M$  iff

$$\bar{R}(X, Y)Z = R(X, Y)Z - [\langle LY, Z \rangle LX - \langle LX, Z \rangle LY] \quad (16)$$

The following theorem generalizes the “theorema egregium” of Gauss, and actually, it may be generalized to the case where  $M$  is a  $k$ -submanifold of  $\bar{M}$  (see [Hic63b]).

**Theorem 6.7.** Let  $M$  be a hypersurface in the Riemannian manifold  $\bar{M}$ , let  $P$  be a 2-dimensional subspace of  $\mathbf{T}_m M$ , and let  $K(P)$  and  $\bar{K}(P)$  be the Riemannian curvature of  $P$  in  $M$  and  $\bar{M}$  respectively. Let  $N$  be a unit  $C^\infty$  normal on a neighborhood of  $m$ , and let  $LX = \bar{D}_X N$  for  $X \in \mathbf{T}_m M$ . If  $X$  and  $Y$  form an orthonormal base of  $P$ , then

$$\bar{K}(P) = K(P) - (\langle LY, Y \rangle \langle LX, X \rangle - \langle LX, Y \rangle^2) \quad (17)$$

*Proof.* Combine the definition of Riemannian curvature with the Gauss curvature equation (13). □

When  $\bar{M}$  is a 3-manifold, the above theorem shows the determinant of  $L$  is independent of the imbedding (i.e., independent of  $L$ ) but depends only on the Riemannian structure of  $\bar{M}$  and  $M$ .

A related result is a form of the Lemma of Synge.

**Theorem 6.8.** Let  $k > 1$ , and let  $M$  be a  $k$ -submanifold of the Riemannian  $n$ -manifold  $\bar{M}$ . Let  $g$  be a geodesic of  $\bar{M}$  that lies in  $M$ , let  $T$  be the unit tangent to  $g$ , let  $X$  be a unit field tangent to  $M$  which is parallel in  $M$  along  $g$  and orthogonal to  $T$ , and let  $P$  be the subspace spanned by  $X$  and  $T$ . Then  $\bar{K}(P) \geq K(P)$  along  $g$ , and  $\bar{K}(P) = K(P)$  iff  $X$  is parallel along  $g$  in  $\bar{M}$ .

*Proof.* We prove the theorem for  $k = n - 1$ , leaving the other cases to problem 55. Let  $N$  be a  $C^\infty$  unit normal on a neighborhood of a point on  $g$  and let  $L(Z) = \bar{D}_Z N$ . Here  $\bar{D}_T T = 0$ , so  $\mathbf{D}_T T = 0$  and  $\langle LT, T \rangle = 0$ . By the previous theorem,

$$\bar{K}(P) = K(P) + \langle LX, T \rangle^2 \geq K(P)$$

If equality holds then  $\langle LX, T \rangle = 0$ , so  $\bar{\mathbf{D}}_T X = \mathbf{D}_T X = 0$ , and conversely.  $\square$

There is a basic “rigidity” theorem for hypersurfaces of  $\mathbb{R}^n$  which is our next goal. This theorem is a uniqueness theorem, and there is a corresponding existence theorem that is proved in Chapter 9. When  $n = 3$ , the theorem was first proved by O. Bonnet (1867).

Intuitively, this theorem states that if two hypersurfaces of  $\mathbb{R}^n$  are isometric and their normals are “bending the same”, then by a “rigid motion” one can superimpose the two manifolds.

**Theorem 6.9.** Let  $M$  and  $M'$  be connected hypersurfaces in  $\mathbb{R}^n$  for  $n \geq 3$ . Let  $N$  and  $N'$  be  $C^\infty$  unit normal fields on  $M$  and  $M'$ , respectively. Let  $F$  be a diffeomorphism on  $M$  onto  $M'$  that preserves the first and second fundamental forms. Then there is an isometry  $G$  of  $\mathbb{R}^n$  with  $F = G|_M$ .

*Proof.* During this proof let us use “primes” to denote concepts belonging to  $M'$  which correspond to familiar concepts for  $M$ ; i.e., let  $L(X) = \bar{\mathbf{D}}_X N$  for  $X \in \mathbf{T}_p M$  and  $L'(Y) = \bar{\mathbf{D}}_Y N'$  for  $Y \in \mathbf{T}_{p'} M'$ . The hypothesis states if  $X, Z \in \mathbf{T}_p M$  then

$$\langle F_* X, F_* Z \rangle = \langle X, Z \rangle, \quad \langle L'(F_* X), F_* Z \rangle = \langle LX, Z \rangle$$

Combining these statements,

$$\langle L'(F_* X), F_* Z \rangle = \langle LX, Z \rangle = \langle F_* LX, F_* Z \rangle$$

for all  $Z$ , which implies  $L' \circ F_* = F_* \circ L$ . Thus the hypothesis could be rephrased as a demand that  $F$  be an isometry of  $M$  onto  $M'$  whose Jacobian commutes with the Weingarten maps. Since an isometry is connexion-preserving,

$$F_*(\mathbf{D}_X Z) = \mathbf{D}'_{F_* X} F_* Z$$

for vectors  $X$  and fields  $Z$  tangent to  $M$ .

If  $p \in M$ , we extend the Jacobian of  $F$  to be a linear map of  $\mathbf{T}_p\mathbb{R}^n$  onto  $\mathbf{T}_{p'}\mathbb{R}^n$  where  $p' = F(p)$ . Let  $W \in \mathbf{T}_p\mathbb{R}^n$ , then  $W = W_t + aN_p$  where  $W_t$  is tangent to  $M$ , so define

$$F_*(W) = F_*(W_t) + aN'_{p'}$$

If  $X \in \mathbf{T}_pM$  and  $W$  is a  $C^\infty$  field of  $\mathbb{R}^n$ -vectors on  $M$ , then

$$F_*(\bar{D}_X W) = \bar{D}_{F_*X}(F_*W)$$

where  $\bar{D}$  is a natural covariant differentiation on  $\mathbb{R}^n$ . This follows since

$$\bar{D}_X W = \bar{D}_X W_t + \bar{D}_X(aN) = D_X W_t - \langle LX, W_t \rangle N + (Xa)N + aLX$$

and

$$\begin{aligned} F_*(\bar{D}_X W) &= D'_{F_*X} F_*W_t - \langle F_*LX, F_*W_t \rangle N' + F_*X(a \circ F^{-1})N' + (a \circ F^{-1})L'F_*X \\ &= \bar{D}_{F_*X} F_*W_t + \bar{D}_{F_*X}((a \circ F^{-1})N') \\ &= \bar{D}_{F_*X} F_*W \end{aligned}$$

Now let  $e_1, \dots, e_n$  be the usual orthonormal fields on  $\mathbb{R}^n$  and define  $C^\infty$  functions  $b_{rs}$  on  $M$  by

$$F_*(e_s)_p = \sum_{r=1}^n b_{rs}(p)(e_r)_{p'}$$

The functions  $b_{rs}$  are  $C^\infty$  since  $F, M, M'$  are  $C^\infty$ , and the  $n$  by  $n$  matrix  $b_{rs}(p)$  is orthogonal since  $F$  is an isometry. If  $X \in \mathbf{T}_pM$  then  $\bar{D}_X e_s = 0$  since  $e_s$  are parallel fields. Thus

$$0 = F_*(\bar{D}_X e_s) = \bar{D}_{F_*X}(F_*e_s) = \sum_{r=1}^n \left[ (Xb_{rs})e_r + b_{rs}\bar{D}_{F_*X}e_r \right] = \sum_r (Xb_{rs})e_r$$

so  $Xb_{rs} = 0$  for all  $r, s$ . Since  $X$  and  $p$  are arbitrary and  $M$  is con-

nected, the functions  $b_{rs}$  are constant on  $M$  and thus the Jacobian of  $F$  is a constant orthogonal transformation relative to the natural base  $e_1, \dots, e_n$  of  $\mathbb{R}^n$ .

Next define a map  $G : \mathbb{R}^n \rightarrow \mathbb{R}^n$  which is a translation followed by an orthogonal map by letting for one  $p \in M$  and requiring  $(G_*)_p = (F_*)_p$ . This completely determines  $G$  and the Jacobian of  $G$  is constant and hence equal to the Jacobian of  $F$  at all points. Since  $M$  is connected,  $F = G|_M$ .  $\square$

## 6.6 Cartan Viewpoint and Coordinate Viewpoint

In this section, let  $M$  be a hypersurface of a Riemannian  $n$ -manifold  $\bar{M}$ . Let  $p \in M$ , let  $\bar{U}$  be a special coordinate neighborhood of  $p$  in  $\bar{M}$  and  $U$  the corresponding neighborhood of  $p$  in  $M$  with  $U \subset \bar{U}$ . Apply the Gram-Schmidt process to the coordinate vector fields on  $\bar{U}$  to obtain an orthonormal base field  $e_1, \dots, e_n$  on  $\bar{U}$  with  $e_1(m), \dots, e_{n-1}(m)$  a base of  $\mathbf{T}_m M$  for  $m \in U$  and  $e_n(m)$  normal to  $\mathbf{T}_m M$  (thus  $e_n$  provides a local normal for the neighborhood  $U$ ). Let  $f : U \rightarrow \bar{U}$  be the inclusion map.

Applying the results of section 5.2, let  $\bar{w}_1, \dots, \bar{w}_n$  be the dual 1-forms associated with  $e_1, \dots, e_n$  and let  $\bar{w}_{ij}$  for  $1 \leq i, j \leq n$  be the connexion 1-forms associated with the Riemannian connexion  $\bar{\mathbf{D}}$  on  $\bar{U}$ , so

$$\bar{\mathbf{D}}_X e_j = \sum_{i=1}^n \bar{w}_{ij}(X) e_i \quad (18)$$

for  $1 \leq j \leq n$ .

Let  $w_{ij} = \bar{w}_{ij}|_U$  and  $w_i = \bar{w}_i|_U$  for  $1 \leq i, j \leq n$ , i.e.,  $w_{ij} = f^* \bar{w}_{ij}$  and  $w_i = f^* \bar{w}_i$ . If  $X$  is tangent to  $M$  at  $m \in U$ , by the Gauss equation,

$$\mathbf{D}_X e_j = \sum_{i=1}^{n-1} w_{ij}(X) e_i \quad (19)$$

$$V(X, e_j) = w_{nj}(X) e_n \quad (20)$$

---

## Section 6.6: Cartan Viewpoint and Coordinate Viewpoint

---

for  $1 \leq j \leq n-1$ . Thus  $w_{ij}$  for  $i, j \leq n$  are the connexion forms for the induced Riemannian connexion  $\mathbf{D}$  on  $M$ . Moreover,

$$L(X) = \bar{\mathbf{D}}_X e_n = \sum_{i=1}^{n-1} w_{in}(X) e_i \quad (21)$$

since  $L(X) \in \mathbf{T}_m M$ , so  $w_{nn} = 0$  on  $U$ . Also  $w_n = 0$  on  $U$  since  $e_n$  is normal to  $M$ . Equation (18) is the Gauss equation and equation (21) is the Weingarten equation.

Let I, II, III be the first, second, and third fundamental forms, respectively. Then for  $X, Y \in \mathbf{T}_m M$ ,  $m \in U$ ,

$$\begin{aligned} \text{I}(X, Y) &= \sum_{i=1}^{n-1} w_i(X) w_i(Y) \\ \text{II}(X, Y) &= \langle LX, Y \rangle = \sum_{i=1}^{n-1} w_{in}(X) w_i(Y) \\ \text{III}(X, Y) &= \langle LX, LY \rangle = \sum_{i=1}^{n-1} w_{in}(X) w_{in}(Y) \end{aligned}$$

Notice

$$0 = X \langle e_i, e_j \rangle = \langle \mathbf{D}_X e_i, e_j \rangle + \langle e_i, \mathbf{D}_X e_j \rangle = w_{ji}(X) + w_{ij}(X)$$

for all  $X$  tangent to  $M$ , i.e.  $w_{ji} = -w_{ij}$  for connexion forms belonging to an orthonormal base (and this again shows  $w_{nn} = 0$ ). Thus we can write II and III in terms of  $w_{ni}$  if we wish.

Certain relations are implied by the Cartan structural equations. The equation

$$d\bar{w}_n = - \sum_{j=1}^n \bar{w}_{nj} \wedge \bar{w}_j = 0$$

(on  $\mathbf{T}_m M$ ) implies II is symmetric. The equation

$$d\bar{w}_{nn} = - \sum_{j=1}^n \bar{w}_{nj} \wedge \bar{w}_{jn} = 0$$

(on  $T_m M$ ) implies III is symmetric. For  $i, j \leq n$ ,

$$d\bar{w}_{ij} = - \sum_{s=1}^n \bar{w}_{is} \wedge \bar{w}_{sj} + \bar{R}_{ij}$$

when restricted to vectors on  $T_m M$ , gives

$$f^* d\bar{w}_{ij} = dw_{ij} = - \sum_{s=1}^{n-1} w_{is} \wedge w_{sj} + R_{ij} = - \sum_{s=1}^n w_{is} \wedge w_{sj} + \bar{R}_{ij}$$

Thus

$$R_{ij} = -w_{in} \wedge w_{nj} + \bar{R}_{ij} \quad (22)$$

which is the Gauss curvature equation from this point of view. For  $i \leq n$ ,

$$f^* d\bar{w}_{in} = dw_{in} = - \sum_{s=1}^{n-1} w_{is} \wedge w_{sn} + \bar{R}_{in} \quad (23)$$

is the Codazzi-Mainardi equation.

For the coordinate viewpoint, let  $x_1, \dots, x_n$  be the special coordinate system on  $\bar{U}$  such that  $x_1, \dots, x_{n-1}$  give coordinates on  $U$ . Let  $X_i = \frac{\partial}{\partial x_i}$  for  $1 \leq i \leq n-1$  and let  $X_n = e_n$  be the unit normal (on  $U$ ). Now apply the above analysis to the base field  $X_1, \dots, X_n$  (and this time  $w_{ij} \neq -w_{ji}$  necessarily since the base  $X_1, \dots, X_n$  is not necessarily orthonormal).

## 6.7 Canonical Spaces of Constant Curvature

We exhibit the three classical examples of  $n$ -dimensional ( $n \geq 2$ ) simply connected complete spaces with constant Riemannian curvature  $K = 0$ ,  $K > 0$ , and  $K < 0$ ; i.e., the Riemannian curvature  $K(P)$  of all plane sections is a constant.

For  $K = 0$ , let  $M = \mathbb{R}^n$  with the usual Riemannian metric. This is usually called *Euclidean space* or *flat space*.

---

## Section 6.7: Canonical Spaces of Constant Curvature

---

For  $K > 0$ , let

$$M = \left\{ a \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} a_i^2 = \frac{1}{K} \right\}$$

i.e.  $M$  is the  $n$ -dim sphere of radius  $\frac{1}{\sqrt{K}}$  about the origin in  $\mathbb{R}^{n+1}$ . It is a Riemannian manifold via the induced metric from  $\mathbb{R}^{n+1}$ . This is called *spherical space* or *Riemann space*. Letting  $N$  be the unit outer normal on  $M$ , then  $L(X) = \sqrt{K}X$  for all vectors tangent to  $M$ , and all points are umbilic. By equation (17) above,

$$K(P) = \langle LX, X \rangle \langle LY, Y \rangle = K$$

where  $X$  and  $Y$  are unit orthogonal vectors spanning  $P$ . Since  $M$  is compact, it is complete. An alternate proof that  $M$  has constant curvature is provided by the group of orthogonal transformations on  $\mathbb{R}^{n+1}$ , which provides isometries that will map any point  $m$ , and plane section  $P$  at  $m$ , into any other point  $m'$  and plane section  $P'$ . Since an isometry preserves the curvature, this would show  $M$  has constant Riemannian curvature but would not evaluate this constant.

For  $K < 0$ , let

$$M = \left\{ a \in \mathbb{R}^n : \sum_{i=1}^n a_i^2 < -\frac{4}{K} \right\}$$

Let  $x_1, \dots, x_n$  be the usual coordinate functions on  $\mathbb{R}^n$ , i.e.,  $x_i(a) = a_i$ , let  $X_i = \frac{\partial}{\partial x_i}$  for  $1 \leq i \leq n$ , and define a metric on  $M$  by the functions

$$g_{ij} = \langle X_i, X_j \rangle = \frac{\delta_{ij}}{A^2}, \quad \text{where} \quad A = 1 + \frac{K}{4} \sum_{r=1}^n x_r^2$$

Then  $M$  with this metric is called *hyperbolic space* or *Poincaré space*. Thus  $M$  is obtained by a conformal change of the usual metric tensor on an open ball in  $\mathbb{R}^n$ , and  $M$  is simply connected since it is contractible.

One proves  $M$  has constant negative Riemannian curvature  $K$

by a direct computation which we outline. Let  $K_{ij}$  be a Riemannian curvature of the plane section spanned by  $X_i, X_j$  at any point in  $M$ . Let

$$R(X_i, X_j)X_r = \sum_k R_r^k(X_i, X_j)X_k = \sum_k R_{rij}^k X_k$$

define functions  $R_{rij}^k$ . Then  $K_{ij} = A^2 R_{jij}^i$ , and compute via the classical formulae for  $R_{rij}^k$  in terms of  $\Gamma_{jk}^i$ , and  $\Gamma_{jk}^i$  in terms of  $g_{ij}$ . These formulae show

$$\Gamma_{jk}^i = 0, \quad \text{unless two indices are equal}$$

and

$$\Gamma_{ij}^i = \Gamma_{ji}^j = \Gamma_{ii}^i = -\frac{Kx_i}{2A}, \quad \Gamma_{ii}^j = \frac{Kx_j}{2A}.$$

Then

$$R_{jij}^i = \frac{K}{A} - \frac{K^2}{4A^2} \sum_r x_r^2, \quad K_{ij} = K.$$

Also by direct computation one shows

$$R_{jkr}^i = 0, \quad \text{unless } k = i, r = j \text{ or } k = j, r = i$$

Then letting  $e_i = AX_i$  gives an orthonormal base  $e_1, \dots, e_n$  at each point of  $M$ . Let  $P$  be any plane section at  $m \in M$  and let  $f_1, f_2$  be an orthonormal base of  $P$  which we extend to an orthonormal base of  $T_m M$ . Then the base  $e_i$  is related to the base  $f_j$  via an orthogonal matrix, and one uses this fact to show  $K(f_1, f_2) = K$ . Thus  $M$  has constant negative curvature. To show  $M$  is complete, let  $K = -B^2$ , and one shows the curve

$$g(t) = \left( 2 \frac{\sinh \frac{B}{2} t}{B \cosh \frac{B}{2} t}, 0, \dots, 0 \right)$$

is a geodesic defined for all  $t$  and parameterized by arc length. Such a geodesic is obtained on every ray emanating from the origin  $\mathcal{O}$



by symmetry. Thus

$$\bar{B}_M(0, t) = \bar{B}_{\mathbb{R}^n} \left( 0, 2 \frac{\sinh \frac{B}{2}t}{B \cosh \frac{B}{2}t} \right)$$

which is a compact set, so  $M$  is complete. (Here,

$$\bar{B}_m(p, r) = \{m \in M : d_M(m, p) \leq r\}$$

where  $d_M$  is the distance function in  $M$ .) Note that the mapping  $g$ , when generalized to all rays in  $\mathbb{R}^n$ , exhibits explicitly the exponential map of  $T_0M$  onto  $M$  (see section 9.3).

For  $K > 0$ , let  $M = \mathbb{R}^n$ , and let  $g_{ij} = \delta_{ij}/A^2$  define a Riemannian metric on  $M$  as above. The above computations show  $M$  has constant Riemannian curvature  $K$  and  $M$  is trivially simply connected. But  $M$  is not complete since  $\bar{B}_M \left( 0, \frac{2\pi}{\sqrt{K}} \right) = \mathbb{R}^n$  is not compact. Thus we have an example of a conformal change of metric which changes a complete Riemannian manifold into a non-complete Riemannian manifold.

## 6.8 Existence

The objective of this section is to show a paracompact connected Hausdorff  $C^\infty$  manifold admits a Riemannian metric. This is accomplished by constructing a “partition of the unit function”. The function  $e^{-1/x^2}$  is the principal tool which is used to show there are “many”  $C^\infty$  functions on a  $C^\infty$  manifold.

**Lemma 6.10.** Given real numbers  $0 < b < c$ , there exists a  $C^\infty$  function  $f : \mathbb{R} \rightarrow \mathbb{R}$  with  $f(t) = 0$  for  $t \leq b$ ,  $0 \leq f(t) \leq 1$  for all  $t$ , and  $f(t) = 1$  for  $t \geq c$ .

*Proof.* Consider the  $C^\infty$  function

$$g(x) = \begin{cases} 0 & x \leq 0 \\ e^{-1/x^2} & x > 0 \end{cases}$$

We outline a sequence of operations which leads to the desired functions, and we illustrate (and number) the graphs of these intermediate functions in Fig. 6.1. Translate  $g$  so its graph moves  $\frac{1}{2}(c - b)$  units to the left (this is no. (1)). Reflect the graph of (1) about the  $y$ -axis to obtain (2). Multiply (1) and (2) to obtain (3). Integrate (3) to obtain (4). Multiply (4) by a scale factor to obtain (5). Translate the graph of (5) to the right to obtain the desired function  $f$ .  $\square$

**Lemma 6.11.** Given real numbers  $0 < b < c$ , there exists a  $C^\infty$  function

$F : \mathbb{R}^n \rightarrow \mathbb{R}$  with  $F(p) = 0$  for  $|p| \leq b$ ,  $0 \leq F(p) \leq 1$  for all  $p$ , and  $F(p) = 1$  for  $|p| \geq c$ .

*Proof.* Let  $F(p) = f(|p|)$  where  $f$  is obtained from lemma 6.10.  $\square$

**Lemma 6.12.** If  $M$  is a Hausdorff  $C^\infty$  manifold and  $m \in M$ , then there is a coordinate neighborhood  $U$  of  $m$  and a  $C^\infty$  function  $f : M \rightarrow \mathbb{R}$  such that  $f(p) > 0$  for  $p \in U$  and  $f(p) = 0$  for  $p \notin U$ .

*Proof.* Let  $V$  be any coordinate neighborhood of  $m$  with coordinate map

$\phi : V \rightarrow \mathbb{R}^n$  such that  $\phi(m) = 0$ . Choose real numbers  $0 < b < c$  such that  $B(0, c) \subset \phi(V)$ . Apply lemma 6.11 to obtain  $F$  and let  $G = 1 - F$ . Then let  $U = \phi^{-1}(B(0, c))$  and let  $f = G \circ \phi$  on  $U$  while  $f(p) = 0$  for  $p \notin U$ .  $\square$

**Lemma 6.13.** If  $M$  is a paracompact Hausdorff  $C^\infty$  manifold then there exists a locally finite covering  $\{U_\alpha\}$ , where  $U_\alpha$  are open coordinate neighborhoods, and a collection of non-negative real valued  $C^\infty$  functions  $\{g_\alpha\}$  such that  $g_\alpha(p) = 0$  for  $p \notin U_\alpha$  and  $\sum_\alpha g_\alpha = 1$ . The collection  $\{g_\alpha\}$  is called a *partition of unity* for the covering  $\{U_\alpha\}$ .

*Proof.* Combining lemma 6.12 and the definition of paracompactness, one obtains the desired covering  $\{U_\alpha\}$  with  $C^\infty$  functions




Figure 6.1: Constructing a  $C^\infty$  Step Function

$f_\alpha : M \rightarrow \mathbb{R}$  such that  $f_\alpha > 0$  on  $U_\alpha$  and  $f_\alpha = 0$  on  $M - U_\alpha$ . The function  $F = \sum_\alpha f_\alpha$  is a well-defined non-vanishing  $C^\infty$  function on  $M$  since the covering  $\{U_\alpha\}$  is locally finite. Finally let  $g_\alpha = f_\alpha/F$ .  $\square$

**Theorem 6.14.** If  $M$  is a connected Hausdorff  $C^\infty$  manifold then the following are equivalent:

- (a)  $M$  is paracompact.
- (b)  $M$  admits a Riemannian metric.
- (c)  $M$  is second-countable (completely separable).

*Proof.* We show (a) implies (b) and give references for the other implications whose proofs are purely topological.

Assuming (a), apply lemma 6.13 to obtain a locally finite cover  $\{U_\alpha\}$  with the partition of unity  $\{g_\alpha\}$ . On each coordinate neighborhood  $U_\alpha$ , define a local Riemannian metric tensor  $\langle -, - \rangle_\alpha$  by demanding the coordinate map be an isometry. Then the tensor  $g_\alpha \langle -, - \rangle_\alpha$  is a global  $C^\infty$  tensor on  $M$  that vanishes outside  $U_\alpha$ . At any point  $m \in M$ , for  $X, Y \in \mathbf{T}_m M$ , let

$$\langle X, Y \rangle = \sum_\alpha g_\alpha(m) \langle X, Y \rangle_\alpha$$

This defines a  $C^\infty$  Riemannian metric tensor on  $M$  which shows (a) implies (b).

Assuming (b), then from section 2.6 we know  $M$  is a metric space and hence must be paracompact (see [Kel17, p. 160]). Thus (b) implies (a). That (c) implies (a) follows from [HY12, p. 79]. To show (b) implies (c), we refer the reader to Chapter 6 in [Kel17]. The metric can be used to define a uniform structure on  $M$  which must admit a countable base (see [Kel17, p. 186]).  $\square$

For theorems concerning the imbedding of manifolds in other manifolds, see [Ste64], [AM12] or [Sma61].

## Problems

All manifolds will be Riemannian unless otherwise stated.

49. If  $M$  is semi-Riemannian and  $D$  satisfies (4), then  $D$  is metric preserving. Show that  $D$  is metric preserving iff for parallel fields  $Y$  and  $Z$  along a curve  $\sigma$ , the function  $\langle Y, Z \rangle$  is constant on  $\sigma$ .
50. Let  $R$  and  $R'$  be two linear map valued skew-symmetric 2-covariant tensors whose corresponding  $K$  and  $K'$  satisfy properties (a) through (d) on p. 103. Show  $K = K'$  iff  $R = R'$ .
51. (i) If  $f$  is a  $C^\infty$  strictly conformal map, show  $f_*$  has no kernel and preserves angles.  
 (ii) If  $f$  is a complex analytic map, show

$$\langle f_*X_p, f_*Y_p \rangle = |f'(p)|^2 \langle X_p, Y_p \rangle$$

where  $f : \mathbb{C} \rightarrow \mathbb{C}$ .

52. Let  $f : M \rightarrow M'$  be a strictly conformal map with scale function  $F$ .
- (i) Show  $f$  is (Riemannian) connexion preserving iff  $F$  is constant and  $f(M)$  is flat.
- (ii) If  $f$  is an isometry, show  $f$  preserves the curvature tensor and the Riemannian curvature.
53. With the standard hypothesis of section 3.3, show if  $f$  is connexion preserving, then  $M$  is a sphere, a plane, or a right circular cylinder (see [Hic63b]).
54. Let  $M \subset \mathbb{R}^n$  be a hypersurface, let  $N$  be a  $C^\infty$  unit normal, let  $g \in C^\infty(M, \mathbb{R})$ , and define  $f_t : M \rightarrow \mathbb{R}^n$  by

$$f_t(p) = p + tg(p)N_p$$

(a) Show that

$$(f_t)_*X = X + t(Xg)N + tgLX$$

for  $X$  tangent to  $M$ .

(b) If  $f_t$  is an isometry for  $t > 0$ , show that  $M$  is flat.

55. Generalize the first two theorems in section 6.5 to the case of a  $k$ -submanifold that is framed in an  $n$ -manifold for  $1 < k < n$  (see [Hic63a]). In the second theorem, if  $k = 2$  and  $n = 3$ , show  $\bar{K}(P) = K(P)$  iff  $g$  is a line of curvature on  $M$ .

56. If  $u$  and  $v$  are orthonormal coordinates with domain  $A$  on a 2-manifold (thus  $\frac{\partial}{\partial u}$  and  $\frac{\partial}{\partial v}$  are orthonormal), show the coordinate curves are geodesics and  $K \equiv 0$  on  $A$ .

57. (K. Leisenring)

(i) Show that

$$f(u, v) = (\cos u \cos v, \cos u \sin v, \sin u \cos v, \sin u \sin v)$$

is an isometric imbedding of the flat torus  $T$  into the unit sphere  $S^3$  in  $\mathbb{R}^4$ .

(ii) Show the total imbedded curvature of  $f(T)$  in  $S^3$  is a constant negative one.

58. Let  $M$  be connected with symmetric connexion  $D$  and let  $L_p : \mathbf{T}_m M \rightarrow \mathbf{T}_m M$  be a  $C^\infty$  linear map valued function on  $M$ . If  $\text{Tor}_L \equiv 0$  and all points are  $L$ -umbilic, show  $L$  is a constant multiple of the identity map.

59. (i) Show that every isometry of  $\mathbb{R}^n$  can be factored uniquely into an orthogonal map followed by a translation.

(ii) If  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is orthogonal, show  $f_* = f$  in a natural way.

60. If  $e_1, \dots, e_n$  is an orthonormal base field with dual base  $w_1, \dots, w_n$  and  $M$  has constant Riemannian curvature  $K$ , show the associated curvature forms  $R_{ij} = Kw_i \wedge w_j$ .

61. If  $M$  has constant Riemannian curvature  $K$  and one defines a metric on  $M \times M$  by

$$\langle (X_1, Y_1), (X_2, Y_2) \rangle = \langle X_1, X_2 \rangle + \langle Y_1, Y_2 \rangle$$

does  $M \times M$  have constant curvature?

62. If  $x_1, \dots, x_n$  are coordinates on a hypersurface  $U \subset \mathbb{R}^{n+1}$ , let

$$x_i = \frac{\partial}{\partial x_i}, \quad g_{ij} = \langle X_i, X_j \rangle, \quad b_{ij} = \langle LX_i, X_j \rangle, \quad LX_j = \sum_i a_{ij} X_i$$

Show that

$$a_{ij} = \sum_r (g^{-1})_{ir} b_{rj} \quad \text{(Weingarten equation)}$$

$$R_{jkh}^i = \sum_r (g^{-1})_{ir} (b_{hj} b_{rk} - b_{kj} b_{rh}) \quad \text{(Gauss curvature equation)}$$

$$\frac{\partial b_{ir}}{\partial x_s} - \frac{\partial b_{is}}{\partial x_r} = \sum_r (b_{kr} \Gamma_{is}^k - b_{ks} \Gamma_{ir}^k) \quad \text{(Codazzi-Mainardi equation)}$$

63. If  $M$  is a Hausdorff  $C^\infty$  manifold,  $A \subset M$  is compact,  $B \subset M$  is open, and  $A \subset B$ , show there exists  $f \in C^\infty(M, \mathbb{R})$  with  $f(A) = 0$ ,  $f(M - B) = 1$ , and  $0 \leq f(M) \leq 1$ .





## 7. Operators on Forms and Integration

This chapter develops more structure on a manifold. To conserve space, the treatment is fairly blunt and many computational details are omitted. In the first four sections  $M$  is a  $C^\infty$   $n$ -manifold and  $A$  is an open set in  $M$ .

### 7.1 Exterior Derivative

For  $p \geq 0$  we define the exterior differentiation map  $\mathbf{d} : \Omega^p(A) \rightarrow \Omega^{p+1}(A)$  where  $\Omega^p(A)$  is the set of  $C^\infty$   $p$ -forms on  $A$ . If  $f \in \Omega^0(A)$  and  $X$  is a  $C^\infty$  field on  $A$ , then  $\mathbf{d}f(X) = Xf$ . For  $p > 1$ , letting  $w$  be a  $(p-1)$  form on  $A$  and  $X_1, \dots, X_p$  be  $C^\infty$  fields on  $A$ , then

$$\begin{aligned} \mathbf{d}w(X_1, \dots, X_p) = & \sum_{j=1}^p (-1)^{j+1} X_j w(X_1, \dots, \widehat{X}_j, \dots, X_p) + \\ & \sum_{i < j} (-1)^{i+j} w([X_i, X_j], X_1, \dots, \widehat{X}_i, \dots, \widehat{X}_j, \dots, X_p), \end{aligned} \tag{1}$$

where  $\widehat{X}$  indicates that the field  $X$  is omitted as an argument.

Notice that the definition is consistent with the partial definition in section 5.2. One proves that  $\mathbf{d}w$  is in  $\Omega^p(A)$  by using the characterization theorem in Chapter 4. We outline the argument. That  $\mathbf{d}w$  is linear with respect to addition is trivial. That  $\mathbf{d}w$  is alternating can be shown by switching two arguments and examining

the terms that don't immediately change signs (this must be done carefully). That  $\mathbf{d}w$  is linear over the ring  $\Omega^0(A)$  then need only be checked in one slot.

**Proposition 7.1.** The operation  $d$  has the following properties:

- (1)  $\mathbf{d}(w + v) = \mathbf{d}w + \mathbf{d}v$ , where  $w$  and  $v$  are in  $\Omega^p(A)$ .
- (2)  $\mathbf{d}(w \wedge v) = ((\mathbf{d}w) \wedge v) + (-1)^p(w \wedge \mathbf{d}v)$ , for  $w$  in  $\Omega^p(A)$  and  $v$  any form on  $A$ . (Any operator with this property is called an anti-derivation.)
- (3)  $\mathbf{d}^2 = \mathbf{d} \circ \mathbf{d} = 0$ .

*Proof.* Property (1) follows trivially from the definitions of  $\mathbf{d}$  and addition of functionals. For the other two properties we first obtain a local representation of  $\mathbf{d}$ . Let  $x_1, \dots, x_n$  be a coordinate system on an open set  $U$ , and let  $X_i = \frac{\partial}{\partial x_i}$ . Then on  $U$ , a  $(p-1)$ -form  $w$  may be represented by

$$w = \sum a_{i_1, \dots, i_{p-1}} \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_{p-1}},$$

where the sum is over all indices such that  $1 \leq i_j \leq n$  and  $i_1 < i_2 < \dots < i_{p-1}$  and  $a_{i_1, \dots, i_{p-1}} = w(X_{i_1}, \dots, X_{i_{p-1}})$ .

$$\mathbf{d}w = \sum da_{i_1, \dots, i_{p-1}} \wedge \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_{p-1}},$$

which is proved by applying both sides to  $(X_{k_1}, \dots, X_{k_p})$  for  $k_1 < k_2 < \dots < k_p$ . Since  $[X_r, X_s] = 0$ ,

$$\mathbf{d}w(X_{k_1}, \dots, X_{k_p}) = \sum_{j=1}^p (-1)^{j+1} X_{k_j} a_{k_1, \dots, \widehat{k_j}, \dots, k_p},$$

while

$$\begin{aligned} & \left[ \sum da_{i_1, \dots, i_{p-1}} \wedge \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_{p-1}} \right] (X_{k_1}, \dots, X_{k_p}) \\ &= \mathbf{d}a_{k_2, \dots, k_p}(X_{k_1}) - \mathbf{d}a_{k_1, \widehat{k_2}, \dots, k_p}(X_{k_2}) + \dots = \mathbf{d}w(X_{k_1}, \dots, X_{k_p}). \end{aligned}$$

To prove property (2), first let  $f$  and  $g$  be functions in  $\Omega^0(A)$  and note

$$d(fg) = (df)g + f(dg)$$

follows from the derivation property of vectors. Next observe that because of (1) and the local representation above, one need only verify (2) for forms of the type

$$w = f dx_1 \wedge \dots \wedge dx_p \text{ and } v = g dy_1 \wedge \dots \wedge dy_r,$$

where  $x_i$  and  $y_i$  are functions chosen from the members of a coordinate system. Then

$$w \wedge v = fg dx_1 \wedge \dots \wedge dx_p \wedge dy_1 \wedge \dots \wedge dy_r,$$

and

$$\begin{aligned} d(w \wedge v) &= d(fg) \wedge dx_1 \wedge \dots \wedge dx_p \wedge dy_1 \wedge \dots \wedge dy_r \\ &= (gdf + f dg) \wedge dx_1 \wedge \dots \wedge dx_p \wedge dy_1 \wedge \dots \wedge dy_r \\ &= dw \wedge v + (-1)^p w \wedge dv. \end{aligned}$$

For property (3) we first show  $d^2 f = 0$  for a  $C^\infty$  function  $f \in \Omega^0(A)$ . Locally,

$$df = \sum_{j=1}^n \frac{\partial f}{\partial x_j} dx_j,$$

so

$$\begin{aligned} d^2 f &= \sum_{i,j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j} dx_i \wedge dx_j \\ &= \sum_{i < j} \left[ \frac{\partial^2 f}{\partial x_i \partial x_j} - \frac{\partial^2 f}{\partial x_j \partial x_i} \right] dx_i \wedge dx_j \\ &= 0. \end{aligned}$$

For any  $w$  we may represent  $dw$  locally as a sum of products of

$df$ 's for functions  $f$ ; hence by (2) each term in  $d^2w$  has a factor  $d^2f = 0$ , so  $d^2w = 0$ .  $\square$

Letting  $\Omega(M) = \sum_{k=0}^n \Omega^k(M)$  be the direct sum of the modules of forms of homogeneous type, endowed with its exterior multiplication structure and exterior derivative operator  $d$ , one obtains a graded differential algebra which is called the *Cartan differential algebra* of  $M$ . If  $f : M \rightarrow N$  is  $C^\infty$ , then  $d \circ f^* = f^* \circ d$  on  $\Omega(N)$ , and it is sufficient to check this only on 0-forms and 1-forms.

There are other ways to define  $d$ , indeed one natural way is to define  $d$  via a local representation, get the desired properties, and then show it is independent of the local representation (see [Che46, p. 146]). Then the invariant formula we took as definition must be verified. Our treatment in this and the following sections is similar to that of [Pal54].

## 7.2 Contraction

Let  $X$  be a  $C^\infty$  vector field on the open set  $A$ . An operator  $C_X$ , called *contraction by  $X$* , which maps  $\Omega^p(A)$  into  $\Omega^{p-1}(A)$  is defined as follows: (a) if  $f \in \Omega^0(A)$ , let  $C_X f = 0$ , and (b) if  $w \in \Omega^p(A)$  for  $p > 0$ , let

$$(C_X w)(X_1, \dots, X_{p-1}) = w(X, X_1, \dots, X_{p-1})$$

.

**Proposition 7.2.** The operator  $C$  has the following properties:

- (1)  $(C_X)^2 = 0$ .
- (2)  $C_X(w + v) = C_X w + C_X v$ ,
- (3)  $C_{X+Y} = C_X + C_Y$ ,
- (4)  $C_{fX} = fC_X$ ,
- (5)  $C_X(w \wedge z) = (C_X w) \wedge z + (-1)^p(w \wedge C_X z)$ ,

for  $f$  in  $\Omega^0(A)$ ,  $X$  and  $Y$  in  $\mathbf{T}^{1,0}A$ ,  $w$  and  $v$  in  $\Omega^p(A)$ , and  $z$  in  $\Omega^q(A)$ .

*Proof.* Properties (1) through (4) are trivial. Property (5) follows by induction on  $p$ , and it is sufficient to prove it when  $w$  is a product of  $p$  1-forms by the local representation of forms.  $\square$

The operator  $C_X$  can be defined on covariant tensors and mixed tensors in an obvious way (with only (2), (3), and (4) valid in general), and one can let  $C_X$  be zero on pure contravariant tensors. Properties (3) and (4) indicate  $C$  is a tensor map (an anti-derivation valued 1-form of degree  $-1$  on  $\Omega(A)$ ).

There is another form of “contraction” induced by the natural identification of tensors of type (1,1) and linear maps. Let  $W$  be an  $n$ -dimensional vector space over  $\mathbb{R}$ . For  $r > 0$ ,  $s > 0$ ,  $1 \leq i \leq r$ ,  $1 \leq j \leq s$  define  $\text{tr}^{i,j} : \mathbf{T}^{r,s}W \rightarrow \mathbf{T}^{r-1,s-1}W$  by taking  $\theta$  in  $\mathbf{T}^{r,s}W$ ,  $w_1, \dots, w_{r-1}$  in  $W^*$ , and  $X_1, \dots, X_{s-1}$  in  $W$  and letting

$$(\text{tr}^{i,j}\theta)(w_1, \dots, w_{r-1}, X_1, \dots, X_{s-1}) = \sum_{k=1}^n \theta(w_1, \dots, w_{i-1}, z_k, w_i, \dots, w_{r-1}, X_1, \dots, X_{j-1}, Z_k, X_j, \dots, X_{s-1}) \quad (2)$$

where  $Z_1, \dots, Z_n$  is a base of  $W$  and  $z_1, \dots, z_n$  the dual base of  $W^*$ . One checks easily that  $\text{tr}^{i,j}\theta$  is well-defined independently of the particular base used. If  $\theta \in \mathbf{T}^{1,1}W$ , let  $\text{tr}^{1,1}\theta = \text{tr}\theta$ . The above operator induces an operator

$\text{tr}^{i,j} : \mathbf{T}^{r,s}A \rightarrow \mathbf{T}^{r-1,s-1}A$  for an open set  $A$  in  $M$ .

### 7.3 Lie Derivative

Let  $X$  be a  $C^\infty$  vector field on the open set  $A$ . An operator  $\mathcal{L}_X$ , called the *Lie derivative via  $X$* , which maps  $\mathbf{T}^{r,s}A$  into itself, is defined as follows:

- (a) if  $f \in \Omega^0(A)$ ,  $\mathcal{L}_X Y = Xf$ ;  
 (b) if  $Y \in \mathbf{T}^{1,0}A$ ,  $\mathcal{L}_X Y = [X, Y]$ ;  
 (c) if  $w \in \mathbf{T}^{0,1}A$ ,  $(\mathcal{L}_X w)(Y) = Xw(Y) - w([X, Y])$ ;  
 (d) if  $\theta$  in  $\mathbf{T}^{r,s}A$ ,  $w_1, \dots, w_t$  in  $\mathbf{T}^{0,1}A$ , and  $Y_1, \dots, Y_s$  in  $\mathbf{T}^{1,0}A$ , then  $\mathcal{L}_X \theta$  is defined by solving for it in the equation

$$\begin{aligned} \mathcal{L}_X[\theta(w_1, \dots, w_r, Y_1, \dots, Y_s)] &= (\mathcal{L}_X \theta)(w_1, \dots, Y_s) \\ &+ \theta(\mathcal{L}_X w_1, w_2, \dots, Y_s) + \dots + \theta(w_1, \dots, Y_{s-1}, \mathcal{L}_X Y_s). \end{aligned} \quad (3)$$

We call  $\mathcal{L}_X$  a complete derivation because of the property (d) and note all terms in 3 are well-defined by (a), (b), and (c) except the  $\mathcal{L}_X \theta$  term (indeed, (c) is “defined” by (d)). One shows  $\mathcal{L}_X \theta$  is a tensor by checking the linearity over  $\Omega^0(A)$ .

**Proposition 7.3.** The operator  $\mathcal{L}_X$  has the following properties:

- (1)  $\mathcal{L}_X$  preserves forms,
- (2)  $\mathcal{L}_X(w + z) = \mathcal{L}_X w + \mathcal{L}_X z$ ,
- (3)  $\mathcal{L}_X(w \otimes v) = (\mathcal{L}_X w) \otimes v + w \otimes \mathcal{L}_X v$ ,
- (4)  $\mathcal{L}_X(\alpha \wedge \beta) = (\mathcal{L}_X \alpha) \wedge \beta + \alpha \wedge \mathcal{L}_X \beta$ ,

where  $w$  and  $z$  are tensors of the same type,  $v$  is any tensor, and  $\alpha$  and  $\beta$  are forms.

*Proof.* An exercise (for (4) use  $\mathcal{L}_X[(\alpha \otimes \beta)^\pi] = [\mathcal{L}_X(\alpha \otimes \beta)]^\pi$ ). □

There is a more geometric definition of the Lie derivative  $\mathcal{L}_X$  on covariant tensors which we now discuss. Suppose the vector field  $X$  is defined and  $C^\infty$  on all of  $M$ . For each  $m$  in  $M$  let  $f_m(t)$  be the integral curve of  $X$  (section 1.5) through  $m$  with  $f_m(0) = m$ . We know  $f_m$  defined for  $t$  in a neighborhood of zero, but suppose each  $f_m$  is defined for all  $t$  and  $\mathbb{R}$ . Then for each  $t$  in  $\mathbb{R}$  we could define a map  $F_t : M \rightarrow M$  by  $F_t(m) = f_m(t)$ , with the properties  $F_t \circ F_s = F_{t+s}$  and

$F : M \times \mathbb{R} \rightarrow M$  by  $F(m, t) = F_t(m)$  would be  $C^\infty$  (from the fact that  $X$  was  $C^\infty$  and the  $C^\infty$  dependence of solutions of ordinary differential equations upon initial conditions). Each  $F_t$  would be a diffeo, since  $(F_t)^{-1} = F_{-t}$  and  $F_0$  is the identity map. A map  $F$  with the above properties is called a 1-parameter group of differentiable transformations of  $M$ , and  $X$  is called its infinitesimal generator.

In general  $f_m$  is not defined for all  $t$ , but one does obtain a local 1-parameter group of local transformations in a neighborhood of each  $m$  in  $M$ ; i.e., for each  $m$  in  $M$  there is a neighborhood  $U$  of  $m$ , a real number  $b > 0$ , and a map  $F : U \times (-b, b) \rightarrow M$  such that

- (1)  $F$  is  $C^\infty$ ,
- (2) for  $t$  in  $(-b, b)$ ,  $F_t : U \rightarrow F_t(U)$  is a diffeo,
- (3) for  $t, s$ , and  $t + s$  in  $(-b, b)$ ,  $F_t \circ F_s = F_{t+s}$ , and
- (4) for fixed  $p \in U$ ,  $f_p(t) = F_t(p)$  is an integral curve of  $X$ .

For more details see [Pal54]; [Pal57] and [Nom21, p. 5].

**Lemma 7.4.** Let  $Y$  be a  $C^\infty$  field in a neighborhood of  $m$  in  $M$ . We choose  $U$  and  $b$  in the preceding paragraph to be sufficiently small so the image of  $F$  is contained in the domain of  $Y$ . Then

$$[X, Y]_m = \lim_{t \rightarrow 0} \frac{(F_{-t})_* Y_{F(m,t)} - Y_m}{t}.$$

*Proof.* See [Nom21, p. 8]. □

Assuming lemma 7.4, which gives us another geometric interpretation of the bracket, it is trivial to show the following lemma.

**Lemma 7.5.** Let  $w$  be a  $C^\infty$   $p$ -form at  $m$ . Then

$$(\mathcal{L}_X w)_m = \lim_{t \rightarrow 0} \frac{(F_t^* w)_m - w_m}{t}$$

where

$$(F_t^* w)_m(Y_1, \dots, Y_p) = w_{F(m,t)}((F_t)_* Y_1, \dots, (F_t)_* Y_p)$$

The following is a useful relation between  $\mathbf{d}$ ,  $\mathcal{L}_X$ , and  $C_X$ .

**Proposition 7.6.** If  $X$  is a  $C^\infty$  field on  $A$ , then  $\mathcal{L}_X = \mathbf{d} \circ C_X + C_X \circ \mathbf{d}$  when applied to  $C^\infty$  forms on  $A$ .

*Proof.* We verify this equality on functions (0-forms) and 1-forms. This is sufficient to prove the proposition, since locally a form is a sum of products of functions and 1-forms, and the operators which we equate above are both derivations; hence their value on any form is determined by the values on functions and 1-forms.

For  $f \in \Omega^0(A)$ ,

$$\mathbf{d}C_X(f) + C_X\mathbf{d}(f) = 0 + \mathbf{d}f(X) = Xf = \mathcal{L}_X f.$$

For  $w \in \Omega^1(A)$ ,

$$\begin{aligned} (\mathbf{d}C_X + C_X\mathbf{d}w)(Y) &= Yw(X) + \mathbf{d}w(X, Y) \\ &= Yw(X) + Xw(Y) - Yw(X) - w([X, Y]) \\ &= (\mathcal{L}_X w)(Y) \end{aligned}$$

□

## 7.4 General Covariant Derivative

Let  $\mathbf{D}$  be a connexion on  $M$ , and let  $X$  be a  $C^\infty$  field on the open set  $A$ . An operator  $\mathbf{D}_X$ , called the covariant derivative via  $X$ , which maps  $\mathbf{T}^{r,s}A$  into itself, is defined by using the recipe for defining  $\mathcal{L}_X$ . The definition of  $\mathbf{D}_X$  proceeds exactly as the definition for  $\mathcal{L}_X$  except for (b), and if  $Y$  in  $\mathbf{T}^{1,0}A$ ,  $\mathbf{D}_X Y$  is given by the connexion  $\mathbf{D}$  (see section 5.1).

When  $\mathbf{D}_X$  is substituted for  $\mathcal{L}_X$  in proposition 7.3 of the previous section, one obtains valid properties for  $\mathbf{D}_X$ .

An operator  $\Delta$ , called the general covariant derivative operator, which maps  $\mathbf{T}^{r,s}A$  into  $\mathbf{T}^{r,s+1}A$  is induced by  $\mathbf{D}$ . If  $\theta$  is in  $\mathbf{T}^{r,s}A$ ,



$w_1, \dots, w_r$  are in  $\mathbf{T}^{0,1}A$ , and  $Y_1, \dots, Y_{s+1}$  are in  $\mathbf{T}^{1,0}A$ , then

$$(\Delta\theta)(w_1, \dots, w_r, Y_1, \dots, Y_{s+1}) = (\mathbf{D}_{Y_{s+1}}\theta)(w_1, \dots, w_r, Y_1, \dots, Y_s) \quad (4)$$

That  $\Delta\theta$  is a tensor is left as a problem. If  $\theta$  and  $\phi$  are tensors of the same type, then  $\Delta(\theta + \phi) = \Delta\theta + \Delta\phi$ , but  $\Delta$  is not a tensor. (see problem 64)

If  $1 \leq i \leq p$  and  $1 \leq j \leq q$ , then

$$\Delta \circ \text{tr}^{i,j} = \text{tr}^{i,j} \circ \Delta \quad (5)$$

on  $\mathbf{T}^{p,q}A$ .

An operator  $\text{div}$ , called the *divergence*, which maps  $\mathbf{T}^{r,s}A$  into  $\mathbf{T}^{r-1,s}A$ , for  $r > 0$  and  $s \geq 0$ , is defined by  $\text{div} = \text{tr}^{r,s+1} \circ \Delta$ . We write  $\text{div}\theta = \text{tr}(\Delta\theta)$ , where we assume the trace is taken on the last covariant slot and the last contravariant slot. A tensor  $\theta$  is *conservative* if  $\text{div}\theta = 0$ .

The *Riemann-Christoffel curvature tensor* of type  $(1,3)$  is the tensor  $K \in \mathbf{T}^{1,3}A$  defined by

$$K(w, X, Y, Z) = w(R(Y, Z)X) \quad (6)$$

for  $w \in \mathbf{T}^{0,1}A$  and  $X, Y, Z \in \mathbf{T}^{1,0}A$ . The *second Bianchi identity* is the equation

$$(\Delta K)(w, X, Y, Z, W) + (\Delta K)(w, X, W, Y, Z) + (\Delta K)(w, X, Z, W, Y) = 0 \quad (7)$$

which is valid if  $\mathbf{D}$  is symmetric, and it is proved by noting the expression

$$\mathbf{D}_W(R(Y, Z)X) - R(Z, [Y, W])X - R(Y, Z)(\mathbf{D}_W X), \quad (8)$$

when written on three lines, permuting  $W, X, Z$  cyclically from line to line, and then adding the three lines, yields zero.

The *Ricci tensor* is the 2-covariant tensor

$$\text{Ric}(X, Y) = (\text{tr}^{1,2}K)(X, Y) = -(\text{tr}^{1,3}K)(X, Y) \quad (7.1)$$

(and this is the negative of the “classical” Ricci tensor). Notice  $(\text{tr}^{1,1}K)(X, Y) = \text{tr}R(X, Y)$ . The Ricci curvature of a vector  $X$  is the number  $\text{Ric}(X, X)$  (and this agrees with the “classical” Ricci curvature). If  $\mathbf{D}$  is symmetric, the first Bianchi identity implies

$$\text{Ric}(X, Y) = \text{Ric}(Y, X) + \text{tr}R(X, Y). \quad (10)$$

If  $\mathbf{D}$  is Riemannian, then  $R(X, Y)$  is skew-symmetric by (c), so  $\text{Ric}$  is symmetric. Hence there exists a self-adjoint linear map  $R^*$ , called the Ricci map, defined on each  $\mathbf{T}_m M$  with  $\text{Ric}(X, Y) = \langle R^*(X), Y \rangle$ ; indeed

$$R^*(X) = \sum_{j=1}^n R(X, Z_j)Z_j \quad (11)$$

for an orthonormal base  $Z_1, Z_2, \dots, Z_n$ . By (11),  $R^*$  is  $C^\infty$ . The scalar curvature  $S(m)$  at each  $m \in M$  is defined by  $S(m) = \text{tr}(R^*)_m$ .

A (semi-) Riemannian metric induces many operations called “raising” and “lowering” of indices which we now explain. The non-singular metric tensor induces a non-singular linear map  $G$  of  $\mathbf{T}_m M$  onto  $\mathbf{T}_m^* M$  for each  $m$ , i.e., if  $X$  in  $\mathbf{T}_m M$ , then  $G(X)(Y) = \langle X, Y \rangle$ . We let  $G_*$  denote the inverse map of  $\mathbf{T}_m^* M$  onto  $\mathbf{T}_m M$ . If  $w \in \mathbf{T}_m^* M$ , then  $\langle G_* w, X \rangle = w(X)$ . If  $1 \leq i \leq r, 1 \leq j \leq s+1$ , and  $\theta$  is in  $\mathbf{T}^{r,s}$  define  $G^{i,j}\theta$  in  $\mathbf{T}^{r-1,s+1}$  by

$$\begin{aligned} (G^{i,j}\theta)(w_1, \dots, w_{r-1}, X_1, \dots, X_{s+1}) \\ = \theta(w_1, \dots, w_{i-1}, G(X_j), w_i, \dots, w_{r-1}, X_1, \dots, \widehat{X}_j, \dots, X_{s+1}). \end{aligned} \quad (12)$$

Similarly, define  $G_*^{i,j} : \mathbf{T}^{r,s} \rightarrow \mathbf{T}^{r+1,s-1}$  for  $1 \leq i \leq r+1$  and  $1 \leq j \leq s$  by taking the form in the  $i^{\text{th}}$  covariant slot (of the new tensor); applying  $G_*$ , and inserting it into the  $j^{\text{th}}$  contravariant slot (of the old tensor). Thus  $G_*^{1,1} = G$  on  $\mathbf{T}^{1,0}$ , and the  $(1,1)$ -tensor  $\bar{R}$  associated with  $R^*$  is given by  $\bar{R} = G_*^{1,1}\text{Ric}$  (where  $\bar{R}(w, X) = w(R^* X)$ ). If  $f$  is in

$C^\infty(M, \mathbb{R})$ , the gradient field of  $f$  is the field  $\text{grad } f = G_*(df)$  and the Laplacian of  $f$  is the function  $\text{del } f = \text{div}(\text{grad } f)$ ; (sometimes the notation  $\text{del } f = \Delta f$  is used).

The operators  $G^{i,j}$  and  $G_*^{i,j}$  commute with  $\Delta$  when possible, i.e.,

$$\Delta \circ G^{i,j} = G^{i,j} \circ \Delta \text{ on } \mathbf{T}^{r,s} \text{ if } j \leq s + 1 \text{ and} \quad (13)$$

$$\Delta \circ G_*^{i,j} = G_*^{i,j} \circ \Delta \text{ on } \mathbf{T}^{r,s} \text{ if } i \leq r + 1. \quad (14)$$

As an example of the use of these operations we prove that

$$\Delta S = 2 \text{div } \bar{R} \quad (15)$$

which is used in general relativity. Let  $Z_1, \dots, Z_n$  be an orthonormal base of  $\mathbf{T}_m M$  and  $w_1, \dots, w_n$  be the dual base. The second Bianchi identity implies

$$\sum_{i,j} [\Delta K(w_j, Z_i, Z_j, X) + \Delta K(w_i, Z_i, X, Z_j, Z_i) + K(w_j, Z_i, Z_i, X, Z_j)] = 0$$

The first term of the sum gives  $(\Delta S)(X)$ , while the other two each give  $(-\text{div } \bar{R})(X)$ . For

$$\begin{aligned}
 (\Delta S)(X) &= (\Delta \mathbf{tr}^{1,1} G_*^{1,1} \mathbf{tr}^{1,2} K)(X) \\
 &= (\mathbf{tr}^{1,1} G_*^{1,1} \mathbf{tr}^{1,2} \Delta K)(X) \\
 &= \sum_{i,j} \Delta K(w_j, Z_i, Z_j, Z_i, X), \\
 (\mathbf{div} \bar{R})(X) &= (\mathbf{tr}^{1,2} \Delta G_*^{1,1} \mathbf{tr}^{1,2} K)(X) \\
 &= \sum_{i,j} \Delta K(w_j, Z_i, X, Z_j, Z_i), \text{ and} \\
 (\Delta K)(w_j, Z_i, Z_i, X, Z_j) &= (\Delta G^{1,1} K)(Z_j, Z_i, Z_i, X, Z_j) \\
 &= (\Delta G^{1,1} K)(Z_i, X, Z_j, Z_i, Z_j) \\
 &= -(\Delta G^{1,1} K)(Z_i, Z_i, X, Z_j, Z_j) - (\Delta G^{1,1} K)(Z_i, Z_j, Z_i, X, Z_j) \\
 &= -(\mathbf{div} \bar{R})(X)
 \end{aligned}$$

by (c) and (a') in section 6.2.

## 7.5 Integration of Forms and Stokes' Theorem

One integrates  $p$ -forms over  $p$ -chains, or singular  $p$ -chains, which we now define. Let  $I^p = \{a \in \mathbb{R}^p : 0 \leq a_i \leq 1\}$  denote the unit  $p$ -square for  $p > 0$ , and  $I^0 = \{0 \in \mathbb{R}\}$ . A  $C^\infty$   $p$ -cube on  $M$  is an  $M$ -valued  $C^\infty$  function defined on an open neighbourhood of the unit  $p$ -square  $I^p$  in  $\mathbb{R}^p$ . A real  $C^\infty$   $p$ -chain  $c$  is a finite formal linear combination of  $C^\infty$   $p$ -cubes with real coefficients, thus  $c = r_1 \sigma_1 + r_2 \sigma_2 + \dots + r_k \sigma_k$  where  $r_j \in \mathbb{R}$  and  $\sigma_j$  are  $C^\infty$   $p$ -cubes. The set  $C_p(M, \mathbb{R})$  of all real  $C^\infty$   $p$ -chains is an abelian group (actually an  $\mathbb{R}$ -module) where one defines addition by adding the coefficients of corresponding  $p$ -cubes.

There are fancier ways of defining  $C_p(M, \mathbb{R})$ . Let  $Q_p$  be the set of  $C^\infty$   $p$ -cubes on  $M$ . Then  $C_p(M, \mathbb{R})$  is isomorphic to set of all functions mapping  $Q_p$  into  $\mathbb{R}$  which are zero except on a finite number of elements, and the addition and scalar multiplication structure on this function space is obvious. Similarly, one could define  $C(M, \mathbb{Z})$ ,

the set of integral  $C^\infty$   $p$ -chains or  $C^\infty$   $p$ -chains over the integers. Then  $C_p(M, \mathbb{R}) = \mathbb{R} \otimes_{\mathbb{Z}} C_p(M, \mathbb{Z})$ . More generally one could define  $C^\infty$   $p$ -chains over any ring  $A$  with an identity element, and then by using the tensor product obtain the  $A$ -module of  $C^\infty$   $p$ -chains on  $M$  over any  $A$ -module. There are corresponding groups obtained from  $C^r$   $p$ -chains for any integer  $r \geq 0$ . These groups are fundamental objects of the cubical singular homology theory for  $M$  and are studied in algebraic topology, (see Eilenberg and Steenrod). Because of our differential geometry bias, we restrict ourselves to real  $C^\infty$   $p$ -chains, and let  $\mathbf{C}_p = \mathbf{C}_p(M, \mathbb{R})$ .

The support of a  $p$ -cube  $\sigma$  is the set  $|\sigma| = \sigma(I^p)$ , the image of  $I^p$  under  $\sigma$ . The support of a  $p$ -chain  $c$  is the set  $|c| = \bigcup_i |\sigma_i|$  for  $\sigma_i$  in  $c$ , where we say  $\sigma_i \in c$  if the coefficient of  $\sigma_i$  is non-zero, i.e., adopting the functional viewpoint  $c(\sigma_i) \neq 0$  iff  $\sigma_i \in c$ .

To define the boundary map  $\partial : C_p \rightarrow C_{p-1}$ , define maps  $\alpha_i^1$  and  $\alpha_i^0$  from  $I^{p-1}$  into  $I^p$  for  $i = 1, \dots, p$  by

$$\alpha_i^\epsilon(t_1, \dots, t_{p-1}) \in (t_1, \dots, t_{i-1}, \epsilon, t_i, \dots, t_{p-1}) \quad (16)$$

where  $\epsilon = 1$  or  $0$ . If  $\sigma$  in  $Q_p$ , define

$$\partial\sigma = \sum_1^p (-1)^{i+1} (\sigma \circ \alpha_i^1 - \sigma \circ \alpha_i^0),$$

and call the  $(n-1)$ -cubes  $\sigma \circ \alpha_i^1$  and  $\sigma \circ \alpha_i^0$  faces of  $\sigma$ . We extend  $\partial$  to all of  $C_p$  by demanding it be linear, i.e.,

$$\partial(c_1 + c_2) = \partial c_1 + \partial c_2 \quad \text{and} \quad \partial(rc) = r\partial c \quad \text{for } r \in \mathbb{R}$$

. A straightforward computation shows  $\partial^2 = 0$ .

For  $p > 0$ , let  $\sigma$  be a  $C^\infty$   $p$ -cube, let  $w$  be a  $p$ -form, and let  $u_1, \dots, u_p$  be the natural coordinate function on  $\mathbb{R}^p$ . Since  $\sigma^*w$  is a  $p$ -form on a neighborhood of  $I^p$ , we may define a  $C^\infty$  function  $f$  on  $I^p$  by

$\sigma^*w = f du_1 \wedge du_2 \wedge \dots \wedge du_p$ . Then

$$\int_{\sigma} w \equiv \int_I p^*w \equiv \int_I p \tag{17}$$

where the integral on the right is the standard Riemann integral of  $f$  over  $I^p$  developed in advanced calculus. For a  $p$ -chain

$$c = \sum_1^{k_i} \sigma_i \implies \int_c w = \sum_1^k r_i \int_{\sigma_i} w;$$

thus for fixed  $w$ , the integral over  $w$  is an  $\mathbb{R}$ -homomorphism of  $C_p$  into  $\mathbb{R}$ . Since  $\sigma^*$  is linear, it is trivial that  $\int_c (w_1 + w_2) = \int_c w_1 + \int_c w_2$  for  $p$ -forms  $W_i$  and a  $p$ -chain  $c$ .

For  $p = 0$ , let  $f$  be a function on  $M$  and  $\sigma_m$  the  $\mathbf{0}$ -cube with  $\sigma_m(0) = m$ , then

$$\int_{\sigma} f = f(m) = \sigma_m^* f(0),$$

and we extend the integral of  $f$  over any real 0-chain to be linear (as extended above).

Let  $C^p = \text{Hom}_{\mathbb{R}}(C_p, \mathbb{R})$ , which is the  $\mathbb{R}$ -module of all  $\mathbb{R}$ -linear homomorphisms of  $C_p$  into  $\mathbb{R}$ . The set  $C^p$  is called the module of real  $C^\infty$   $p$ -cochains of  $M$ . The adjoint  $\delta$  of the boundary operator  $\partial$  is called the coboundary operator and is defined by  $\delta f(c) = f(\partial c)$  for  $p$ -cochain  $f$  and a  $(p + 1)$ -chain  $c$ . Thus  $\delta : C^p \rightarrow C^{p+1}$  and  $\delta^2 = 0$ .

We define the Stokes' map  $S : \Omega^p(M) \rightarrow C^p$  which maps  $p$ -forms on  $M$  into  $C^\infty$   $p$ -cochains on  $M$  by  $[S(w)](c) = \int_c w$ , for  $c$  in  $C_p$ . The following theorem shows the Stokes' map commutes with the differential coboundary operator, i.e.,  $S \circ d = \delta \circ S$ .

**Theorem 7.7 (Stokes' Theorem).** Let  $w$  be a  $C^\infty$   $p$ -form and  $\sigma$  be a  $C^\infty$   $(p + 1)$ -cube, then

$$\int_{\sigma} dw = \int_{\partial\sigma} w \tag{18}$$

---

Section 7.5: Integration of Forms and Stokes' Theorem

---

**Proof.** Define  $C^\infty$  functions  $a_1, \dots, a_{p+1}$  on  $I^{p+1}$  by

$$\sigma^* w = \sum_1^{p+1} a_i \mathbf{d}u_1 \wedge \mathbf{d}u_2 \wedge \dots \wedge \widehat{\mathbf{d}u_i} \wedge \dots \wedge \mathbf{d}u_{p+1}.$$

Then

$$\begin{aligned} \mathbf{d}(\sigma^* w) &= \sum_{i=1}^{p+1} \left( \sum_{i=1}^{p+1} \frac{\partial a_i}{\partial u_j} du_j \right) \wedge \mathbf{d}u_1 \wedge \dots \wedge \widehat{\mathbf{d}u_i} \wedge \dots \wedge \mathbf{d}u_{p+1} \\ &= \left[ \sum_{i=1}^{p+1} (-1)^{i+1} \frac{\partial a_i}{\partial u_i} \right] \mathbf{d}u_1 \wedge \dots \wedge \mathbf{d}u_{p+1}. \end{aligned}$$

Hence

$$\begin{aligned} \int_\sigma \mathbf{d}w &= \int_{I^{p+1}} \sigma^* \mathbf{d}w = \int_{I^{p+1}} \mathbf{d}\sigma^* w = \int_{I^{p+1}} \left[ \sum_1^{p+1} (-1)^{i+1} \frac{\partial a_i}{\partial u_i} \right] \\ &= \sum_1^{p+1} (-1)^{i+1} \left[ \int_0^1 \int_0^1 \dots \int_0^1 \frac{\partial a_i}{\partial u_i} \mathbf{d}u_1 \mathbf{d}u_2 \dots \mathbf{d}u_{p+1} \right] \\ &= \sum_1^{p+1} (-1)^{i+1} \int_{I^p} (a_i \circ \alpha_i^1 - a_i \circ \alpha_i^0) \end{aligned}$$

where we use Fubini's theorem and integrate first with respect to  $i$ th coordinate to obtain the last equality.

For the other side we must compute  $\int_{\sigma \circ \alpha_i^\epsilon} w = \int_{I^p} (\alpha_i^\epsilon)^* \circ \sigma^*(w)$  for  $\epsilon = 0$  or  $1$ . Notice

$$(\alpha_i^\epsilon)^* \mathbf{d}u_j = \mathbf{d}((\alpha_i^\epsilon)^* u_j) = \mathbf{d}(u_j \circ \alpha_i^\epsilon) = \begin{cases} \mathbf{d}u_j & j < i \\ 0 & j = i \\ \mathbf{d}u_{j-1} & j > i \end{cases}$$

Thus  $(\alpha_i^\epsilon)^* \sigma^* w = (a_i \circ \alpha_i^\epsilon) \mathbf{d}u_1 \wedge \dots \wedge \mathbf{d}u_p$  and

$$\begin{aligned} \int_{\partial\sigma} w &= \sum_{i=1}^{p+1} (-1)^{i+1} \left[ \int_{\sigma \circ \alpha_i^1} w - \int_{\sigma \circ \alpha_i^0} w \right] \\ &= \sum_{i=1}^{p+1} (-1)^{i+1} \int_{I^p} (a_i \circ \alpha_i^1 - a_i \circ \alpha_i^0) \end{aligned}$$

which proves the desired equality.  $\square$

We remark that Stokes' theorem is simply a generalized "fundamental theorem of calculus." Let  $f : M \rightarrow M'$  be a  $C^\infty$  map, let  $w$  be a  $p$ -form on  $M'$  and  $\sigma$  a  $p$ -cube in  $M$ , then it is trivial to show  $\int_{f \circ \sigma} w = \int_{\sigma} f^* w$ , which is essentially the classical substitution rule that deals with the behavior of integrals with respect to mappings.

The Stokes' map induces a map at the cohomology level that yields an algebra-isomorphism of the differential cohomology groups of a manifold with the real singular cohomology groups. This fact is called the de Rham theorem (see [Wei52] and problem 71).

## 7.6 Integration in a Riemannian Manifold

Let  $M$  be a Riemannian manifold, let  $\sigma$  be a  $C^\infty$  curve in  $M$ , and let  $f$  be a real valued  $C^\infty$  function on the image of  $\sigma$ , i.e., let  $f \circ \sigma$  be  $C^\infty$ . Consider a "piece" of  $\sigma$ , which we assume to be parameterized by arc length on the interval  $[a, b]$ , and define

$$\int_{\sigma|_{[a,b]}} f = \int_a^b f \circ \sigma(s) ds \tag{19}$$

where  $\sigma|_{[a,b]}$  denotes the restriction of  $\sigma$  to the interval  $[a, b]$ . Call the integral just defined the integral of  $f$  over  $\sigma$  restricted to  $[a, b]$  and when the interval is understood, we write simply  $\int_{\sigma} f$ . If  $f$  is a  $C^\infty$  real valued function  $f$  defined on a broken  $C^\infty$  curve  $\sigma$ , we define  $\int_{\sigma} f$  to be the sum of the integrals of  $f$  over the finite number of  $C^\infty$  sub-curves determining  $\sigma$ . Notice that by assuming



$\sigma$  parametrized by arc length we are integrating over oriented or directed curves.

We wish to integrate real valued  $C^\infty$  functions over other subsets of  $M$ , and in some cases over  $M$  itself. This could be accomplished by using the Riemannian metric to define a measure on  $M$ , but for our purposes we need not be so general. First we define orientable manifolds and then utilize the theory developed above for integrating forms over chains.

An  $n$ -dimensional manifold  $M$  is *orientable* if there is a non-vanishing  $C^\infty$   $n$ -form  $w$  on  $M$ . When  $M$  is orientable and we have selected  $w$ , we say  $M$  is *oriented (by  $w$ )* and  $w$  is an *orientation* of  $M$ . If  $M$  is oriented by  $w$ , then an ordered base  $e_1, \dots, e_n$  of  $\mathbf{T}_m M$  is *positively oriented* if  $w_m = b w_1 \wedge \dots \wedge w_n$  where  $b > 0$  and  $w_i$  are the 1-forms dual to  $e_j$ . We say  $M$  is *non-orientable* when  $M$  is not orientable. If  $M$  is oriented and  $e_1, \dots, e_n$  a positively oriented base of  $M$ , then one verifies easily that a base  $f_1, \dots, f_n$  of  $M$  is positively oriented if and only if  $\det(b_{ij}) > 0$  where  $f_j = \sum_i b_{ij} e_i$ .

For example,  $\mathbb{R}^n$  is orientable, and we orient it by choosing  $W = du_1 \wedge \dots \wedge du_n$  where  $u_i$  are the natural coordinate functions. It is a topological result that any complete (or closed) hypersurface in  $\mathbb{R}^n$  is orientable.

Let  $M$  and  $M'$  be oriented  $n$ -manifolds. A non-singular  $C^\infty$  map  $f$  of  $M$  into  $M'$  is *orientation preserving* if  $f_*$  maps a positively oriented base onto a positively oriented base.

Let  $M$  be an oriented Riemannian  $n$ -manifold. For  $m$  in  $M$  let  $e_1, \dots, e_n$  be a positively oriented orthonormal base of  $\mathbf{T}_m M$  with dual base  $w_1, \dots, w_n$ . Define the  $n$ -form  $v$  by  $v_m = w_1 \wedge \dots \wedge w_n$ . The form  $v$  is a well-defined (independent of the particular base)  $C^\infty$   $n$ -form on  $M$  called the *volume element*.

A major problem now confronts us: the problem of “triangulating” or “cubulating” a manifold. This is a theory for breaking the manifold into “nice pieces” over which one can integrate functions. For this purpose we define fundamental  $n$ -chains. Let  $\text{Int}(A)$  denote

the interior of a set  $A$ .

Let  $M$  be an oriented  $C^\infty$   $n$ -manifold. A fundamental  $n$ -chain in  $M$  is a chain  $c = \sigma_1 + \dots + \sigma_k$  such that:

- (1) each  $\sigma_i$  is an  $n$ -cube that is an orientation preserving diffeo onto its image;
- (2)  $\text{Int}(|\sigma_i|) \cap \text{Int}(|\sigma_j|)$  is empty for  $i \neq j$ .

Figure 7.1 gives a schematic diagram of a fundamental 2-chain (with the images of the faces of the canonical 2-cube numbered).




Figure 7.1: A Fundamental 2-Chain

If  $M$  is an oriented Riemannian  $n$ -manifold,  $c$  is an  $n$ -chain, and  $f$  is a  $C^\infty$  real valued function whose domain contains  $|c|$ , then define

$$\int_c f = \int_c f v \tag{20}$$

where  $v$  is the volume element on  $M$ . Let a subset  $A$  of  $M$  be fundamental if there exists fundamental  $n$ -chain  $c$  with  $|c| = A$ . Notice a fundamental set is compact.

**Proposition 7.8.** If  $c$  and  $\tau$  are two fundamental  $n$ -chains with  $|c| = |\tau| = A$ , and  $f$  is a  $C^\infty$  function whose domain contains  $A$ , then  $\int_c f v = \int_\tau f v$ . Thus define  $\int_A f = \int_c f v$ .

**Proof (King Lee).** Let  $c = \sigma_1 + \dots + \sigma_r$  and  $\tau = \gamma_1 + \dots + \gamma_s$ , and throughout this proof let  $1 \leq i \leq r$  and  $1 \leq j \leq s$ . If  $A_{ij} = |\sigma_i| \cap |\gamma_j|$ , let  $B_{ij} = (\sigma_i)^{-1}(A_{ij})$  and  $C_{ij} = (\gamma_j)^{-1}(A_{ij})$ . Then  $\gamma_j^{-1} \circ \sigma_i$  is a diffeo of  $B_{ij}$  onto  $C_{ij}$  and

$$\int_{B_{ij}} (\sigma_i)^* f v = \int_{B_{ij}} (\sigma_i)^* (\gamma_j \circ \gamma_j^{-1})^* f v = \int_{B_{ij}} (\gamma_j^{-1} \circ \sigma_i)^* f v = \int_{C_{ij}} (\gamma_j)^* f v$$

Hence

$$\int_c f v = \sum_i \int_{\sigma_i} f v = \sum_{i,j} \int_{B_{ij}} (\sigma_i)^* f v = \sum_{i,j} \int_{C_{ij}} (\gamma_j)^* f v = \int_\tau f v.$$

□

If  $M$  is a compact oriented  $n$ -manifold, then  $M$  is a fundamental set (this is hard; see [Cai]). Thus if  $M$  is a compact oriented Riemannian manifold and  $f$  is a  $C^\infty$  real valued function on  $M$ , then  $\int_M f$  is well-defined. To handle the non-compact case, define the support of a function  $f$  to be the set  $S_f$  that is the closure of the set  $\{p \in M : f(p) \neq 0\}$ . Since any compact set of  $M$  is contained in a fundamental set (a non-trivial remark), if  $M$  is oriented and Riemannian,  $f$  is  $C^\infty$  with compact support, and  $S_f \subset$  fundamental set  $A$ , then  $\int_M f = \int_A f$  is well-defined (independent of  $A$ ).

The area, volume, or measure (depending on the appropriate dimension) of a fundamental set  $A$  is the number  $\int_A f$ , where  $f \equiv 1$  on  $M$ . For a deeper study of integration theory on manifolds see the book of [Whi16].

## Problems

Let  $M$  be a  $C^\infty$   $n$ -manifold and let  $U$  be an open subset of  $M$ .

64. If  $X$  and  $Y$  are in  $\mathbf{T}^{1,0}M$ ,  $f, g \in C^\infty(M, \mathbb{R})$  and  $w \in \mathbf{T}^{0,1}M$  show

(i)  $\mathcal{L}_{fX}w = w(X)\mathbf{d}f + f(\mathcal{L}_Xw),$

(ii)  $\mathcal{L}_{fX}Y = f(\mathcal{L}_X Y) - \mathbf{d}f(Y)X,$

(iii)  $\mathcal{L}_{fX}g = f\mathcal{L}_X g,$

(iv) and  $\Delta(fw) = f\Delta w + w \otimes \mathbf{d}f.$

Thus  $\mathcal{L}$  and  $\Delta$  are not tensors.

65. If  $X$  is a  $C^\infty$  vector field on  $U$ ,  $m \in U$ ,  $Z_1, \dots, Z_n$  a base of  $T_m M$  with dual base  $w_1, \dots, w_n$

(i) Show  $(\operatorname{div} X)_m = \sum_{i=1}^n w_i(\mathbf{D}_{Z_i} X).$

(ii) Show that the divergence of a  $C^\infty$  field on  $\mathbb{R}^3$  agrees with the advanced calculus definition.

66. Let  $A$  be in  $\mathbf{T}^{1,1}U$ , let  $Z_1, Z_2, \dots, Z_k$  be a  $C^\infty$  base field on  $U$  and let  $w_1, w_2, \dots, w_k$  be the dual base on  $U$ . Show

$$\mathbf{D}_X w_j = - \sum_k w_j(\mathbf{D}_X Z_k)w_k \text{ and } \sum_j [A(\mathbf{D}_X w_j, Z_j) + A(w_j, \mathbf{D}_X Z_j)] = 0.$$

67. Let  $M$  be Riemannian, let  $X_1, \dots, X_{n-1}, T$  be an orthonormal base, and let  $P_i$  be the plane section spanned by  $X_i$  and  $T$ . Show

$$\operatorname{Ric}(T, T) = \sum_{i=1}^{n-1} K(P_i).$$

68. Prove formulas (5), (7), (13), and (14).

69. If  $\mathbf{D}$  has zero torsion, show

$$\mathbf{d}w(X, Y) = (\mathbf{D}_X w)(Y) - (\mathbf{D}_Y w)(X).$$

70. If  $M$  is Riemannian and  $G(X, Y) = \langle X, Y \rangle,$

(i) show that a connexion  $\mathbf{D}$  is metric preserving iff  $\Delta G = 0.$

(ii) Given arbitrary  $A \in \mathbf{T}^{0,3}M$  and  $B \in \mathbf{T}^{1,2}M$  with

$$A(X, Y, Z) = A(Y, X, Z) \text{ and } B(w, X, Y) = -B(w, Y, X)$$

for all  $w, X, Y, Z$ , show there exists a unique connexion  $\mathbf{D}$  on  $M$  with  $\Delta G = A$  and  $B(w, X, Y) = w(\text{Tor}_{\mathbf{D}}(X, Y))$ .

**71. (Poincaré lemma)** Show every closed  $p$ -form on  $\mathbb{R}^n$  is exact for  $p > 0$  as follows: for  $b$  in  $\mathbb{R}$  let  $g_b : \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$  by  $g_b(t_1, \dots, t_n) = (t_1, \dots, t_n, b)$ , let  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  by

$$f(t_1, \dots, t_{n+1}) = (t_{n+1}t_1, t_{n+1}t_2, \dots, t_{n+1}t_n),$$

let  $T = \frac{\partial}{\partial u_n}$ , and for  $p > 0$ , define the linear map  $K : \Omega^p(\mathbb{R}^n) \rightarrow \Omega^{p-1}(\mathbb{R}^n)$  by

$$K(w) = \int_0^1 (g_b)^* \circ C_T \circ f^*(w) \mathbf{d}b,$$

and show  $\mathbf{d}K + K\mathbf{d}$  equals the identity map on  $\Omega^p(\mathbb{R}^n)$ .

**72.** Let  $M$  be an oriented Riemannian 2-manifold. If  $\sigma$  is an oriented  $C^\infty$  curve in  $M$  with unit tangent  $T$ , let  $T, N$  be an orthonormal oriented base along  $\sigma$  and define the signed geodesic curvature of  $\sigma$  to be the  $C^\infty$  function  $b$  with  $\mathbf{D}_T T = bN$  on  $\sigma$ .

(i) If  $Z, W$  is an oriented orthonormal parallel base field along  $\sigma$  and  $T = (\cos \theta)Z + (\sin \theta)W$ , show  $b = \frac{\mathbf{d}\theta}{\mathbf{d}s} = T\theta$  on  $\sigma$ .

(ii) If  $x, y$  is an oriented orthogonal coordinate system on  $U \in M$ , let  $E = \langle X, X \rangle$  and  $G = \langle Y, Y \rangle$ . If  $b_1$  and  $b_2$  denote the geodesic curvature along the  $x$ -coordinate and  $y$ -coordinate curves, respectively, show

$$b_1 = -\frac{1}{2E\sqrt{G}} \frac{\partial E}{\partial y}, b_2 = \frac{1}{2E\sqrt{G}} \frac{\partial E}{\partial y}$$

and

$$K = \frac{1}{\sqrt{EG}} \left[ \frac{\partial b_1 \sqrt{E}}{\partial y} - \frac{\partial b_2 \sqrt{G}}{\partial x} \right]$$

(iii) Show the  $y$ -curve are geodesics (with  $y$  as parameter) iff  $G$  is constant.

73. If  $M$  is Riemannian,  $(\phi, U)$  is a coordinate pair,

$$x_i = u_i \circ \phi, g_{ij} = \left\langle \frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j} \right\rangle,$$

$g = \det(g_{ij})$ ,  $f$  is in  $C^\infty(M, \mathbb{R})$ , and  $A$  is a fundamental set with  $A \subset U$ , show

$$\int_A f = \int_{\phi(A)} (f \circ \phi^{-1}) \sqrt{g \circ \phi^{-1}} \mathbf{d}u_1 \mathbf{d}u_2 \dots \mathbf{d}u_n.$$

74. Let  $M$  be a surface in  $\mathbb{R}^3$  with sphere map  $\eta$ . For  $m$  in  $M$  let  $A(r)$  be the area of  $B(m, r)$ , the ball about  $m$  of radius  $r$  and let  $A_\eta(r)$  be the area of  $\eta(B(m, r))$ . Show

$$K(m) = \lim_{r \rightarrow 0} \left[ \frac{A_\eta(r)}{A(r)} \right].$$

## 8. Gauss-Bonnet Theory and Rigidity

In this chapter,  $M$  will denote a connected orientated Riemann  $n$ -manifold.

### 8.1 Gauss-Bonnet Formula

In this section, let  $n = 2$ , let  $A$  be a fundamental set in  $M$ , and let  $c$  be a fundamental 2-chain with  $|c| = A$ . The oriented curve  $\gamma = \partial c$  is called the *bounding curve* of  $A$ . A *vertex* of  $c$  is a point in  $M$  that is the image of a vertex in  $I^2$  under a 2-cube in  $c$ . A *face* of  $c$  is the support of a 2-cube in  $c$ . An *edge* of  $c$  is the face of a 1-cube in  $\partial\sigma$  for some 2-cube  $\sigma$  in  $c$ . A *boundary edge* of  $c$  is an edge that is in  $\gamma$ . A *corner point* of  $\gamma$  is a vertex of  $c$  belonging to exactly two boundary edges. At a corner point  $p$  of  $\gamma$ , let  $T_i(p)$  (the “tangent in”) and  $T_o(p)$  (the “tangent out”) be the unit tangents at  $p$  of the 1-cubes in  $\gamma$ , defined by the orientation, going “into” and “out from”  $p$ , respectively. The *exterior corner angle*  $\alpha(p)$  is the angle such that  $\cos \alpha(p) = \langle T_i(p), T_o(p) \rangle$  and  $0 < \alpha < \pi$  or  $-\pi < \alpha < 0$  according as  $T_i, T_o$  is a positively or negatively ordered base. If  $T_o = T_i$ , then  $\alpha = 0$ , and if  $T_o = -T_i$  then  $\alpha = -\pi$  (see 8.1).

In the proof of the Gauss-Bonnet formula that follows, the differential geometry involved is simple. The crux of the theorem is the Hopf Umlaufsatz (see discussion after proof). As usual, a *simple closed curve* is a homeomorphic image of the circle  $S^1$  in  $R^2$ .




Figure 8.1: Corner Angles

**Theorem 8.1 (Gauss-Bonnet formula).** Let  $A$  be contained in a coordinate domain  $U$  of  $M$ , let the bounding curve  $\gamma$  of  $A$  be a simple closed curve, and let  $\alpha_1, \dots, \alpha_r$  be the exterior corner angles of  $\gamma$ . Then

$$\int_{\gamma} k = 2\pi - \sum_{j=1}^r \alpha_j - \int_A K \quad (1)$$

where  $k$  is the signed geodesic curvature function on  $\gamma$  and  $K$  is the Riemannian (Gaussian) curvature function on  $A$ .

**Proof.** Let  $e_1, e_2$  be a  $C^\infty$  positively oriented base field on  $U$ . Let  $\gamma_1, \dots, \gamma_r$  be the  $C^\infty$  pieces of  $\gamma$  with each  $\gamma_j$  parameterized by arc length on the interval  $[s_j, s_{j+1}]$ ,  $\gamma_j(s_{j+1}) = \gamma_{j+1}(s_{j+1})$  for  $j = 1, \dots, r-1$ , while  $\gamma_r(s_{r+1}) = \gamma_1(s_1)$ , and  $\alpha_j$  the exterior corner angle at  $\gamma(s_j)$ . Let  $T$  be the unit tangent to  $\gamma$ . By making a constant rotation of  $e_1, e_2$ , if necessary, we may assume  $T(s_1^+) = e_1$ . Define  $\zeta(s)$  on  $[s_1, s_2]$  so  $\zeta$  is  $C^\infty$ ,  $\zeta(s_1^+) = 0$ , and  $T = (\cos \zeta)e_1 + (\sin \zeta)e_2$ . This  $\zeta$  is well-defined, since we have given its initial value and it is  $C^\infty$ , since locally it is given by  $\zeta(s) = \cos^{-1} \langle T(s), e_1(s) \rangle$  for a proper branch of the inverse cosine. Thus we obtain  $\zeta(s_2^-)$ . Let  $\zeta(s_2^+) = \zeta(s_2^-) + \alpha_1$  and extend  $\zeta$  to  $[s_2, s_3]$  so  $\zeta$  is  $C^\infty$  and  $T = (\cos \zeta)e_1 + (\sin \zeta)e_2$ , as before. Continuing this process, we extend  $\zeta$  to  $[s_1, s_{r+1}]$  with  $\zeta$  in  $C^\infty$  at all interior points except  $s_i$  where it has a jump precisely equal to  $\alpha_i$  for  $i = 2, \dots, r$ . Since  $\gamma$  is a simple closed curve, we use the Hopf Umlaufsatz to



obtain  $\zeta(s_{r+1}^-) + \alpha_1 = \zeta(s_1^+) + 2\pi$ . We include a schematic diagram (8.2):




Figure 8.2: Fundamental Set

On each  $C^\infty$  piece of  $\gamma$  we have the positively ordered orthonormal base field,  $T, N$ , and the signed geodesic curvature  $k$  is defined by  $D_T T = kN$ . In terms of  $\zeta$ ,

$$T = (\cos \zeta)e_1 + (\sin \zeta)e_2, \text{ while } N = (-\sin \zeta)e_1 + (\cos \zeta)e_2.$$

Let  $w_1, w_2$  be the dual 1-forms to the base  $e_1, e_2$  and let  $w_{12} = -w_{21}$  be the corresponding connection 1-form on  $U$  (note  $w_{11} = w_{22} = 0$  for the Riemann connection  $D$ ). Thus  $v = w_1 \wedge w_2$  is the volume element on  $U$ . Moreover, by the Cartan structural equations,  $dw_{12} = R_{12}$ , and

$$K = \langle R(e_1, e_2)e_2, e_1 \rangle = \left\langle \sum_{i=1}^2 R_{i2}(e_1, e_2)e_i, e_1 \right\rangle = R_{12}(e_1, e_2)$$

thus  $R_{12} = Kw_1 \wedge w_2$ .

Since  $k = \langle \mathbf{D}, N \rangle$  and

$$\mathbf{D}_T T = (T\zeta)N + (\cos \zeta)w_{21}(T)e_2 + (\sin \zeta)w_{12}(T)e_1,$$

then

$$k = (T\zeta) - w_{12}T, \tag{2}$$

which is a Cartan formula for the geodesic curvature. Then

$$\begin{aligned} \int_{\gamma} k &= \sum_{j=1}^r \int_{s_j}^{s_{j+1}} \frac{d\zeta}{ds} ds - \int_{\partial c} w_{12} = \sum_{j=1}^r [\zeta(s_{j+1}^-) - \zeta(s_j^+)] - \int_c dw_{12} \\ &= 2\pi - \sum_{j=1}^r \alpha_j - \int_A K, \end{aligned}$$

where we use Stokes' theorem for the second equality. □

The Gauss-Bonnet formula almost proves the Hopf Umlaufsatz (see [Hop26]), which states if  $\gamma$  is a simple closed smooth ( $C^1$ ) curve in  $\mathbb{R}^2$ , then  $\int_{\gamma} k = \pm 2\pi$ , depending on the orientation of  $\gamma$ . We need the topological result that  $\gamma$  disconnects the plane into two components and the map  $\gamma$  may be extended into a homeomorphism of the interior of the disc  $B(0,1)$ , which then maps onto a set  $A$ , which is fundamental and has  $\gamma$  as bounding curve. Then letting  $e_1 = i, e_2 = j$  (advanced calculus notation), we have  $w_{12} = 0, K = 0$ , and all  $\alpha_i = 0$ , so  $\int_{\gamma} k = 2\pi$  if  $\gamma$  positively oriented. The reader may also be interested in the papers of [Whi37], [Gri58], and [Tit].

The Gauss-Bonnet formula was first proved by Bonnet in 1848. Somewhat earlier Gauss had proved the following result on geodesic triangles.

**Theorem 8.2 (Gauss).** Let  $A$  be a fundamental set of  $M$  bounded by three (non-closed) geodesics, i.e.,  $A$  is a geodesic triangle, and

let  $\beta_1, \beta_2, \beta_3$  be the interior angles at the corners. Then

$$\int_A K = \beta_1 + \beta_2 + \beta_3 - \pi,$$

and this number is called the excess of the triangle.

*Proof.* The Gauss-Bonnet formula is applicable. Since  $k = 0$  and  $\alpha_i = \pi - \beta_i$ , we have

$$0 = 2\pi - \sum_{j=1}^3 (\pi - \beta_j) - \int_A K.$$

□

**Corollary 8.3.** Let  $B$  be the sum of the interior angles of a geodesic triangle  $A$  on  $M$ . Then  $B$  is  $> \pi$ ,  $= \pi$ , or  $< \pi$ , according as  $K > 0$ ,  $= 0$ , or  $< 0$  on  $A$ . If  $K$  is constant and not zero on  $A$ , then the area of  $A$  equals the excess of  $A$  divided by  $K$ .

We obtain some simple applications of the Gauss-Bonnet formula by applying it to the cases when  $M$  is diffeo to the sphere or the torus. In the former case  $\int_M K = 4\pi$ , and in the latter case  $\int_M K = 0$ . These are special cases of the Gauss-Bonnet theorem which we prove later in this section. We sketch the proofs of these facts.

When  $M$  is diffeo to  $S^2$ , we let  $\gamma$  be the image of the equator (under the diffeo),  $A_1$  the image of the “northern” hemisphere, and  $A_2$  the image of the “southern” hemisphere (see 8.3). Supposing  $\gamma$  to be the bounding curve of  $A_1$ , we have

$$\int_{\gamma} k = 2\pi - \int_{A_1} K \quad \text{and} \quad \int_{-\gamma} k = - \int_{\gamma} k = 2\pi - \int_{A_2} K.$$

Hence

$$\int_M K = \int_{A_1} K + \int_{A_2} K = 4\pi.$$




Figure 8.3: Sphere and Torus

When  $M$  is diffeo to the torus, let  $A_1$  be the image of the “top half” and  $A_2$  the image of the “bottom half” of the torus so  $A_1$  and  $A_2$  are bounded and separated by the image  $\gamma$  of the “inside” and “outside” curve on the torus (see 8.3). Again letting  $\gamma$  be the bounding curve of  $A_1$ , connecting and closing  $\gamma$  via a cut curve  $\beta$  (see 8.3), and taking a limit, we obtain

$$\int_{\gamma} k = 2\pi - 2\pi - \int_{A_1} K \text{ and } -\int_{\gamma} k = -\int_{A_2} K \text{ so } \int_M K = 0.$$

Our next task is to free the Gauss-Bonnet formula from the special neighborhood  $U$ . The proof follows from the [Sam55]. Define the Euler characteristic,  $\chi_c(A)$ , of  $A$  with respect to  $c$  by  $\chi_c(A) = V - E + F$ , where  $V$  is the number of vertices of  $c$ ,  $E$  the number of edges, and  $F$  the number of faces.

**Theorem 8.4.** Let  $A$  be a fundamental set on  $M$ , let the bounding curve  $\gamma$  of  $A$  be a finite disjoint union of simple closed curves, and let  $\alpha_1, \dots, \alpha_r$  be the exterior corner angles of  $\gamma$ . Then

$$\int_{\gamma} k = 2\pi\chi_c(A) - \sum_{i=1}^r \alpha_i - \int_A K. \tag{3}$$

This expression proves  $\chi_c(A)$  is independent of  $c$ , so define  $\chi(A) = \chi_c(A)$  to be the Euler characteristic of  $A$  and drop the subscript  $c$

in the above formula.

*Proof.* Let  $c = \sigma_1 + \dots + \sigma_F$  and note from the definition of a fundamental 2-chain we may apply the Gauss-Bonnet formula to each set  $|\sigma_j|$  (for  $\sigma_j$  defines a coordinate neighborhood of  $|\sigma_j|$ ). Let  $\alpha_1^j, \dots, \alpha_4^j$  denote the four exterior angles for  $\sigma_j$ . Then

$$\int_{\gamma} k = \sum_{j=1}^F \int_{\partial\sigma_j} k = \sum_{j=1}^F (2\pi - \sum_{i=1}^4 \alpha_i^j) - \sum_{j=1}^F \int_{\sigma_j} K$$

or

$$\int_{\gamma} k = 2\pi F - \sum_{j=1}^F \sum_{i=1}^4 \alpha_i^j - \int_A K.$$

Thus the problem is one of bookkeeping with the term  $\sum \alpha_i^j$ .

Let  $\beta_i^j$  be the interior angle corresponding to each  $\alpha_i^j$ , thus  $\beta_i^j = \pi - \alpha_i^j$ , and let  $\beta_s = \pi - \alpha_s$  be the interior angles at the corners of  $\gamma$ . In the following we sum over  $i = 1, \dots, 4$  and  $j = 1, \dots, F$ . The sum

$$\begin{aligned} \sum_{ij} \beta_i^j &= 2\pi(V - r) + \sum_1^r \beta_s = 2\pi(V - r) + \sum_1^r (\pi - \alpha_s) \\ &= 2\pi V - \pi r - \sum_1^r \alpha_s, \end{aligned}$$

since  $r$  is the number of vertices of  $c$  on  $\gamma$  (as well as the number of angles and edges on  $\gamma$ ) so  $(V - r)$  is the number of vertices interior to  $A$ , each of which contributes  $2\pi$  to the total sum.

We now show that  $rF = (2E - r)$ , which is the number of terms in the sum  $\sum_{ij} \beta_i^j$ . This is done by assigning to each  $\beta_i^j$  an edge, namely, its "starting" edge, which is well-defined by the orientation. More precisely, if  $T$  and  $T'$  are the unit vectors at the vertex of  $\beta_i^j$  which are tangent to the edge curves of  $\beta_i^j$ , then  $T$  and  $T'$  are independent, since  $c$  is a fundamental chain, so  $(\sigma_j)_*$  is non-singular on its domain (which is slightly larger than  $I^2$ ). Thus  $T$  is the "starting" edge of  $\beta_i^j$  if and only if  $T, T'$  is a positively oriented

bases (see 8.4).




Figure 8.4: Starting Edges

Then each edge on the boundary  $\gamma$  belongs to exactly one  $\beta_i^j$ , while each edge not on the boundary belongs to exactly two  $\beta_i^j$ . Thus  $rF = r + 2(E - r)$ , since  $r$  is the number of edges on the boundary.

Finally,

$$\sum_{ij} \alpha_i^j = \sum_{ij} (\pi - \beta_i^j) = \pi(2E - r) - 2\pi V + \pi r + \sum_i \alpha_s = 2\pi(E - V) + \sum_1^r \alpha_s.$$

Hence,

$$\int_{\gamma} k = 2\pi(F - E + V) - \sum_1^r \alpha_s - \int_A K.$$

□

**Theorem 8.5 (Gauss-Bonnet Theorem).** Let  $M$  be a compact connected oriented Riemannian 2-manifold with Riemannian (Gaussian) curvature function  $K$ . Then

$$\int_M K = 2\pi\chi(M).$$

*Proof.* We apply the preceding theorem to a fundamental chain on

$M$  which will have no boundary and no exterior angles. □

The above theorem is an important example of a theorem relating differential geometry and topology. The Euler characteristic is a topological invariant which does not depend on either the differentiable structure or the Riemannian structure on  $M$ . The theorem may be used to prove many “negative” statements: for example, there does not exist a Riemannian metric on the torus with  $K > 0$  everywhere (nor does there exist one with  $K < 0$  everywhere) since  $\chi(M) = 0$  (which we computed above for the induced Riemannian metric). The theorem has been generalized for dimensions greater than two and provides one of the first successes of the global theory of fiber bundles.

## 8.2 Index Theorem

This section is also based on [Sam55]. Let  $n = 2$  and let  $W$  be a  $C^\infty$  vector field on  $M$ . If  $W_m = 0$ , then  $m$  is a singularity of  $W$ . Assuming  $W$  has only isolated singularities, we define the index of  $W$  at  $m$ ,  $J(W, m)$  as follows.

Let  $U$  be a coordinate domain, with coordinate radius  $b > 0$ , about  $m$  with  $W \neq 0$  on  $U - \{m\}$ . Assume the coordinate map is orientation preserving, and let  $\sigma_r$  be the oriented coordinate circle of radius  $r$  about  $m$  with  $0 < r < b$  and  $\sigma_r$  defined on  $[0, 1]$ . Let  $X$  be a unit vector field on  $U$ . Since  $W$  does not vanish on  $\sigma_r$ , by using the proper inverse cosine function one obtains a  $C^\infty$  function  $\theta$  on  $[0, 1]$  with  $\langle W(s), X(s) \rangle = |W(s)| \cos \theta(s)$  on  $[0, 1]$ . Let

$$2\pi J_X(W, m, r) = \theta(1) - \theta(0) \tag{4}$$

For  $0 < r < b$ ,  $J_X(W, m, r)$  is a continuous integer-valued function, and hence yields a constant  $J_X(W, m)$ . If  $m$  is not a singular point, then for small  $r > 0$ ,  $\theta$  is close to the constant  $\cos^{-1}(\frac{\langle W_m, X_m \rangle}{|W_m|}) \pmod{2\pi}$ ; hence  $\theta(1) = \theta(0)$ , and  $J_X(W, m) = 0$ . If  $Y$  is another unit vector field

on  $U$ , then

$$J_X(W, m) = J_Y(W, m) + J_X(Y, m) = J_Y(W, m),$$

since  $Y$  has no singularities. Thus  $J_X(W, m)$  is independent of  $X$ . An analogous argument shows  $J(W, m)$  can be computed by using any simple closed  $C^\infty$  curve  $\sigma$  about  $m$  with  $\sigma$  in  $U$ , and thus  $J(W, m)$  is an integer depending only on  $W$  and  $m$  (see 8.5).




Figure 8.5: Examples of  $J(W, m)$

If  $W$  has only a finite number of singularities, define the index of  $W$ ,  $J(W)$ , by  $J(W) = \sum_m J(W, m)$ .

**Theorem 8.6 (Index Theorem).** If  $M$  is a compact connected oriented Riemannian 2-manifold and  $W$  is a  $C^\infty$  vector field on  $M$  with a finite number of singularities, then the index of  $W$  equals the Euler characteristic of  $M$ .

*Proof.* Take an oriented fundamental chain  $c = \sigma_1 + \dots + \sigma_r$  with at most one singularity  $m_i$  of  $W$  in the interior of each  $|\sigma_i|$ . Let  $\gamma_i$  be the bounding curve of  $\sigma_i$ , and define functions  $\theta_i, \zeta_i, \xi_i$  on the domain of  $\gamma_i$  so that

- $\theta = \zeta_i + \xi_i$ ,
- $\theta_i$  is an angle between  $W$  and  $e_i$ ,
- $\zeta_i$  is an angle between the tangent  $T_i$  of  $\gamma_i$
- and  $\xi_i$  will be piece-wise  $C^\infty$  and  $\theta_i$  is continuous.



By integrating over the pieces of  $\gamma_i$  we obtain

$$\begin{aligned} 2\pi J(W, m_i) &= \int_{\gamma_i} \frac{d\theta_i}{ds} = \int_{\gamma_i} \frac{d\zeta_i}{ds} + \int_{\gamma_i} \frac{d\xi}{ds} \\ &= \int_{\gamma_i} k + \int_{\sigma_i} K + \int_{\gamma_i} \frac{d\xi_i}{ds} \end{aligned}$$

Adding over the 2-cubes in  $c$  gives

$$2\pi J(W) = \int_M K \tag{5}$$

since the integrals over the bounding curves cancel one another. By the Gauss-Bonnet theorem,  $J(W) = \chi(M)$ .  $\square$

Omitting the last line of the proof, we note  $2\pi J(W) = \int_M K$  implies  $J(W)$  is independent of  $W$  as long as  $W$  has only a finite number of singularities. Then for any oriented fundamental chain  $c$  we can define a particular  $W$  which has a singularity for each face, edge, and vertex with index 1,  $-1$ , and 1 respectively. We indicate in 8.6 how  $W$  is defined on each 2-cube. Actually  $W$  would be precisely defined by defining a field on a neighborhood of  $I^2$  and carrying this to each  $|\sigma_i|$  via the map  $\sigma_i$ .




Figure 8.6: The Canonical Vector Field on a 3-cube

Thus  $W$  is defined by “going out from each vertex and in to the

center of each face.” From 8.6 we see  $J(W) = V - E + F = \chi_c(M)$ . Thus we again prove  $\chi_c(M)$  is independent of  $c$ , and  $2\pi\chi(M) = 2\pi J(W) = \int_M K$  reproves the Gauss-Bonnet theorem.

**Corollary 8.7.** If  $M$  is a manifold as described in the theorem and there exists a non-vanishing  $C^\infty$  vector field on  $M$ , then  $\chi(M) = 0$ . Thus any surface that is diffeomorphic to the 2-sphere has no non-vanishing  $C^\infty$  vector fields.

Actually, a differentiable manifold (any dimension) admits a nonzero continuous vector field if and only if its Euler characteristic is zero (see [Ste51, p. 203], and [AH37, p. 549]).

### 8.3 Gauss-Bonnet Form

In the proof of the Gauss-Bonnet formula we found that  $R_{12}$  is a local representation of the global form  $Kv$  on an oriented Riemannian 2-manifold  $M$ . One might ask if there are other global forms obtainable in this way, or if there is an analogous form on an  $n$ -manifold. We answer these questions now.

Let  $e_1, \dots, e_n$  and  $f_1, \dots, f_n$  be two sets of positively oriented orthonormal  $C^\infty$  base fields on an open set  $U$  in  $M$ , and let  $f_j = \sum_{i=1}^n b_{ij}e_i$  define  $C^\infty$  functions  $b_{ij}$  on  $U$ . Notice that determinant  $(b_{ij}) = 1$  and  $(b_{ij})^{-1} = (b_{ji})$  since  $(b_{ij})$  is orthogonal. We let  $R_{ij}$  and  $\bar{R}_{ij}$  denote the local curvature forms associated with  $e_i$ 's and  $f_i$ 's, respectively, thus  $R(X, Y)e_j = \sum R_{ij}(X, Y)e_i$ . Then for  $m \in U$ ,  $X$  and  $Y$  in  $\mathbf{T}_mM$ , we have

$$\begin{aligned} R_{ij}(X, Y) &= \langle R(X, Y)e_j, e_i \rangle = \left\langle R(X, Y) \left( \sum_r b_{jr}f_r \right), \sum_s b_{is}f_s \right\rangle \\ &= \sum_{r,s} b_{jr}b_{is} \langle R(X, Y)f_r, f_s \rangle = \sum_{r,s} b_{is}\bar{R}_{sr}(X, Y)b_{jr}. \end{aligned} \tag{6}$$

Thus  $R_{ij} = \sum_{r,s} b_{is}\bar{R}_{sr}b_{jr}$  relates the local curvature forms of the two bases on  $U$ .

If  $n$  is even, we define an  $n$ -form  $Q$  on  $U$  by

$$Q = \sum (-1)^\pi R_{\pi(1)\pi(2)} \wedge R_{\pi(3)\pi(4)} \wedge \cdots \wedge R_{\pi(n-1)\pi(n)} \quad (7)$$

where we sum over all permutations  $\pi$  in  $P_n$ , the group of permutations on the set  $\{1, 2, \dots, n\}$ . The representation of  $Q$  in terms of the forms  $\bar{R}_{ij}$  is,

$$\begin{aligned} Q &= \sum_{r_i=1}^n \sum (-1)^\pi b_{\pi(1)r_1} \bar{R}_{r_1 r_2} b_{\pi(2)r_2} b_{\pi(3)r_3} \bar{R}_{r_3 r_4} b_{\pi(4)r_4} \cdots \\ &= (\det b_{ij}) \sum (-1)^\pi \bar{R}_{\pi(1)\pi(2)} \wedge \bar{R}_{\pi(3)\pi(4)} \wedge \cdots \wedge \bar{R}_{\pi(n-1)\pi(n)} \end{aligned}$$

Since  $(\det b_{ij}) = 1$ ,  $Q$  is independent of the particular base field used to define it; thus  $Q$  defines a global  $n$ -form on  $M$  which is called the Gauss-Bonnet form. Note if  $n = 2$ , then locally  $Q = R_{12} - R_{21} = 2R_{12} = 2Kv$ .

**Theorem 8.8 (Generalized Gauss-Bonnet).** If  $M$  is an even dimensional ( $n = 2k$ ) compact connected oriented Riemannian manifold, then

$$\int_M Q = 2^n \pi^k (k!) \chi(M).$$

For a proof see [Che51]. Other pertinent references are [Hop26], [All40], [AW43], [Fen40], [Che45], and [All50].

Let  $M$  be as in the theorem and assume further that  $M$  is a hypersurface in  $\mathbb{R}^{n+1}$  with unit normal field  $N$ . Using the notation from section 4,

$$R_{ij} = -w_{i,n+1} \wedge w_{n+1,j} = w_{i,n+1} \wedge w_{j,n+1}$$

and

$$L(X) = \sum_{i=1}^n w_{i,n+1}(X) e_i = \eta_*(X)$$

where  $\eta$  is the sphere map induced by the normal  $N$  (section 2.2).

Thus  $\eta^*W_i = W_{i,n+1}$  and

$$\begin{aligned} Q &= \sum (-1)^\pi w_{\pi(1),n+1} \wedge \cdots \wedge w_{\pi(n-1),n+1} \wedge w_{\pi(n),n+1} \\ &= n! w_{1,n+1} \wedge w_{2,n+1} \wedge \cdots \wedge w_{n,n+1} = n! \eta^*(v_S), \end{aligned}$$

where  $v_S$  is the volume element of the unit sphere  $S^n$  oriented by its outer normal, and we assume  $N_m$  is parallel to the outer normal at  $\eta(m)$ . Integrating,

$$\int_M Q = (n!) \int_M \eta^*(v_S).$$

If  $n = 2$ , then

$$\int_M Q = 2 \int_M K v = 4\pi \chi(M) = 2 \int_M \eta^*(v_S),$$

thus

$$\int_M \eta^*(v_S) = \frac{(V_2)\chi(M)}{2}$$

where  $V_2$  is the “volume” of the unit 2-sphere. This is the *Hopf index theorem* for dimension 2.

In the general case ( $M^n$  imbedded in  $\mathbb{R}^{n+1}$  as above), we let  $X$  be any unit vector field of  $\mathbb{R}^{n+1}$  that is  $C^\infty$  on  $M$ , and we define the *index* of  $X$  on  $M$ ,  $\mathcal{I}(X)$ , by

$$\mathcal{I}(X) = \frac{1}{V_n} \int_M \eta_X^*(v_S), \tag{8}$$

where  $V_n$  is the “volume” of the unit  $n$ -sphere in  $\mathbb{R}^{n+1}$ , and  $\eta_X$  is the  $C^\infty$  map of  $M$  into  $S^n$  induced by the vector field  $X$ .

**Theorem 8.9 (Hopf index theorem).** If  $M^n$  is an even-dimensional compact connected submanifold of  $\mathbb{R}^{n+1}$ , then twice the index of the normal field  $N$  on  $M$  is the Euler characteristic of  $M$ , or  $2\mathcal{I}(N) = \chi(M)$ .

*Proof.* Assuming the [Gauss-Bonnet theorem](#) and letting  $n = 2k$ , we

see

$$2\mathcal{I}(N) = 2 \int_M \frac{Q}{V_n n!} = 2^{n+1} \pi^k k! \frac{\chi(M)}{V_n n!} = \chi(M),$$

since  $V_n = \frac{2^{n+1} \pi^k k!}{n!}$  (see problem 75). □

## 8.4 Characteristic Forms

A general reference for this section is [Che51] with related treatments in [Adl57] and [Car25]. The “wedge” product symbol between forms will be omitted in this section.

For  $k > 1$ , define local forms

$$Q_k = \sum_{i_j=1}^n R_{i_1 i_2} R_{i_2 i_3} R_{i_3 i_4} \cdots R_{i_k i_1},$$

where the  $R_{i_j}$  belong to a local positively oriented orthonormal base field  $e_1, \dots, e_n$ . As above (equation 7), one shows  $Q_k$  is independent of this particular base field and thus  $Q_k$  is a global  $2k$ -form on  $M$ . Moreover  $dQ_k = 0$ , i.e., each  $Q_k$  is a closed  $2k$ -form. To prove this, use

$$dR_{ij} = \sum_{t=1}^n (R_{it} w_{rj} - R_{jr} w_{ti})$$

which follows from the second structural equation (section 5.2). Then,

$$dQ_k = \sum [ (dR_{i_1 i_2}) R_{i_2 i_3} \cdots R_{i_k i_1} + R_{i_1 i_2} (dR_{i_2 i_3}) R_{i_3 i_4} \cdots R_{i_k i_1} + \cdots ].$$

Consider one of the sums (all indices are summed from 1 to  $n$ ),

$$A = \sum R_{i_1 r} w_{r i_2} R_{i_2 i_3} \cdots R_{i_k i_1}.$$

If  $k$  is even, the products in  $A$  are formed from an odd number of forms that are skew-symmetric in their indices; hence switching all the indices changes the sign, and adding, one gets  $A = -A$  so  $A = 0$ .

If  $k$  is odd, the argument just used shows  $Q_k = 0$ .

**Proposition 8.10.** For even  $k$ , the forms  $Q_k$  define global closed  $2k$ -forms on  $M$ . For odd  $k$ ,  $Q_k = 0$ .

Let  $W_D$  denote the subalgebra of the Cartan differential algebra  $F$  (or  $F(M)$ ) which is generated over the real field by the forms  $Q_k$  for  $k = 2, 4, \lfloor n/2 \rfloor$ , and call  $W_D$  the algebra of characteristic forms for the connexion  $D$ . Elements in  $W_D$  are called characteristic forms, and they are closed forms since the generators are all closed. By going to the differential cohomology we can free ourselves of the connexion  $D$  which we now do.

Let  $\Omega^p$  denote the module of  $C^\infty$   $p$ -forms on  $M$ . Let  $Z^p$  denote the closed forms in  $\Omega^p$ , thus  $Z^p = \{\alpha \in \Omega^p : d\alpha = 0\}$ ; and let  $B^p$  denote the exact forms in  $\Omega^p$ , so

$$B^p = \{\alpha \in \Omega^p : \text{there is } \beta \in F^{p-1} \text{ with } d\beta = \alpha\}.$$

Since  $d^2 = 0, B^p \subset Z^p$ ; hence let  $H^p = Z^p/B^p$  and call  $H^p$  the  $p$ -dimensional differential cohomology group of  $M$ . If  $\alpha$  in  $Z^p$ , denote its image in  $H^p$  by  $\bar{\alpha}$ ; hence  $\bar{\alpha}$  is the coset  $\alpha + B^p$  which is called a (differential) cohomology class on  $M$ . Let  $H^* = \bigoplus_{p=0}^n H^p$  (direct sum) and notice the multiplication in  $F$  carries over to  $H^*$ .

Thus  $\bar{W}_D$  defines a set of classes called (differential) characteristic cohomology classes, and this set we show is independent of  $D$  (the Riemannian structure) and depends only on the manifold  $M$ . It is customary to speak of  $\bar{W}_D$  as the image of the *Wiel* homomorphism. This we explain.

Let  $\mathfrak{gl}(n, \mathbb{R})$  be the set of  $n$  by  $n$  matrices over the real field  $\mathbb{R}$ . Our notation is the customary one for this set when it is thought of as the Lie algebra of the general linear group  $\text{GL}(n, \mathbb{R})$ . If  $A = (a_{ij})$  in  $\mathfrak{gl}(n, \mathbb{R})$  we let  $u_{ij}(A) = a_{ij}$ . Then a polynomial function  $P$  on  $\mathfrak{gl}(n, \mathbb{R})$  is a polynomial in the functions  $u_{11}, u_{12}, \dots, u_{nn}$ ; for example,  $P(A) = \det(A)$  is a polynomial function. An invariant polynomial  $P$  on  $\mathfrak{gl}(n, \mathbb{R})$  is a polynomial function  $P$  such that  $P(BAB^{-1}) = P(A)$

for all non-singular orthogonal matrices  $B$ . Referring to the way we define the characteristic forms  $Q_k$ , we see that every invariant polynomial  $P$  can be used to define a global differential form  $Q$  on  $M$  by using the curvature forms from a Riemannian connexion  $D$  on  $M$  and letting  $Q = P(R_{11}, R_{12}, \dots, R_{nn})$ . Let us use  $\mathcal{W}_D$  for this map, so  $Q = \mathcal{W}_D(P)$ . Letting  $\mathcal{J}$  denote the set of invariant polynomials on  $\mathfrak{gl}(n, \mathbb{R})$ , we then claim to have a homomorphism  $\mathcal{W}_D : \mathcal{J} \rightarrow F(M)$  with  $\mathcal{W}_D(\mathcal{J}) = W_D(M) = W_D$ . This is the *Weil homomorphism*.

**Theorem 8.11.** The Weil homomorphism is well-defined from the set of invariant polynomials on  $\mathfrak{gl}(n, \mathbb{R})$  onto the set of characteristic differential forms on  $M$ ; moreover, the Weil homomorphism is independent of the connexion at the cohomology level, i.e.,  $\overline{W}_{D_1} = \overline{W}_{D_2}$  for two Riemannian connexions  $D_1$  and  $D_2$ .

*Proof.* Let  $f_A(\lambda)$  denote the characteristic polynomial of a matrix  $A$  and define polynomials  $E_r(A)$  to be the coefficients of  $f_A(\lambda)$ , thus

$$f_A(\lambda) = \det(\lambda I - A) = \lambda^n + E_{n-1}(A)\lambda^{n-1} + \dots + E_0(A).$$

From linear algebra we know  $E_r(A)$  are invariant polynomials on  $\mathfrak{gl}(n, \mathbb{R})$ ; moreover, they generate a ring of invariant polynomials. In terms of the characteristic roots of  $A$ ,  $E_r(A)$  is the  $r$ th elementary symmetric function of these roots, i.e.,  $E_1(A) = a_1 + \dots + a_n$ ,  $E_2(A) = \sum_{i < j} a_i a_j$ , etc. By Newton's theorem on symmetric functions, the functions  $E_r(A)$  are expressible as polynomials in the functions  $P_r(A)$ , where  $P_r(A) = (a_1)^r + (a_2)^r + \dots + (a_n)^r$ . But  $P_r(A)$  is the trace of  $A^r$ , and we can write this trace in terms of the elements of  $A$  by

$$P_r(A) = \sum a_{i_1 i_2} a_{i_2 i_3} \dots a_{i_r i_1}$$

summing over all  $i_j = 1, \dots, n$ . Hence  $\mathcal{W}_D(\mathcal{J})$  is generated by the forms  $Q_k$ ,  $\mathcal{W}_D$  is well-defined, and  $\mathcal{W}_D(\mathcal{J}) = W_D$ .

To show  $\overline{W}_D$  is independent of the Riemannian connexion, we take two such connexions  $D_1$  and  $D_0$ , let  $Q_k^i = W_{D_i}(P_k)$  for  $i = 0, 1$ ,

and show  $Q_k^1 - Q_k^0 = dG$ , where  $G$  in  $\Omega^{2k-1}(M)$ . Thus  $\bar{Q}_k^1 = \bar{Q}_k^0$ , which implies  $\bar{W}_{D_1} = \bar{W}_{D_0}$ .

Let  $\langle X, Y \rangle_1$  and  $\langle X, Y \rangle_0$  be the Riemannian metrics associated with  $D_1$  and  $D_0$ , respectively, and for  $0 \leq t \leq 1$  define  $\langle X, Y \rangle_t = t\langle X, Y \rangle_1 + (1-t)\langle X, Y \rangle_0$ . Then  $\langle X, Y \rangle_t$  is a Riemannian metric for each  $t$ , and its Riemannian connexion  $D_t$  is given by  $D_t = tD_1 + (1-t)D_0$ . This can be shown easily by verifying that  $D$  has zero torsion and preserves the metric  $\langle X, Y \rangle_t$ . For any base field  $e_1, \dots, e_n$  on an open set  $U$  of  $M$  let  $w_{ij}^t$  and  $R_{ij}^t$  be the connexion and curvature forms associated with  $D_t$ . Then

$$(D_t)_X e_j = \sum_i w_{ij}^t(X) e_i = t \sum_i w_{ij}^1(X) e_i + (1-t) \sum_i w_{ij}^0(X) e_i,$$

so  $w_{ij}^t = tw_{ij}^1 - (1-t)w_{ij}^0$ . From the second Cartan structural equation we obtain

$$R_{ij}^t = tR_{ij}^1 + (1-t)R_{ij}^0 + t(t-1) \sum_k \theta_{ik} \theta_{kj}$$

where  $\theta_{ij} = w_{ij}^1 - w_{ij}^0$ . The 1-forms  $\theta_{ij}$  are the local forms belonging to the difference tensor  $B(X, Y) = (D_1)_X Y - (D_0)_X Y$ , i.e.,  $B(X, e_j) = \sum_i \theta_{ij}(X) e_i$ . Since  $B$  is a tensor, if  $f_1, \dots, f_n$  is another base field on  $U$  with  $f_j = \sum_i b_{ij} e_i$  and

$$B(X, f_j) = \sum_i \bar{\theta}_{ij}(X) f_i, \text{ then } \theta_{ij}(X) = \sum_{r,s} b_{is} \theta_{sr}(X) (b^{-1})_{rj}.$$

For each even  $k$  and each  $t$ , choose  $e_1, \dots, e_n$  to be an orthonormal base field relative to the metric  $\langle X, Y \rangle_t$ , and define a  $(2k-1)$ -form on  $U$  by

$$G_k^t = \sum \theta_{i_1 i_2} R_{i_2 i_3}^t R_{i_3 i_4}^t \cdots R_{i_k i_1}^t,$$

summing over all  $i_j = 1, \dots, n$ . Since the  $\theta_{ij}$  transform exactly like the  $R_{ij}$  when changing to another orthonormal base, the forms  $G_k^t$  are global forms on  $M$  by the argument that was used to show  $Q_k$  are global forms. Note  $\theta_{ij}$  are not skew symmetric.



In an obvious way, define for each  $t$ , a  $2k$ -form  $\left(\frac{d}{dt}\right)Q_k^t$ , i.e.,

$$\frac{d}{dt}Q_k^t = \frac{d}{dt}\left(\sum R_{i_1 i_2}^t R_{i_2 i_3}^t \cdots R_{i_k i_1}^t\right) = k \sum \left(\frac{d}{dt}R_{i_1 i_2}^t\right) R_{i_2 i_3}^t \cdots R_{i_k i_1}^t$$

where

$$\frac{d}{dt}R_{ij}^t = R_{ij}^1 - R_{ij}^0 + (2t-1) \sum_k \theta_{ik} \theta_{kj}.$$

Then  $Q_k^1 - Q_k^0 = dG_k$  where  $G_k = k \int_0^1 G_k^t dt$ .

To compute  $dG_k^t$ , use the second Cartan structural equation to obtain

$$d\theta_{ij} = R_{ij}^1 - R_{ij}^0 - \sum [\theta_{ik} \theta_{kj} + \theta_{ij} w_{kj}^0 + w_{ik}^0 \theta_{kj}].$$

Also,

$$dR_{ij}^t = \sum (R_{ik}^t w_{kj}^t - w_{ik}^t R_{kj}^t) = \sum [tR_{ik}^t \theta_{kj} - t\theta_{ik} R_{kj}^t + R_{ik}^t w_{kj}^0 - w_{ik}^0 R_{kj}^t],$$

since  $w_{ij}^t = t\theta_{ij} + w_{ij}^0$ . Hence

$$\begin{aligned} dG_k^t &= \sum (R_{i_1 i_2}^1 - R_{i_1 i_2}^0 \theta_{i_1 k} \theta_{k i_2} - \theta_{i_1 k} w_{k i_2}^0 - w_{i_1 k} \theta_{k i_2}) R_{i_2 i_3}^t \cdots R_{i_k i_1}^t \\ &\quad - \left[ \sum \theta_{i_1 i_2} (tR_{i_2 k}^t \theta_{k i_3} - t\theta_{i_2 k} R_{k i_3}^t + R_{i_2 k}^t w_{k i_3}^0 - w_{i_2 k}^0 R_{k i_3}^t) R_{i_3 i_4}^t \cdots R_{i_k i_1}^t \right] - \cdots \\ &\quad - \left[ \sum \theta_{i_1 i_2} R_{i_2 i_3}^t \cdots R_{i_{k-1} i_k}^t (tR_{i_k j}^t \theta_{j i_1} - t\theta_{i_k j} R_{j i_1}^t + R_{i_k j}^t w_{j i_1}^0 - w_{i_k j}^0 R_{j i_1}^t) \right] \\ &= \sum \left[ R_{i_1 i_2}^1 - R_{i_1 i_2}^0 + (2t-1) \sum \theta_{i_1 k} \theta_{k i_2} \right] R_{i_2 i_3}^t \cdots R_{i_k i_1}^t = \frac{1}{k} \frac{d}{dt} Q_k^t. \end{aligned}$$

□

## 8.5 Rigidity Problems

Two submanifolds of  $\mathbb{R}^n$  are congruent or symmetric if there is an isometry of  $\mathbb{R}^n$  mapping one onto the other that is orientation preserving or reversing, respectively. Let us say a submanifold  $M$  of  $\mathbb{R}^n$  is *rigid* if any submanifold  $M'$  that is isometric to  $M$  is actually congruent or symmetric to  $M$ . Natural questions arise which

are called *rigidity problems*. For example, which submanifolds are rigid, or when are two isometric submanifolds congruent or symmetric?

Our principal reference for this section is [Che51]. The standard procedure in the following theorems is to somehow set up the hypothesis of the fundamental rigidity theorem proved in section 6.5. Given an isometry  $f$  between submanifolds, the first fundamental form is preserved by hypothesis, and our task is to show the second fundamental form is preserved, or that  $f_*$  commutes with the fundamental linear transformations  $L$ .

**Theorem 8.12.** If  $n \geq 3$  and  $M$  is an oriented hypersurface in  $\mathbb{R}^{n+1}$  with positive Riemannian curvature, then  $M$  is rigid.

*Proof.* Let  $f : M \rightarrow M'$  be an isometry and let  $L' = L \circ f_*$ . Since  $f$  is an isometry, the Gauss curvature equations give  $R(X, Y)Y = R'(X, Y)Y$  or

$$\langle LY, Y \rangle L(X) - \langle LY, X \rangle L(Y) = \langle L'Y, Y \rangle L'(X) - \langle L'Y, X \rangle L'(Y),$$

where  $X, Y$  in  $T_m M$ . Choose an orthonormal base  $X_1, \dots, X_n$  of vectors at  $m$ , and let  $LX_i = k_i X_i$ . We show  $L'$  is invariant on each subspace  $P_{ij}$  spanned by  $X_i$  and  $X_j$  for  $i \neq j$ . Let  $b_{rs} = \langle L'X_r, X_s \rangle$ , and the Gauss curvature equations imply

$$\begin{aligned} k_i k_j X_i &= b_{jj} L'X_i - b_{ij} L'X_j \\ k_i k_j X_j &= -b_{ij} L'X_i + b_{ii} L'X_j. \end{aligned}$$

Then  $K(P_{ij}) = k_i k_j = b_{ii} b_{jj} - b_{ij}^2 > 0$  implies  $L'X_i$  and  $L'X_j$  lie in  $P_{ij}$ . Since  $n \geq 3$ , there is a third index  $r$  with  $L'X_i$  in  $P_{ir}$ ; hence  $L'X_i$  lies in  $P_{ir} \cap P_{ij}$ , and thus  $X_i$  is an eigenvector of  $L'$ . For all  $i$ , let  $L'X_i = h_i X_i$ . Then  $k_i k_j = h_i h_j > 0$  for all  $i \neq j$ ; hence  $k_i^2 = h_i^2$ , so  $h_i = \pm k_i$ . The positive curvature condition also implies  $h_i = k_i$  for all  $i$ , or  $h_i = -k_i$  for all  $i$ . Thus  $L = \pm L'$  and we apply the fundamental rigidity theorem.  $\square$

If in the above theorem we assume  $M$  is complete (or closed),

then we need not assume it is oriented. For  $n = 2$ , the Cohn-Vossen theorem provides a similar result with the additional requirement that  $M$  be compact. We now examine some global functions and forms on an oriented surface  $M$  in  $\mathbb{R}^3$  before proving the Cohn-Vossen theorem.

Let  $N$  be the unit normal on  $M$ , let  $p \in M$ , and let  $e_1, e_2$  be a positively oriented orthonormal base field in the neighborhood  $U$  about  $p$ . Identifying  $p$  with the vector from the origin to  $p$ , define local functions  $y_1, y_2$  on  $U$ , and a global function  $y_3$  on  $M$ , by  $p = y_1(p)e_1 + y_2(p)e_2 + y_3(p)N$ . Define global 1-forms  $\alpha$  and  $\beta$  on  $U$  by

$$\alpha(X) = \langle p, e_1 \rangle \langle X, e_2 \rangle - \langle p, e_2 \rangle \langle X, e_1 \rangle$$

and  $\beta(X) = \alpha(LX)$ . One checks that  $\alpha$  is independent of the particular positively oriented base  $e_1, e_2$  used to define it, so  $\alpha$  and  $\beta$  are global 1-forms on  $M$ . We now compute  $d\alpha$  and  $d\beta$ . Let  $w_i, w_{ij}$  be the local forms belonging to the base  $e_1, e_2$  so  $w_{ij} = -W_{ji}$ . Then

$$L(X) = \bar{D}_X(N) = w_{13}(X)e_1 + w_{23}(X)e_2, w_{i3} = b_{1i}W_1 + b_{2i}w_2,$$

and  $b_{ij} = \langle Le_i, e_j \rangle$ . Thus  $\alpha = y_1w_2 - y_2w_1$  and  $\beta = y_1w_{23} - y_2w_{13}$ .

Since  $y_i = \langle p, e_i \rangle$  we have

$$\begin{aligned} dy_i(X) &= X \langle p, e_i \rangle \\ &= \langle \bar{D}_X p, e_i \rangle + \langle p, \bar{D}_X e_i \rangle \\ &= \langle X, e_i \rangle + \left\langle p, \sum_1^3 w_{ri}(X)e_r \right\rangle \\ &= w_i(X) + \sum_{r=1}^3 y_r w_{ri}(X). \end{aligned}$$

Thus, using the Cartan structural equations,

$$\begin{aligned} \mathbf{d}\alpha &= (w_1 + y_2w_{21} + y_3w_{31})w_2 - y_1(w_{21}w_1) - (w_2 + y_1w_{12} + y_3w_{32})w_1 \\ &\quad + y_2(w_{12}w_2) = 2w_1w_2 - y_3Hw_1w_2 = (2 - y_3H)v \end{aligned}$$

where  $v$  is the volume element. Similarly,  $\mathbf{d}\beta = (H - 2y_3K)v$ .

If  $M$  is compact, then

$$\int_M (H - 2y_3K) = \int_M \mathbf{d}\beta = \int_{\partial M} \beta = 0,$$

and

$$\int_M (2 - y_3H) = \int_M \mathbf{d}\alpha = \int_{\partial M} \alpha = 0,$$

by Stokes' theorem. The equality  $\int_M (H/2) = \int_M y_3K$  is called *Minkowski's formula*, and the other integral implies the area of  $M$  is  $\int_M y_3(H/2)$ . For other formulae of this type see [BF34].

The above paragraph provides two examples of “Chern's formula for theorems in differential geometry,” i.e., take a global 1-form  $w$  such that  $\mathbf{d}w = Fv$  where  $F$  is an “interesting” function, then state  $\int_M F = 0$ . Another example is that  $\int_M K = 0$  is a necessary condition that  $w_{12}$  be a global 1-form.

**Theorem 8.13 (Cohn-Vossen).** A compact surface of positive Gaussian curvature is rigid.

*Proof.* Let  $f : M \rightarrow M'$  be an isometry of such surfaces, and assume the origin to be inside  $M$  so  $y_3 > 0$ . Let  $L' = L_{M'} \circ f_*$ , then  $L$  and  $L'$  are positive definite on  $M$  since  $K = K' > 0$ . Let  $\Delta = \det(L - L')$ . We show  $L = \pm L'$  by showing  $\Delta = 0$  and apply the following lemma: if  $A$  and  $B$  are two positive definite quadratic forms on  $\mathbb{R}^2$  with  $\det A = \det B$ , then  $\det(A - B) \leq 0$ , and  $\det(A - B) = 0$  implies  $A = \pm B$ .

Let  $\beta' = \alpha \circ L'$  and, as above, we compute  $\mathbf{d}\beta' = [H' - y_3(2K - \Delta)]v$ . Hence

$$\int H' = \int \overline{y_3}(2k - \Delta) = \int H - \int y_3\Delta,$$

all integrals taken over  $M$ . Thus  $\int H' - \int H \geq 0$  since  $y_3 \leq 0$ , so  $\int H' \geq \int H$ . By symmetry we can reverse the inequality so  $\int H' = \int H$  and  $\int y_3 \Delta = 0$ , which implies  $\Delta = 0$ .  $\square$

**Theorem 8.14.** If  $f$  is an isometry between two oriented surfaces that preserves the mean curvature and the third fundamental form, and the mean curvature is never identically zero on any neighborhood, then the surfaces are congruent.

*Proof.* Let  $f : M \rightarrow M'$  and let  $L' = L_{M'} \circ f_*$ . Equality of the third fundamental forms implies  $\langle L^2 X, Y \rangle = \langle (L')^2 X, Y \rangle$  for all  $X, Y$  in  $\mathbf{T}_m M$  so  $L^2 = (L')^2$ . Using the characteristic equation for  $L$  and  $L'$  we have

$$HL = L^2 + KI = (L')^2 + K'I = H'L' = HL'.$$

Thus if  $H(m) \neq 0$ , then  $L = L'$  at  $m$ , and since  $H$  never vanishes identically on any neighborhood, we have  $L = L'$  on  $M$  by continuity.  $\square$

There is a theorem, similar to the preceding result, which states if  $f$  is a diffeo between two compact convex hypersurfaces that preserves the mean curvature and the third fundamental form, then the hypersurfaces are congruent or symmetric. For the proof of this result we refer the reader to [Che51], p. 29. Problem 77 shows one can relax the compactness assumption in the Cohn-Vossen theorem by assuming the third fundamental form is preserved.

The above theorems were included chiefly for their accessibility. Much better theorems have been proved (see [Pog56]) with weaker differentiability assumptions.

## Problems

75. Prove that the volume  $V_n$  of the unit sphere in  $\mathbb{R}^{n+1}$  is equal to  $\frac{2^{n+1} \pi^k k!}{n!}$  for even  $n = 2k$ .

76. If  $M$  is a compact surface in  $\mathbb{R}^3$  with constant mean curvature and  $y_3 > 0$  on  $M$ , show  $M$  is a sphere (see section 8.5)
77. If  $f$  is an isometry between two oriented surface in  $\mathbb{R}^3$  of positive Gaussian curvature which preserves the third fundamental form, show the surfaces are congruent or symmetric.
78. Use an integral argument to show there exists no compact minimal surface in  $\mathbb{R}^3$ .

## 9. Existence Theory

### 9.1 Involutive Distributions and the Frobenius Theorem

We prove the standard theorem on the existence of “integral manifolds” of a distribution following [Che46, p. 88]. The theorem also appears in [AM12, p. 147] with the terminology altered slightly.

In this section let  $M$  be a  $C^\infty$   $n$ -manifold. A  $k$ -dimensional distribution on a set  $A$  in  $M$  is a function  $P$  that assigns each point  $p$  in the  $k$ -dimensional subspace  $P_p$  of the tangent space  $\mathbf{T}_p M$ . We say  $P$  is  $C^\infty$  on  $A$  if  $A$  is open, and for each  $p \in A$  there are  $k$  independent  $C^\infty$  vector fields  $X_1, \dots, X_k$  which span  $P_m$  for all  $m$  in some neighborhood of  $p$ . A vector field  $X$  with domain  $B$  lies in  $P$  or is in  $P$  if  $B \subset A$  and  $X_p$  is in  $P_p$  for all  $p \in B$ . A  $C^\infty$  distribution  $P$  is integrable (involutive or closed) when it is closed under the bracket operation, i.e, if  $X$  and  $Y$  are any  $C^\infty$  fields with common domain that lie in  $P$ , then  $[X, Y]$  lies in  $P$ . A submanifold  $V$  of  $M$  is an integral submanifold or integral manifold of  $P$  if  $V$  is contained in the domain of  $P$ , and  $V_p = P_p$  for all  $p \in V$ ; thus the subspace of the tangent space  $\mathbf{T}_p M$  which belongs to  $V_p$  is exactly the subspace  $P_p$ .

The theorem proved below implies a  $C^\infty$  distribution has integral manifolds if and only if it is involutive. A slightly stronger statement is made involving the existence of a special coordinate system. First some terminology: if  $x_1, \dots, x_n$  is a coordinate system

on  $M$  with domain  $U$ , then define a slice of  $U$  to be any subset of  $U$  on which  $r$  of the functions  $x_1 \dots x_n$  are constant, where  $1 \leq r < n$ . Obviously, each slice of  $U$  is a submanifold of  $U$  (or  $M$ ).

**Theorem 9.1.** Let  $P$  be a  $k$ -dimensional involutive  $C^\infty$  distribution on  $M$ . For any  $m \in M$  there exists a coordinate system  $x_1, \dots, x_n$  with domain  $U$  including  $m$  such that the coordinate fields  $\frac{\partial}{\partial x_j}$  for  $j = 1, \dots, k$  span  $P$  at each point of  $U$ . Thus the slices of  $U$  for which  $x_{k+1}, \dots, x_n$  are constant are integral manifolds of  $P$ .

The theorem is proved by induction on  $k$ . The case  $k = 1$  is covered by the following lemma, and note in this case any distribution is automatically involutive.

**Lemma 9.2.** Let  $X$  be a  $C^\infty$  vector field on  $M$ ,  $p \in M$ , and  $X_p \neq 0$ , then there exists a coordinate system  $y_1 \dots y_n$  on a neighborhood  $U$  of  $p$  with  $X = \frac{\partial}{\partial y_1}$  on  $U$ .

*Proof of Lemma 9.2.* Let  $x_i = u_i \circ \phi$  be a coordinate system on the neighborhood  $V$  of  $p$  with  $x_i(p) = 0$  and  $\frac{\partial}{\partial x_1}(p) = X_p$ . Let  $X = \sum_1^n a_i \left( \frac{\partial}{\partial x_i} \right)$  where  $a_i$  are  $C^\infty$  real valued functions on  $V$  and  $a_1(p) \neq 0$ , and restrict  $V$  if necessary so  $a_1 \neq 0$  on  $V$ . Setting up the system of differential equations for the integral curves  $\sigma$  of  $X$  on  $V$ , we have

$$\frac{d(x_i \circ \sigma)}{dt} = a_i \circ \sigma \quad \text{or} \quad \frac{df_i}{dt} = a_i(f_1(t), \dots, f_n(t))$$

where  $f_i(t) = x_i \circ \sigma(t)$ . Applying an existence theorem from the theory of differential equations ([CL90, Chapter 1]) we obtain an  $r > 0$  and  $n$  functions  $F_i(t, a_1, a_2 \dots a_n)$  which are  $C^\infty$  on the neighborhood  $W$  of the origin in  $\mathbb{R}^{n+1}$  where  $|t| < r$  and  $|a_i| < r$  such that for  $i = 1, \dots, n$ :

- (1)  $F_i(0, a_1, a_2, \dots, a_n) = a_i$ ,
- (2)  $(F_1(b), \dots, F_n(b)) \in \phi(V)$  for  $b \in W$



(3) Letting

$$F(t, a_2, \dots, a_n) = \phi^{-1}[F_1(t, 0, a_2, \dots, a_n), \dots, F_n(t, 0, a_2, \dots, a_n)]$$

define a map  $F$  of  $B(0, r)$  in  $\mathbb{R}^n$  into  $V$ ; then for fixed  $a_2, \dots, a_n$  the curves

$$\sigma_{(a_2, \dots, a_n)}(t) = F(t, a_2, \dots, a_n)$$

are integral curves of  $X$ , i.e.,  $F_*\left(\frac{\partial}{\partial u_1}\right) = X$

For points  $(0, a_2, \dots, a_n)$  in  $B(0, r)$  we notice that

$$F(0, a_2, \dots, a_n) = \phi^{-1}(0, a_2, \dots, a_n);$$

hence  $F_*\left(\frac{\partial}{\partial u_i}\right)_{\text{origin}} = \frac{\partial}{\partial x_i}(p)$  for  $i = 2, \dots, n$ . Since  $F_*\left(\frac{\partial}{\partial u_1}\right)_b = X_{F(b)}$  for all  $b \in B(0, r)$  we have  $F_* = (\phi^{-1})_*$  at the origin in  $\mathbb{R}^n$ . Hence  $F_*$  is non-singular at the origin and by the Inverse Function Theorem  $F$  is a diffeo between a neighborhood of the origin and a neighborhood  $U$  of  $p$  with  $U \subset V$ . Finally, let  $y_i = u_i \circ F^{-1}$  on  $U$ .  $\square$

Intuitively, in the above proof we have changed the  $x_1, \dots, x_n$  coordinates about  $p$  by leaving the slice where  $x_1 = 0$  fixed, and replacing the “ $x_1$ -coordinate curves” by the integral curves of  $X$  emanating from this slice.

*Proof of Theorem 9.1.* Take the point  $m$  and take  $C^\infty$  fields  $X_1, \dots, X_k$  that span  $P$  on a neighborhood  $U_1$  on  $m$ . Apply the previous lemma to get a coordinate system  $y_1, \dots, y_n$  about  $m$  with domain  $U_2 \subset U_1$  such that  $\frac{\partial}{\partial y_1} = X_1$  on  $U_2$ , and assume  $y_i(m) = 0$ .

If  $k = 1$ , then the coordinate system  $y_1, \dots, y_n$  satisfies the conclusion of the theorem. If  $k > 1$ , we assume the theorem is true for the distributions of dimension less than  $k$ , and we define the  $(k - 1)$  dimensional distribution  $\bar{P}$  on  $U_2$  by

$$\bar{P}_p = \{X \in P_p : X_p y_1 = 0\} \text{ for } p \in U_2.$$

## Chapter 9: Existence Theory

---

This is a  $(k-1)$ -dimensional  $C^\infty$  distribution for it is spanned by the  $(k-1)$  independent  $C^\infty$  fields

$$Y_i = X_i - (X_i y_1) X_1 \text{ for } i = 2, \dots, k$$

It is involutive since if  $Y$  and  $Z$  are in  $\bar{P}$ , then  $[Y, Z]$  is in  $P$  and

$$[Y, Z]y_1 = Y(Zy_1) - Z(Yy_1) = 0 \text{ on } U_2,$$

so  $[Y, Z]$  is in  $\bar{P}$ .

Let  $V_0$  be the slice of  $U_2$  defined by  $y_1 = 0$ . Then for  $p \in V_0$ ,  $\bar{P}_p \subset (V_0)_p$ , so we apply the induction hypothesis to the distribution  $\bar{P}$  on the manifold  $V_0$  to obtain a coordinate system  $z_2, \dots, z_n$  on the neighborhood  $U_3$  about  $m \in v_0$  such that  $\frac{\partial}{\partial z_2}, \dots, \frac{\partial}{\partial z_k}$  span  $\bar{P}$  on  $U_3$ . We define the map  $\pi: U_2 \rightarrow V_0$  by  $\pi(p) = \phi^{-1}(0, y_2(p), \dots, y_n(p))$ , where  $\phi$  is the coordinate map so  $y_i = u_i \circ \phi$ . Let  $U_4 = \pi^{-1}(U_3)$  and define functions  $x_1, \dots, x_n$  on  $U_4$  by

$$x_1 = y_1, x_2 = z_2 \circ \pi, \dots, x_n = z_n \circ \pi$$

Then the functions  $x_1, \dots, x_n$  define a coordinate system in a neighborhood  $U$  of  $m$  with  $U \subset U_4$ ; indeed,  $\frac{\partial}{\partial x_1}(m) = \frac{\partial}{\partial y_1}(m)$ , while  $\frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n}$  span  $(V_0)_m$  at  $m$ .

We show  $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_k}$  span  $P$  on  $U$  by showing they span the same subspaces as  $X_1, Y_2, \dots, Y_k$ . Let  $Y_1 = X - 1$ , then we show

$$Y_i x_j = 0 \text{ for } i = 1, \dots, k \text{ and } j = k+1, \dots, n.$$

Since  $Y_1 = X_1 = \frac{\partial}{\partial x_1}$ , we immediately see  $Y_1 x_j = 0$  for  $j \neq 1$ . Since  $P$  is involutive, there are  $C^\infty$  functions  $g_{irs}$  on  $U$  such that for  $i \leq k$

---

## Section 9.1: Involutive Distributions and the Frobenius Theorem

---

and  $r \leq k$  we have  $[Y_i, Y_r] = \sum_{s=1}^k g_{irs} Y_s$ . thus for  $i = 2, \dots, k$  and  $j > k$ ,

$$Y_1(Y_i x_j) = [Y_1, Y_i]x_j = \sum_{s=1}^k g_{1is}(Y_s x_j).$$

This implies the functions  $Y_i x_j$  satisfy a linear homogeneous system of ordinary differential equations along any  $x_1$ -curve. But on  $V_0$ ,  $x_j = z_j$  for  $j > 1$  and  $Y_i x_j = Y_i z_j = 0$  on  $V_0$  for  $j > k$  because of the choice of coordinates  $z_2, \dots, z_n$ . Hence, by the uniqueness of solutions to systems of the above type,  $Y_i x_j = 0$  for  $i \leq k$  and  $j > k$  □

We use the theorem on involutive distributions to prove the classical Frobenius theorem on (total) partial differential equations (see [Lev77]). This theorem can be stated roughly as follows: there exist unique solution functions  $f_i(x_1, \dots, x_k)$ , with prescribed values at a point, to the system of partial differential equations

$$\frac{\partial f_1}{\partial x_j} = A_{ij}(x_1, \dots, x_k, f_1, \dots, f_d)$$

if and only if for all  $j \leq k, r \leq k$  and  $i \leq d$

$$\frac{\partial A_{ij}}{\partial x_r} + \sum_{s=1}^d \frac{\partial A_{ij}}{\partial f_s} A_{st} = \frac{\partial A_{ir}}{\partial x_j} + \sum_{s=1}^d \frac{\partial A_{ir}}{\partial f_s} A_{sj}$$

(which is merely what the chain rule demands if  $\frac{\partial^2 f_i}{\partial x_r \partial x_j} = \frac{\partial^2 f_t}{\partial x_j \partial x_t}$ ).

**Theorem 9.3 (Frobenius).** For  $1 \leq i \leq d$  and  $1 \leq j \leq k$ , let  $A_{ij}(x_1, \dots, x_k, u_1, \dots, u_d)$  be  $C^\infty$  real valued functions on an open set  $Q$  in  $\mathbb{R}^n$  where  $n = k + d$ , and we have labelled the coordinate functions of  $\mathbb{R}^n$  in order to conveniently express partial derivatives. Let  $(a; b) = (a_1, \dots, a_k, b_1, \dots, b_d)$  be in  $Q$ . Then there exists a unique set of  $C^\infty$  real valued functions  $f_1, \dots, f_d$  defined on a neighborhood  $V$  of  $a$  and satisfying the following three conditions:

- (1)  $f_i(a) = b_i$  or  $f(a) = b$ , where  $f$  is the mapping of  $V$  into  $\mathbb{R}^d$  defined by  $f(p) = (f_1(p), \dots, f_d(p))$
- (2) if  $p \in V$ , then  $(p; f(p))$  in  $Q$ , and
- (3) if  $p$  in  $V$ , then  $\frac{\partial f_i}{\partial x_j}(p) = A_{ij}(p; f(p))$

iff at every point of  $Q$ ,

$$\frac{\partial A_{ij}}{\partial x_r} + \sum_{s=1}^d \frac{\partial A_{ij}}{\partial u_s} A_{sr} = \frac{\partial A_{ir}}{\partial x_j} + \sum_{s=1}^d \frac{\partial A_{ir}}{\partial u_s} A_{sj}. \quad (4)$$

*Proof.* Let  $e_1, \dots, e_n$  be the usual global orthonormal vector fields on  $\mathbb{R}^n$ . We use the functions  $A_{ij}$  to define  $C^\infty$  vector fields  $Y_1, \dots, Y_k$  on  $Q$  by

$Y_r = e_r + \sum_{s=1}^d A_{sr} e_{k+s}$ . These vector fields are independent at each point of  $Q$  and hence they span a  $k$ -dimensional  $C^\infty$  distribution  $P$  on  $Q$ . We form brackets

$$\begin{aligned} [Y_r, Y_q] &= \left[ e_z + \sum_{s=1}^d A_{sr} e_{k+s}, e_q + \sum_{t=1}^d A_{tq} e_{k+t} \right] \\ &= \sum_{t=1}^d \left( \frac{\partial A_{tq}}{\partial x_r} + \sum_{s=1}^d A_{st} \frac{\partial A_{tq}}{\partial u_s} \right) e_{k+t} - \sum_{s=1}^d \left( \frac{\partial A_{sr}}{\partial x_q} + \sum_{t=1}^d A_{tq} \frac{\partial A_{sr}}{\partial u_t} \right) e_{k+s}, \end{aligned}$$

and thus by condition 4,  $[Y_r, Y_q] = 0$ .

Hence the distribution  $P$  is involutive and by the theorem above there exists an integral manifold  $U$  of  $T$  through  $(a; b)$  with  $U \subset Q$ . Let  $\phi: U \rightarrow \mathbb{R}^k$  by  $\phi(a'; b') = a'$ , then  $\phi_*(Y_r) = e_r$  and  $\phi_*$  is nonsingular on the tangent space of  $U$  at  $(a; b)$ . Thus there is a neighborhood  $V$  of  $a$  and a map  $F$  which is a diffeo of  $V$  onto  $F(V) \subset U$  such that  $F \circ \phi$  and  $\phi \circ F$  give the identity map on  $F(V)$  and  $V$ , respectively. Define  $f_1, \dots, f_d$  on  $V$  by  $F(p) = (p; f_1(p), \dots, f_d(p))$ . Then the functions  $f_1, \dots, f_d$  are  $C^\infty$  functions satisfying (1) and (2), and (3) follows since  $F_*(e_t) = Y_t$  for  $r \leq k_0$

The implication of the theorem in the other direction is trivial. □

Actually the Frobenius theorem in turn can be used to prove the theorem on involutive distributions. A  $k$ -dimensional distribution  $P$  about  $m$  can be carried to an open set  $Q$  in  $\mathbb{R}^n$  via a coordinate map. Furthermore one may choose the coordinate map so the induced distribution on  $Q$  is spanned by vectors  $Y_1, \dots, Y_k$  of the type defined above, and this defines functions  $A$ . The involutive condition will then imply  $[Y_r, Y_q] = 0$  since  $[Y_r, Y_q]$  must be a linear combination of  $Y_1, \dots, Y_k$  at each point. This implies the integrability condition 4 of the Frobenius theorem is satisfied which we then apply to get local integral manifolds. One actually has to state the Frobenius theorem to include the  $C^\infty$  dependence of the solution functions on the initial conditions (which follows from the Chevalley theorem) in order to obtain the full equivalence.

A first application of the Frobenius theorem provides a useful theorem concerning the existence of coordinate systems.

**Theorem 9.4.** Let  $M$  be an  $n$ -dimensional  $C^\infty$  manifold and let  $X_1, \dots, X_n$  be a set of independent  $C^\infty$  vector fields on a neighborhood  $U$  of  $m \in M$ . Then there exists a set of coordinate functions  $x_1, \dots, x_n$  defined on a neighborhood  $V$  of  $m$  with  $V \subset U$  and  $X_i = \frac{\partial}{\partial x_i}$  on  $V$  for all  $i$  iff  $[X_i, X_j] = 0$  for all  $i$  and  $j$ .

## 9.2 The Fundamental Existence Theorem for Hypersurfaces

Let  $U$  be an open set in  $\mathbb{R}^n$  on which is defined the real valued  $C^\infty$  functions  $g_{ij}$  and  $b_{ij}$  for  $1 \leq i, j \leq n$  such that the matrices  $(g_{ij})$  and  $(b_{ij})$  are symmetric and  $(g_{ij})$  is positive definite. Roughly speaking, we prescribe conditions which imply the existence of a coordinate system on a hypersurface of  $\mathbb{R}^{n+1}$  such that the matrices  $(g_{ij})$  and  $(b_{ij})$  are the coordinate representations of the first and second

fundamental forms, respectively. We demand that  $(g_{ij})$  and  $(b_{ij})$  satisfy the Gauss curvature and Codazzi-Mainardi equations, and explain this demand. On  $U$  define functions  $\Gamma_{jk}^i$ , in terms of the  $g_{ij}$  by the classical formula (see section 6.2) and define functions

- $w_{ij}(e_k) = \Gamma_{jk}^i$ ,
- $w_{n+1,j}(e_k) = -b_{jk}$ ,
- $w_{j,n+1} = \sum_{r=1}^n (g^{-1})_{jr} b_{rk}$ ,
- $w_{n+1,n+1}(e_k) = 0$ ,

for all  $i, j, k \leq n$ . Then if there was a coordinate system with coordinate fields  $e_1, \dots, e_n$  whose image set was  $U$ , the Gauss curvature equations and Codazzi-Mainardi equations imply (see section 6.6)

$$dw_{ij}(e_r, e_s) = - \sum_{k=1}^{n+1} w_{ik} \wedge w_{ks}(e_r, e_s) \quad (1)$$

and

$$dw_{j,n+1}(e_r, e_s) = - \sum_{k=1}^n w_{jk} \wedge w_{k,n+1}(e_r, e_s) \quad (2)$$

respectively. Thus we can say  $(g_{ij})$  and  $(b_{ij})$  satisfy the Gauss curvature and Codazzi-mainardi equations if 1 and 2 hold for the functions defined on  $U$  where the left sides are computed by

$$dw_{ij}(e_r, e_s) = \frac{\partial}{\partial u_r} w_{ij}(e_s) - \frac{\partial}{\partial u_s} w_{ij}(e_r), \text{ etc.}$$

**Theorem 9.5.** Let  $(g_{ij})$  and  $(b_{ij})$  be defined on  $U$  as described above and suppose they satisfy the Gauss curvature and Codazzi-Mainardi equations. Then for any point  $p \in U$ , there is a neighbourhood  $V \subset U$  and a  $C^\infty$  mapping  $F : V \rightarrow \mathbb{R}^{n+1}$  such that  $F(V)$  is an  $n$ -dimensional submanifold of  $\mathbb{R}^{n+1}$ ,  $F^{-1}$  is a coordinate map on  $F(V)$ , and  $(g_{ij})$  and  $(b_{ij})$  are the coordinate representation matrices of the first and second forms of  $F(V)$ , respectively.

Section 9.2: The Fundamental Existence Theorem for  
Hypersurfaces

---

**Proof.** let  $u_1, \dots, u_n$  be the natural coordinate functions on  $U$ . We seek

$(n + 1) \mathbb{R}^{n+1}$ -valued functions  $e_1, \dots, e_{n+1}$  defined on  $U$  that satisfy the Gauss equations and Weingarten equations, i.e.,

$$\frac{\partial e_i}{\partial u_j} = \sum_{k=1}^{n+1} w_{ki}(e_j) e_k = (\bar{D}_{e_j}(e_i)) \quad (3)$$

where  $j = 1, \dots, n$  and  $i = 1, \dots, n+1$ . Each of the equations in (3) has  $n + 1$  components, and the differentiation operator  $\frac{\partial}{\partial u_j}$  is applied to each component. In order to apply the Frobenius' theorem we compute  $\frac{\partial^2 e_i}{\partial u_k \partial u_j}$ , using (3) to obtain

$$\begin{aligned} \frac{\partial^2 e_i}{\partial u_k \partial u_j} &= \sum_r \left( \frac{\partial w_{ri}(e_j)}{\partial u_k} e_r + w_{ri}(e_j) \frac{\partial e_r}{\partial u_k} \right) \\ &= \sum_r \left( \frac{\partial w_{ri}(e_j)}{\partial u_k} e_r + \sum_s w_{ri}(e_j) w_{sr}(e_k) e_s \right) \end{aligned}$$

where we sum  $r$  and  $s$  from 1 to  $n + 1$ . The integrability conditions are the equation

$$\frac{\partial w_{ri}(e_j)}{\partial u_k} + \sum_s w_{si}(e_j) w_{rs}(e_k) = \frac{\partial w_{ri}(e_k)}{\partial u_j} + \sum_s w_{si}(e_k) w_{rs}(e_j), \quad (4)$$

which follows from (1) and (2).

At the origin in  $\mathbb{R}^{n+1}$  we choose initial vectors  $e_1, \dots, e_{n+1}$ , so

- $\langle e_i, e_j \rangle = g_{ij}(p)$ ,
- $\langle e_i, e_{n+1} \rangle = g_{i,n+1}(p) = 0$ ,
- $e_{n+1} = \frac{\partial}{\partial u_{n+1}}$

for  $i, j \leq n$  and  $e_1, \dots, e_{n+1}$  positively oriented. Applying the Frobenius theorem we obtain a neighborhood  $V_1$  of  $p$  and  $(n + 1) \mathbb{R}^{n+1}$ -valued  $C^\infty$  functions  $e_1, \dots, e_{n+1}$  that satisfy (3).

To check that  $\langle e_i, e_j \rangle = g_{ij}$  and  $\langle e_i, e_{n+1} \rangle = 0$  at all points on  $V_1$  for  $i, j \leq n$ , we must again apply the Frobenius theorem. Let  $G_{ij} = \langle e_i, e_j \rangle$  on  $V_1$  for  $i, j \leq n+1$ . Then by (3) and the product rule we have

$$\frac{\partial G_{ij}}{\partial u_k} = \sum_{r=1}^{n+1} [w_{ri}(e_k)G_{rj} + w_{rj}(e_k)G_{ir}] \quad (5)$$

on  $V_1$ . But from the definition of  $w_{ri}(e_k) = \Gamma_{ik}^r$  in terms of  $g_{ij}$  we find the functions  $g_{ij}$  also satisfy (5) where we define  $g_{i,n+1} \equiv \delta_{i,n+1}$ . By using (1) and (2) we verify the system (3) satisfies the necessary integrability conditions for the Frobenius theorem and since  $G_{ij}(p) = g_{ij}(p)$  we have  $G_{ij} = g_{ij}$  on a neighborhood  $V_2$  on  $p$ .

Define functions  $A_{ij}$  on  $V_2$  for  $i = 1, \dots, n+1$  and  $j = 1, \dots, n$  by  $e_j = (A_{1j}, \dots, A_{n+1j})$ , and consider the system of equations

$$\frac{\partial f_i}{\partial u_j} = A_{ij} \quad (6)$$

Here

$$\frac{\partial A_{ij}}{\partial u_k} = \frac{\partial A_{ik}}{\partial u_j} \text{ since } \frac{\partial e_j}{\partial u_k} = \frac{\partial e_k}{\partial u_j} \text{ (for } \Gamma_{jk}^i = \Gamma_{kj}^i).$$

Thus, letting  $f_i(p) = 0$  for  $i = 1, \dots, n+1$ , we apply the Frobenius theorem again to get  $C^\infty$  functions  $f_1, \dots, f_{n+1}$  on a neighborhood  $V_3$  of  $p$  with  $V_3 \subset V_2$ .

Finally, we define  $F : V_3 \rightarrow \mathbb{R}^{n+1}$  by  $F(m) = (f_1(m), \dots, f_{n+1}(m))$  for  $m \in V_3$ . Then  $F$  is  $C^\infty$  and  $F_*\left(\frac{\partial}{\partial u_j}(m)\right) = e_j(m)$  for  $j = 1, \dots, n$ . Thus  $F$  is a diffeo of a neighborhood  $V$  of  $p$  onto its image  $F(V) \subset \mathbb{R}^{n+1}$  and  $V \subset V_3$ . The map  $F^{-1}$  is a coordinate system on  $F(V)$  with coordinate vectors  $e_1, \dots, e_n$ , so  $g_{ij} = \langle e_i, e_j \rangle$  and

$$\langle Le_i, e_j \rangle = \left\langle \sum_r w_{r,n+1}(e_i)e_r, e_j \right\rangle = \sum_r w_{r,n+1}(e_i)g_{rj} = \sum_{r,s} (g^{-1})_{rs} b_{si} g_{rj} = b_{ji}$$

as desired. □



### 9.3 The Exponential Map

Let  $\mathbf{D}$  be a connexion on  $M^n$ . From section 5.1 we know for each vector  $X$ , tangent to  $M$  at  $m$ , there is a unique geodesic  $g_X(t)$  of the connexion  $\mathbf{D}$ , which is defined on a neighborhood of zero in  $\mathbb{R}$  with  $g_X(0) = m$  and tangent  $X$  at  $t = 0$ . Furthermore, for appropriate  $s \in \mathbb{R}$ ,  $g_{sX}(t) = g_X(st)$  by the nature of the differential equations defining the geodesics. This implies that  $g_{aX}(1)$  is defined if  $g_X(a)$  is defined, thus  $g_Y(1)$  is a well-defined point of  $M$  for  $Y$  near zero in  $\mathbf{T}_m M$ .

*Definition.* For  $Y \in \mathbf{T}_m M$ , we define  $\exp_m Y = g_Y(1)$  when the latter is defined. The map  $\exp_m$  is called the exponential map.

The name “exponential map” is used because in a special case for the general linear group  $\mathrm{GL}(n, \mathbb{R})$  it becomes the classical map,

$$A \mapsto e^A = I + A + \frac{A^2}{2!} + \cdots,$$

from the set of all  $n$  by  $n$  real matrices into the set of non-singular matrices (problem 81).

Our current objective is to obtain some important properties of the exponential map and to state these precisely we must use the tangent bundle  $\mathbf{T}M$  of  $M$ .

*Proposition 9.6.* Let  $N$  be the subset of  $\mathbf{T}M$  such that if  $(m, Y) \in N$  then  $\exp_m(Y)$  is defined and define the map  $\exp : N \rightarrow M$  by  $\exp(m, Y) = \exp_m(Y)$ . Then  $N$  is an open set and  $\exp$  is  $C^\infty$  on  $N$ . In particular let

$$\widehat{M} = \{(m, 0) \in \mathbf{T}(M) : m \in M\},$$

then there is an open set  $\widehat{N} \subset \mathbf{T}M$  such that  $\widehat{M} \subset \widehat{N} \subset N$ .

*Proof.* We do not completely prove the above proposition. Applying the local theory of differential equations, we prove the last statement of the theorem and we prove  $\exp : \widehat{N} \rightarrow M$  is  $C^\infty$ . Then

we sketch the proof that  $\exp$  is  $C^\infty$  on  $N$  and refer the reader to [Lan14].

Using the above notation, if  $g(t)$  is a geodesic in the neighborhood  $U$ , then

$$\frac{d^2(x_k \circ g)}{dt^2} + \sum_{i,j=1}^n \Gamma_{ij}^k \frac{d(x_i \circ g)}{dt} \frac{d(x_j \circ g)}{dt} = 0$$

for  $t$  such that  $g(t) \in U$ . For each point  $(m, 0) \in \mathbf{T}(M)$ , take a coordinate neighborhood  $U_m$  of  $m \in M$  and apply the existence and uniqueness theorem to the above differential equation to obtain a real number  $b > 0$ , a neighborhood  $V_m$  of  $(m, 0) \in \mathbf{T}M$  with  $V_m \subset \pi^{-1}(U_m)$ , a  $C^\infty$  map  $g : (-b, b) \times V_m \rightarrow M$ , such that for fixed  $(p, Y) \in V_m$ , the curve  $g_Y(t) = g(t; p, Y)$  is the unique geodesic defined on  $(-b, b)$  which passes through  $p$  with tangent  $Y$  at  $t = 0$ . Moreover, for  $(p, Y) \in V_m$  and  $a > 0$ , we have  $g_{aY}$  defined on  $(-b/a, b/a)$ , since

$g_{aY}(t) = g_Y(at) = g(at; p, Y)$ . Choose  $a > 0$  so  $a < b$  and let

$$W_m = \{(p, X) \in V_m : (p, X/a) \in V_m\}.$$

Then for  $(p, X) \in W_m$ ,  $\exp(p, X) = g(1; p, X) = g(a; p, X/a)$  is defined and  $\exp$  is  $C^\infty$  on  $W_m$ .

Let  $\hat{N} = U_m \cap W_m$ , and the last sentence of the theorem is proved.

For each  $(p, Y) \in \mathbf{T}M$ , choose  $a > 0$  so  $(p, aY) \in \hat{N}$ , and thus  $g_Y(t)$  is defined in some neighborhood of  $t = 0$ . As usual, for any curve  $g$  let  $T_g$  be its tangent vector and define the natural lifting of a curve  $g \in M$  to a curve  $\bar{g} \in \mathbf{T}M$  by  $Z(p, Y) = T_{\bar{g}_Y}(0)$ . Then  $Z$  is a  $C^\infty$  field on  $\mathbf{T}M$  by the above analysis, and if  $\sigma$  is an integral curve of  $Z$ , then  $\pi \circ \sigma$  is a geodesic in  $M$ . The field  $Z$  is called the geodesic flow field associated with the connexion. The fact that  $\exp$  is  $C^\infty$  on all of  $N$  now follows from Theorem 5 on p. 66 in [Lan14].  $\square$

**Corollary 9.7.** For fixed  $p \in M$ , the map  $\exp_p$  is a diffeo. of a neighborhood of  $0 \in \mathbf{T}_p M$  onto a neighborhood of  $p$ . Furthermore, if  $\eta_0 : \mathbf{T}_p M \rightarrow \mathbf{T}_0 \mathbf{T}_p M$  is the natural map of the tangent space at  $p$

onto its tangent space at 0, then  $(\exp_p)_* \circ \eta_0$  is the identity map on  $\mathbf{T}_p M$ .

*Proof.* The map  $\eta_0$  is defined by choosing any base  $e_1, \dots, e_n$  of  $\mathbf{T}_p M$  and letting  $z_1, \dots, z_n$  be its dual base. Then  $z_1, \dots, z_n$  are a global coordinate system on the vector space viewed as a  $C^\infty$  manifold. Let  $\eta_0(e_i) = \left(\frac{\partial}{\partial z_i}\right)_0$  for all  $i$ . This map  $\eta$  is independent of the particular base  $e_1, \dots, e_n$ ; furthermore, by evaluating the global fields  $\frac{\partial}{\partial z_i}$  at any point  $Y \in \mathbf{T}_m M$ , we obtain a natural isomorphism  $\eta_Y : \mathbf{T}_p M \rightarrow \mathbf{T}_Y \mathbf{T}_p M$ . In these notes, for any  $X \in \mathbf{T}_p M$  we let  $\bar{X}$  be the natural constant vector field on  $\mathbf{T}_p M$  associated with  $X$ , where  $\bar{X}_Y = \eta_Y(X)$ .

Take  $X \in \mathbf{T}_p M$ , then  $\eta_0(X) = \bar{X}_0$ . To compute  $(\exp_p)_* \bar{X}_0$  we note  $\bar{X}_0$  is the tangent vector at  $t = 0$  to the ray  $\gamma(t) = tX \in \mathbf{T}_p M$ . The curve

$\exp_p \circ \gamma(t) = \exp_p tX$  is by definition the geodesic through  $p$  with tangent  $X$  at  $t = 0$ . Thus  $(\exp_p)_* \bar{X}_0 = X$ . Thus  $(\exp_p)_*$  is non-singular and onto at the origin in  $\mathbf{T}_p M$ . The corollary now follows by applying the Inverse Function theorem.  $\square$

**Corollary 9.8.** Let  $G : \widehat{N} \rightarrow M \times M$  by  $G(p, Y) = (p, \exp_p Y)$ . Then  $G$  is  $C^\infty$  and  $G_*$  is non-singular and onto at all points  $(p, 0) \in \Gamma(M)$ .

*Proof.* Let  $\pi_i : M \times M \rightarrow M$  by  $\pi_i(m_1, m_2) = m_i$  for  $i = 1, 2$ . Each  $\pi_i$  is  $C^\infty$ . Since  $\pi_i \circ G = \pi_i$  and  $\pi_2 \circ G = \exp$ , the map  $G$  is  $C^\infty$  on  $\widehat{N}$ .

The tangent space  $\widehat{N}_{(p,0)}$  is naturally isomorphic to  $\mathbf{T}_p M \times \mathbf{T}_0 \mathbf{T}_p M$ , while the tangent space to  $M \times M$  at  $G(p, 0) = (p, p)$  is naturally isomorphic to  $\mathbf{T}_p M \times \mathbf{T}_p M$ . In terms of these natural isomorphic spaces,  $G_*$  is the identity on the first factor and  $(\exp_p)_*$  on the second factor. Hence, by Corollary 9.7,  $G_*$  is non-singular at  $(p, 0)$ .  $\square$

We apply Corollary 9.7 to obtain normal coordinate systems. For any  $m \in M$ , let  $e_1, \dots, e_n$  be a base of  $\mathbf{T}_m M$ , let  $z_1, \dots, z_n$  be

its dual base, and let  $\bar{U}$  and  $U$  be neighborhoods of  $O \in T_m M$  and  $m \in M$ , respectively, such that  $\exp_m$  is a diffeo. of  $\bar{U}$  onto  $U$  whose inverse we denote by  $\exp^{-1}$ . Then define  $C^\infty$  functions  $x_1, \dots, x_n$  on  $U$  by  $x_i = z_i \circ \exp^{-1}$  for all  $i$ . These functions  $x_1, \dots, x_n$  define a normal coordinate system (of the connection  $\mathbf{D}$ ) on  $U$ . The curves  $\sigma \in U$  such that  $x_i \circ \sigma(t) = a_i t$  for constants  $a_1, \dots, a_n$ , are geodesics emanating from  $m$  (at  $t = 0$ ), and if  $\Gamma_{jk}^i$  are the connexion functions on  $U$  for this coordinate system, then  $\Gamma_{jk}^i(m) = 0$ , provided the connexion has zero torsion.

One verifies this last statement by letting  $X_i = \frac{\partial}{\partial x_i}$  and then

$$\mathbf{D}_{X_i}(X_j) = \sum_{k=1}^n \Gamma_{jk}^k X_k$$

by definition of  $\Gamma_{jk}^k$ . Since the curve  $\sigma$  with

$$x_i \circ \sigma(t) = t, x_j \circ \sigma(t) = t, \text{ and } x_k \circ \sigma(t) = 0$$

for  $k \neq i$  or  $j$ , is a geodesic, its tangent  $X_i + X_j$  satisfies the condition

$$0 = \mathbf{D}_{(X_i + X_j)}(X_i + X_j) = \mathbf{D}_{X_i}X_i + \mathbf{D}_{X_i}X_j + \mathbf{D}_{X_j}X_i + \mathbf{D}_{X_j}X_j.$$

Thus at  $m$ ,  $\mathbf{D}_{X_i}X_i = 0$  for all  $i$ , since each coordinate curve emanating from  $m$  is a geodesic, and if  $\mathbf{D}$  has zero torsion, then

$$0 = 2(\mathbf{D}_{X_i}X_j)_m = 2 \sum_k \Gamma_{ji}^k(m) X_k$$

so  $\Gamma_{ji}^k(m) = 0$  for all  $i, j$ , and  $k$ .

We apply Corollary 9.8 to obtain Fermi coordinates along a curve. Let  $\sigma$  be a  $C^\infty$  curve in  $M$  that is univalent on the open interval  $I \subset \mathbb{R}$ . Let  $e_1, \dots, e_n$  be  $C^\infty$  fields on  $\sigma$  that are independent at each  $\sigma(t)$  and  $e_n(t) = T_\sigma(t)$  for all  $t \in I$ . Let  $z_1, \dots, z_n$  be the dual base to  $e_1, \dots, e_n$  for each  $t$ . By Corollary 9.8, there is a neighborhood  $V$  of  $\widehat{M} \subset TM$  such that  $G$  is a diffeo. of  $V$  onto a

neighborhood  $N_M$  of the diagonal in  $M \times M$ . Let

$$U = \{(m, Y) \in V : m = \sigma(t) \text{ and } z_n(Y) = 0 \text{ for some } t \in I\}.$$

Then  $F = G|_U$  is a 1 to 1  $C^\infty$  map of the submanifold  $U$  into  $M \times M$ . Moreover  $F_*$  is non-singular at each point of  $U$ , so  $F$  is an imbedding of  $U$  into  $M \times M$ . The map  $H = \pi_2 \circ F$  then gives a 1 to 1  $C^\infty$  map of  $U$  onto an open neighborhood  $W$  of the image set  $\sigma(I)$ . Define Fermi coordinate  $y_1, \dots, y_n$  on  $p \in W$  by letting  $H^{-1}(p) = (\sigma(t), Y) \in U$  and  $y_i(p) = z_i(Y)$  for  $i = 1, \dots, n-1$  and  $y_n(p) = t$ .

More special types of Fermi coordinates can be defined by taking  $e_1, \dots, e_n$  to be a parallel base along a geodesic  $\sigma$ , and in the Riemannian case, one can take an orthonormal parallel base along a geodesic.

## 9.4 Convex Neighborhoods

This section is devoted to proving the following theorem, due to J. H. C. Whitehead [Whi32].

**Theorem 9.9.** Let  $M$  be a  $C^\infty$  manifold and  $D$  be a  $C^\infty$  connexion on  $M$ . Then for any point  $m$  in  $M$  there is a neighbourhood  $U$  of  $m$  that is convex; i.e., for any two points in  $U$  there is a unique geodesic of  $D$  which joins the two points and lies in  $U$ .

*Proof.* We may assume  $D$  has zero torsion, since by section 5.4 there is a unique torsion-free connexion with the same geodesics. The theorem is local, and we work completely in one coordinate neighbourhood of  $m$ . From the previous section, we choose a normal coordinate system  $x_1, x_2, \dots, x_n$  about  $m$  with domain  $A$ , thus  $x_i(m) = 0$  and  $\Gamma_{jk}^i(m) = 0$  for all  $i, j$  and  $k$ . Let

- $d(p, q)$  be a local metric on  $A$  defined by

$$d(p, q) = \sqrt{\sum_i (x_i(p) - x_i(q))^2},$$

- $f(p) = d(p, m)$ ,
- $B(p, c) = \{q \in A : d(p, q) < c\}$  for  $p \in A$ .
- $|X| = \sqrt{\sum_i (\mathbf{d}x_i(X)^2)}$  for  $p \in A$  and  $X \in \mathbf{T}_p M$ .

By Corollary 9.8 in section 9.3, for each  $p \in A$  there is a real number  $r_p > 0$  so that  $G$  is a diffeomorphism on the set  $(q, X)$  where  $q \in B(p, r_p)$  and  $d(q, \exp_p X) < r_p$ . Take  $c > 0$  so  $\bar{B} = \bar{B}(m, c) \subset A$ . For each  $p \in \bar{B}$  we obtain an integer  $r_p > 0$ . The family of neighbourhoods  $B(p, r_p)$  for  $p \in \bar{B}$  is a covering of the compact set  $\bar{B}$ , hence we may select a finite subcovering of neighbourhoods belonging to  $p_1, p_2, \dots, p_k$ . Let  $s = \min\{r_1, \dots, r_k\}$ . Then for any  $p \in \bar{B}$ ,  $\exp_p$  maps a neighbourhood  $\bar{U}_p$  of the origin in  $\mathbf{T}_p M$  diffeomorphism onto  $B(p, s)$ . This follows since  $p \in B(p_j, r_j)$  for some  $j$  and hence  $G$  is a diffeomorphism on the set  $(q, X)$  for  $q \in B(p_j, r_j)$  and  $d(q, \exp_p X) < r_j$ . We fix  $q = p$ , and  $\exp_p$  is a diffeomorphism of a neighbourhood  $\bar{V}_p$  of  $\mathbf{0}$  in  $\mathbf{T}_p M$  onto  $B(p, r_j)$  and  $s \leq r_j$ . We have proved the following:

**Lemma 9.10.** For any  $c > 0$  with  $\bar{B}(m, c) \subset A$ , there exists an  $s > 0$  such that for  $p \in \bar{B}(m, c)$  the map  $\exp_p$  is a diffeomorphism from a neighbourhood  $\bar{U}_p$  of  $\mathbf{0}$  in  $\mathbf{T}_p M$  onto  $B(p, s) \subset A$ .

We now prove two lemmas that complete the proof of the theorem.

**Lemma 9.11.** There exists a real number  $a$ ,  $0 < a < 1$  and  $\bar{B}(m, a) \subset A$ , such that if  $0 < b < a$  and  $g$  is a geodesic with  $\mathbf{T}g$  and  $f \circ g(0) = b$ ,  $T_{g(0)}f = 0$ , then  $f \circ g$  has a strict relative minimum at  $g(0)$ . Thus if  $g$  is tangent to the “sphere about  $m$  of radius  $b$ ” at  $g(0)$ , then  $g$  lies outside of  $B(m, b)$  near  $g(0)$ .

*Proof.* We may assume  $|T_{g(0)}| = 1$ . Let  $T = \sum_j a_j X_j$  where  $X_j = \frac{\partial}{\partial x_j}$

and

$a_j \circ g = \left(\frac{\mathbf{d}}{\mathbf{d}t}\right)(x_j \circ g)$ , and we assume  $T_{g(0)}$  is extended to a  $C^\infty$  field

in a neighbourhood of  $g(0)$ . Since  $f = \sqrt{\sum_i x_i^2}$  we have

$$Tf = \sum_j a_j (X_j f) = \frac{1}{f} \sum_j a_j x_j$$

and

$$T^2 f = \sum_k a_k \left[ -\frac{x_k}{f^3} \left( \sum_j a_j x_j \right) + \frac{1}{f} \left( \sum_j (X_k a_j) x_j a_k \right) \right].$$

At  $t = 0$ , or at  $g(0)$ ,  $Tf = 0$ ; hence

$$T^2 f = \frac{1}{b} \left[ \sum_k a_k^2 + \sum_{k,j} a_k x_j (X_k a_j) \right]$$

But at  $g(0)$ ,

$$\sum_k a_k^2 = |T|^2 = 1 \quad \text{and} \quad \sum_k a_k (X_k a_j) + \sum_{r,s} \Gamma_{rs}^j a_r a_s = 0,$$

since  $g$  is a geodesic. Thus

$$T^2 f = \frac{1}{b} \left[ 1 - \sum_{j,r,s} x_j \Gamma_{rs}^j a_r a_s \right].$$

Choose  $a > 0$  and  $a < 1$  so for points  $p$  with  $f(p) \leq a$ ,  $|\Gamma_{jk}^i(p)| < 1/2n^3$  for all  $i, j$  and  $k$ , which is possible since  $\Gamma_{jk}^i$  continuous and  $\Gamma_{jk}^i(m) = 0$ . Then, at  $g(0)$ ,

$$\left| \sum_{j,r,s} x_j \Gamma_{rs}^j a_r a_s \right| \leq \left( \frac{1}{2n^3} \right) \left( \sum_{j,r,s} 1 \right) \leq \frac{1}{2},$$

hence  $T^2 f(g(0)) > 0$ , which implies  $f \circ g$  has a strict relative minimum at  $\mathbf{0}$ . □

**Lemma 9.12.** Let  $a$  be given by Lemme 9.11 and apply Lemma 9.10 with  $c = a/2$  to obtain  $s > 0$  with  $s < (2/3)a$ . Then  $B(m, s/2)$  is convex.

*Proof.* Choose any  $p$  and  $q$  in  $B(m, s/2)$ . By Lemma 9.10 there is a geodesic  $g$  defined on some interval  $[0, u]$  with  $g(0) = p$ ,  $g(u) = q$ , and  $g(t) \in B(p, s)$  for all  $t \in [0, u]$ . We show  $f \circ g(t) < s/2$  for all  $t \in [0, u]$ . Let  $v$  be a number in  $[0, u]$  where  $f \circ g$  attains its maximum value. Then  $f \circ g(v) < a$  since

$$f \circ g(v) = d(m, g(v)) \leq d(m, p) + d(p, g(v)) \leq s/2 + s < a.$$

Suppose  $f \circ g(v) \geq s/2$ . Then  $(f \circ g)'(v) = 0$  and  $f \circ g(v) < a$  which implies by Lemma 9.11 that  $f \circ g$  has a strict relative minimum at  $v$  which gives a contradiction. Hence  $B(m, s/2)$  is convex.  $\square$

Our theorem follows.  $\square$

## 9.5 Special Coordinate Systems

Let  $M$  be a Riemannian  $n$ -manifold, let  $\phi$  be a coordinate map on  $M$  with domain  $U$  and  $x_i = u_i \circ \phi$ , and let  $X_i = \frac{\partial}{\partial x_i}$ . The coordinate system  $x_1, \dots, x_n$  is orthogonal if  $\langle X_i, X_j \rangle = 0$  for  $i \neq j$ . If the map  $\phi$  is a conformal map of  $U$  into  $\mathbb{R}^n$  (with respect to the canonical Riemannian metric on  $\mathbb{R}^n$ ), then the coordinate system is *isothermal* or *conformal* (and hence also orthogonal). When  $M^n$  is a hypersurface in some  $\overline{M}^{n+1}$ , the coordinate system is *principal* if each  $X_i$  is a principal vector, and it is *asymptotic* if each  $X_i$  is an asymptotic vector.

In this section we study the existence of such special coordinate systems when  $n = 2$ . Orthogonal systems and conformal systems exist about any point, and the latter may be used to define a Riemann surface structure on  $M$ . Principal coordinates exist of necessity about any non-umbilical point on a surface, while they may or may not exist about an umbilic. We show asymptotic coordinates exist in some special cases, e.g., about a point of a surface which has a neighborhood on which the curvature is a negative constant, and about a non-umbilical point on a negative constant, and about



a non-umbilical point on a minimal surface (problem 88).

**Theorem 9.13 (Gauss 1827).** Let  $\gamma$  be an arbitrary univalent curve in  $M^2$  parameterized by arc length on  $(a, b)$ , let  $X$  be the (unit) tangent to  $\gamma$ , and let  $Y$  be a unit  $C^\infty$  field along  $\gamma$  such that  $\langle X, Y \rangle = 0$ . Then the Fermi coordinate system induced by  $Y$  on a neighborhood of  $\gamma$  is an orthogonal coordinate system about  $\gamma$  which is called a set of “geodesic parallel coordinates.” This proves the existence of orthogonal coordinates about any point on a two-dimensional Riemannian manifold.




Figure 9.1: Fermi Coordinates

*Proof.* Let  $\phi$  be the Fermi coordinate map from the neighborhood  $U$  of  $\gamma$  onto the set  $V$  in  $\mathbb{R}^2$ .

Then for  $(t, s)$  in  $V$ ,  $\phi^{-1}(t, s) = \exp_{\gamma(t)} sY$ . We let  $X$  and  $Y$  be the coordinate fields on  $U$  which extend  $X$  and  $Y$  along  $\gamma$ . Since the  $\gamma$ -curves are geodesics parameterized by arc length,  $D_Y Y = 0$  and  $\langle Y, Y \rangle = 1$ , where  $D$  is the Riemannian connexion. We compute

$$Y \langle X, X \rangle = \langle D_Y X, X \rangle + \langle X, D_Y X \rangle = \langle D_Y X, X \rangle = \frac{1}{2} X \langle Y, Y \rangle = 0,$$

since the torsion is zero, so  $D_Y X - D_X Y = [Y, X] = 0$ . Thus  $\langle X, Y \rangle$  is constant along the  $y$ -curves and since  $\langle X, Y \rangle = 0$  on  $\gamma$  we have  $\langle X, Y \rangle = 0$  on  $U$ . □

One way to paraphrase the above situation is to say “if segments of equal length (lying in  $U$ ) are laid off along geodesics that are orthogonal to a univalent curve  $\gamma$ , then their endpoints determine an orthogonal trajectory to the family of geodesics.

**Theorem 9.14.** If  $m$  is a non-umbilical point on a surface  $M$  in  $\mathbb{R}^3$ , then there exists a set of principal coordinates in a neighborhood  $U$  of  $m$ .

*Proof.* Since  $m$  is non-umbilic, there is a neighborhood  $V$  of  $m$  which contains no umbilics. Assume  $V$  is oriented via a unit field  $N$ , and let  $L(X) = \bar{D}_X N$  as usual, where  $\bar{D}$  is the Riemannian connexion on  $\mathbb{R}^3$ . Let  $X$  and  $Y$  be  $C^\infty$  orthonormal principal vector fields on  $V$  with  $L(X) = kX$ ,  $L(Y) = hY$ , and  $k < h$ , which corresponds to the notation of Chapter 3. We seek non-vanishing  $C^\infty$  functions  $f$  and  $g$  defined on a neighborhood of  $m$  such that the fields  $Z = fX$  and  $W = gY$  satisfy the condition  $[Z, W] = 0$ . Finding  $f$  and  $g$ , we can apply theorem 9.1 to obtain the desired principal coordinates.

We compute

$$[fX, gY] = f(Xg)Y - g(Yf)X + fg(aX - bY),$$

where

$$a = (Yk)/(h - k) \text{ and } b = -(Xh)/(h - k)$$

by theorem 3.2. Hence  $[Z, W] = 0$  if  $(Xg) - bg = 0$  and  $(Yf) - af = 0$ . Thus we may prescribe  $g = 1$  on the integral curve of  $Y$  through  $m$ , and then on each integral curve  $\gamma(t)$  of  $X$  we have the differential equation

$$\frac{dg \circ \gamma(t)}{dt} - (b \circ \gamma)(t)(g \circ \gamma)(t) = 0.$$

From the existence theory of ordinary differential equations we get  $g$  defined and  $C^\infty$  on a neighborhood of  $m$  with  $g > 0$ . Similarly, we obtain  $f$ . □

One can write the differential equations  $Xg = bg$  and  $Yf = af$  as

---

## Section 9.6: Isothermal Coordinates and Riemannian Surfaces

---

first-order linear partial differential equations in terms of a coordinate system  $u, v$  about  $m$ . This follows, since

$$X = b_1 \frac{\partial}{\partial u} + b_2 \frac{\partial}{\partial v} \quad \text{and} \quad Y = a_1 \frac{\partial}{\partial u} + a_2 \frac{\partial}{\partial v}$$

defines  $C^\infty$  functions  $a_i$  and  $b_i$ , and then one must solve,

$$b_1 \frac{\partial g}{\partial u} + b_2 \frac{\partial g}{\partial v} = bg \quad \text{and} \quad a_1 \frac{\partial f}{\partial u} + a_2 \frac{\partial f}{\partial v} = af.$$

**Theorem 9.15.** If  $m$  is contained in the neighborhood  $U$  on a surface with constant  $K = -a^2 < 0$  on  $U$ , then there exists a set of asymptotic coordinates about  $m$ .

*Proof.* Let  $X$  and  $Y$  be orthonormal principal fields on  $U$  with  $LX = kX$  and  $LY = hY$ ,  $k < 0 < h$ . Let

- $b = \sqrt{a^2 + k^2}$ ,
- $Z = b(aX - kY)$ ,
- $W = (-aX - kY)$ .

Then  $\langle LZ, Z \rangle = \langle LW, W \rangle = 0$ , and  $Z$  and  $W$  are clearly independent. Using theorem 3.2, one computes  $[Z, W] = 0$ . Hence, the desired coordinates exist. □

## 9.6 Isothermal Coordinates and Riemannian Surfaces

The principal reference for this section is [Sam55]. Let  $M$  be a Riemannian 2-manifold.

Let  $x, y$  be an arbitrary coordinate system on a neighbourhood  $U$  of  $M$ . We seek functions  $f$  and  $g$  so the map  $p \rightarrow (f(p), g(p))$  will define a conformal coordinate system about  $m$  in  $U$ . If  $f$  and  $g$  exist, let

$$\bullet E = \left\langle \frac{\partial}{\partial f}, \frac{\partial}{\partial g} \right\rangle,$$

- $F = \left\langle \frac{\partial}{\partial f}, \frac{\partial}{\partial g} \right\rangle,$
- and  $G = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right\rangle.$

Then

$$\mathbf{grad} f = \frac{1}{W^2} \left( G \frac{\partial}{\partial f} - F \frac{\partial}{\partial g} \right)$$

where  $W = \sqrt{EG - F^2}$ . If  $f$  and  $g$  are orthogonal coordinates, then  $F = 0$ . If they are also conformal coordinates, then  $E = G$  and  $|\mathbf{grad} f|^2 = 1/E = |\mathbf{grad} g|^2$ . Thus coordinates  $f$  and  $g$  are conformal iff  $\langle \mathbf{grad} f, \mathbf{grad} g \rangle = 0$  and  $|\mathbf{grad} f|^2 = |\mathbf{grad} g|^2$ .

In terms of the  $x, y$  coordinate system,

$$\langle \mathbf{grad} f, \mathbf{grad} g \rangle = g_x(Gf_x - Ff_y) - g_y(Ff_x - Ef_y)$$

where  $g_x = \frac{\partial g}{\partial x}$ , etc., and  $E, F$  and  $G$  now belong to  $x$  and  $y$ , i.e.,  $E = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial x} \right\rangle$ , etc. Thus  $\langle \mathbf{grad} f, \mathbf{grad} g \rangle = 0$  if there is a function  $\rho$  on  $U$  with

$$g_x = \rho(Ff_x - Ef_y) \text{ and } g_y = \rho(Gf_x - Ff_y). \quad (1)$$

Then  $|\mathbf{grad} g|^2 = \rho^2 W^2 |\mathbf{grad} f|^2$ , so let  $\rho = 1/W$ . The equations (1) become a generalization of the Cauchy-Riemann equations. For a particular  $f$ , one can solve the system (1) for  $g$  iff  $g_{xy} = g_{yx}$  or

$$\frac{\partial}{\partial x} \left[ \frac{Gf_x - Ff_y}{\sqrt{EG - F^2}} \right] + \frac{\partial}{\partial y} \left[ \frac{Ef_y - Ff_x}{\sqrt{EG - F^2}} \right] = 0 \quad (2)$$

Equation (2) is the classical *Beltrami equation*, a generalized form of the Laplace equation. Indeed, the left side of (2) is  $W\Delta f$ . Classically,  $\langle \mathbf{grad} f, \mathbf{grad} g \rangle$  is called the *first Beltrami operator* on  $f$  and  $g$  and the Laplacian  $\Delta$  is called the *second Beltrami operator*.

The theory of elliptic partial differential equations gives the existence of non-trivial solutions of (2) about a point in  $U$  which proves the following theorem.

**Theorem 9.16.** There exists a system of isothermal (conformal) coordinates about any point of Riemannian 2-manifold.

On manifolds  $M$  as described in this theorem, if we restrict ourselves to conformal coordinate systems then, when the domains of these coordinate systems intersect, they induce a conformal map from one open set of  $\mathbb{R}^2$  onto another. Since  $\mathbb{R}^2$  is the underlying set for the space of complex numbers, these conformal maps must be given by analytic functions from one open set of onto another. Thus at each point  $m$  of  $M$  we have diffeomorphisms of a neighborhood of  $m$  onto an open set in  $\mathbb{C}$  which are related by analytic functions on the intersection of their domains. When  $M$  is covered by neighborhoods such that the analytic functions induced by overlapping neighborhoods are orientation preserving, then  $M$  is called a *Riemann surface* and the study of these objects leads to a rich theory (see [AL60]).

## Problems

79. Let  $T$  be a  $C^\infty$  vector field on the Riemannian manifold  $M$  and define  $A_T : \mathbf{T}_m M \rightarrow \mathbf{T}_m M$  by  $A_T(X) = \mathbf{D}_X T$ , where  $\mathbf{D}$  is the Riemannian connexion.

(i) Show that  $\operatorname{div} T = \operatorname{tr} A_T$ .

(ii) Show  $A_T$  is self-adjoint iff  $d \circ G(T) = 0$  ( $T$  is closed).

(iii) Let  $(T^\perp)_m = \{X \in \mathbf{T}_m M : \langle X, T_m \rangle = 0\}$ .

(iv) If  $T$  is closed, show  $T^\perp$  is an integrable  $(n - 1)$ -dim distribution on the subset of  $M$  where  $T \neq 0$ .

80. (Frobenius) Let  $W_1, \dots, w_k$  be a set of independent  $C^\infty$  1-forms on a  $C^\infty$   $n$ -manifold  $M$  with  $k < n$ . Define an  $(n - k)$ -dim distribution  $P$  on  $M$  by

$$P_m = \{X \in \mathbf{T}_m M : w_i(X) = 0 \text{ for } i = 1, 2, \dots, k\}$$

Show that  $P$  is integrable iff

$$dw_i = \sum_{1 \leq r \leq s \leq k} a_{irs} w_r \wedge w_s$$

for all  $i$ . (For generalizations of this result, see Kuranishi [Kur57] or Johnson [Joh60].)

81. If  $G = \mathbf{GL}(n, \mathbb{R})$ ,  $I$  is the identity in  $G$ ,  $A \in \mathbf{T}_I G$ , and

$$\sigma : t \rightarrow e^{tA} = I + tA + \frac{(tA)^2}{2!} + \cdots + \frac{(tA)^n}{n!} + \cdots,$$

show  $\sigma(t)$  is a 1-parameter subgroup of  $G$  with tangent  $A$  at  $t = 0$ . Thus show  $e^{tA} = \exp_I(tA)$  for all  $t$  (see problem 46).

82. Show the map  $(m, X) \rightarrow |X|$  is  $C^\infty$  on the set

$$N = \{(m, X) \in \mathbf{T}M : X \neq 0\}.$$

83. If  $M$  is a Riemannian manifold and  $A$  is a compact set in  $M$ , show that there exists a real number  $r > 0$  such that the ball  $B(m, r)$  is convex for all  $m$  in  $A$ .

84. If  $G$  is a Lie group,  $g \in G$ ,  $X$  in the Lie algebra, and  $g = \exp(X)$ , show that  $h^2 = g$  where  $h = \exp(X/2)$ . If

$$h \in \mathbf{SL}(2, \mathbb{R}) = \{g \in \mathbf{GL}(2, \mathbb{R}) : \det(g) = 1\},$$

show the  $\text{tr}(h^2) \geq -2$ . Use this to prove the exp map is not always onto even when the connexion is complete.

85. Let  $D$  be a connexion on  $M$ .

(i) Show the curvature  $R \equiv 0$  iff the horizontal distribution  $H$  on  $B(M)$  is integrable (section 5.5).

(ii) Show that  $R \equiv 0$  implies parallel translation is independent of the path (problem 45).

86. Show that there exists at least one umbilic on any compact convex  $C^\infty$  surface in  $\mathbb{R}^3$ . (It was conjectured by Caratheodory, and proven by Bol and Hamburger respectively, that a compact convex surface has at least two umbilics.)

87. (i) If  $M$  is a surface in  $\mathbb{R}^3$ ,  $U$  a coordinate domain on  $M$  with coordinate fields  $X$  and  $Y$ , show the area of  $U$  is equal to

$$\int_U \sqrt{\langle X, Y \rangle \langle Y, Y \rangle - \langle X, Y \rangle^2}.$$

(ii) Let  $f$  be in  $C^\infty(U, \mathbb{R})$ , and define a normal deformation belonging to  $f$  by  $\phi_t(p) = p + tf(p)N_p$  for  $p \in U$  and  $N$  a  $C^\infty$  unit normal on  $U$ . Let  $J(t)$  be the area of  $\phi_t(p)$ . Show that  $J'(0) = 0$  for all  $f$  iff  $U$  is a minimal surface ( $H \equiv 0$ ).

88. (i) Show that about any non-umbilic point on a minimal surface there exists an isothermal coordinate system  $x, y$  whose coordinate systems are lines of curvature.

(ii) Show the functions  $z = (x + y)/2$  and  $w = (x - y)/2$  define an isothermal coordinate system whose coordinate curves are asymptotic curves which bisect the  $x, y$  coordinate curves.

89. Using the notation of section 3.4, let  $u, v$  be conformal coordinates on domain  $B$  with  $E = G = \langle T_u, T_u \rangle$ .

(i) Show  $T_{uu} + T_{vv} = -HGN$ . If  $f$  is  $C^\infty$  on  $B$ ,

(ii) show  $\Delta f = \frac{f_{uu} + f_{vv}}{G}$ .

(iii) Let  $I : M \rightarrow \mathbb{R}^3$  be the inclusion map of a surface  $M$  into  $\mathbb{R}^3$ , and let  $x_i = u_i \circ I$  for  $i = 1, 2, 3$ . Defining  $\Delta I = (\Delta x_1, \Delta x_2, \Delta x_3)$ , show that  $\Delta I = -HN$  on  $B$ . Thus, if  $M$  is minimal, then the functions  $x_i$  are harmonic on  $B$ .

90. Let  $f_1, f_2, f_3$  be three analytic functions defined on an open set  $B \subset \mathbb{C}$ . Let  $Z : B \rightarrow \mathbb{C}^3$  by  $Z(w) = (f_1(w), f_2(w), f_3(w))$  and define

$X$  and  $Y$  mapping  $B$  into  $\mathbb{R}^3$  by  $X = \operatorname{Re}(Z)$  and  $Y = \operatorname{Im}(Z)$  so  $Z = X + iY$ . If  $Z' \circ Z' = 0$  and  $X_u \circ X_u > 0$  on  $B$ , show the maps  $X$  and  $Y$  each define an immersion of  $B$  into  $\mathbb{R}^3$  whose image locally is a minimal surface. Conversely, if  $M$  is a minimal surface in  $\mathbb{R}^3$  and  $m \in M$ , show that there is an open set  $B \subset \mathbb{C}$  and analytic functions  $f_1, f_2, f_3$  defined on  $B$  such that  $X(B)$  is a neighbourhood of  $m$ .

91. A Weingarten surface is a surface whose principal curvatures are functionally independent. Let  $W : M \rightarrow \mathbb{R}^2$  by  $W(m) = (k(m), h(m))$ , where  $k \leq h$ , and call the image of  $W$  the  $W$ -diagram.

(i) Show there exists no compact Weingarten surface of positive Gauss curvature whose  $W$ -diagram has negative slope (see section 3.1).

(ii) Show a compact surface with  $K > 0$  and  $H$  constant is a sphere.

Hopf [Hop50] has shown a compact surface with (a) constant mean curvature and (b) Euler characteristic zero, is a sphere. It is an open question whether the assumption (b) can be dropped.<sup>1</sup>

92. Let  $X$  and  $Y$  be the coordinate fields for a set of orthogonal coordinates on a surface. Show that there exist conformal coordinate with the same coordinate curves (as images) iff  $YX \left[ \log \left( \frac{E}{G} \right) \right] = 0$ .

---

<sup>1</sup>This conjecture was disproven, first by [Hsi82] and next in  $\mathbb{R}^3$  by [Wen86].



## 10. Topics in Riemannian Geometry

### 10.1 Jacobi Fields and Conjugate Points

In order to study the minimizing properties of geodesics, we study one and two parameter families of curves and the vector fields which they induce. Our main tools are developed in the following three propositions.

Let  $Q$  and  $M$  be  $C^\infty$  manifolds, and let  $f$  be a  $C^\infty$  map of  $Q$  into  $M$ . A  $\mathbf{TM}$ -valued vector field on  $Q$  associated with  $f$ , or a  $\mathbf{TM}_f$  field on  $Q$ , is a  $C^\infty$  function  $A$  from  $Q$  into  $\mathbf{TM}$ , the tangent bundle to  $M$ , such that  $A(p)$  lies in  $\mathbf{T}_{f(p)}M$  for all  $p$  in  $Q$ . The field  $A$  is a tangent  $\mathbf{TM}_f$  field on  $Q$  if  $A = f_*A'$  for some  $C^\infty$  field  $A'$  on  $Q$ .

For the remainder of this section, let  $Q$ ,  $M$  and  $f$  be as just described, and let  $\mathbf{D}$  denote a connexion on  $M$ .

If  $A$  and  $Z$  are  $\mathbf{TM}_f$  fields on  $Q$  and  $A = f_*A'$  is tangent, then we can define  $\mathbf{D}_A Z$  to be a  $\mathbf{TM}_f$  field on  $Q$ . This is possible, since for a particular  $p$  in  $Q$  the field  $Z$  gives a well-defined  $C^\infty$  field along a curve through  $f(p)$  with tangent  $A_p$ . More explicitly, let  $y_1, \dots, y_n$  be a coordinate system on  $M$  about  $f(p)$  and let  $Y_i = \frac{\partial}{\partial y_i}$ . Let  $U$  be an open set about  $p$  such that  $f(U)$  is contained in the domain of the  $y_i$ . Then  $Z = \sum_1^n z_i Y_i$  defines real valued  $C^\infty$  functions  $z_i$  on  $U$  and

$$\mathbf{D}_A Z = \sum_{i=1}^n [(A' z_i) Y_i + z_i (\mathbf{D}_A Y_i)] \quad (1)$$

on  $U$ . Letting equation (1) define  $\mathbf{D}_A Z$  on  $U$ , we leave it to the reader to show this definition is independent of the coordinate system. Notice that  $\mathbf{D}_A Z$  is not necessarily a tangent  $\mathbf{T}M_f$  field even when both  $A$  and  $Z$  are tangent.

If  $A$  and  $B$  are tangent  $\mathbf{T}M_f$  fields on  $Q$ , then we define the tangent  $\mathbf{T}M_f$  field  $[A, B]$  by  $[A, B](p) = f_*([A', B']_p)$  where  $A = f_*A'$ ,  $B = f_*B'$  and  $p$  in  $Q$ .

**Proposition 10.1.** Let  $A, B, X, Z$  be  $\mathbf{T}M_f$  fields on  $Q$ , let  $A$  and  $B$  be tangent, and let  $g$  be a real-valued  $C^\infty$  function on  $Q$ . Then the following equations are valid:

$$\mathbf{D}_{(gA)}X = g(\mathbf{D}_A X), \quad (2)$$

$$\mathbf{D}_A(gX) = (A'g)X + g(\mathbf{D}_A X), \quad (3)$$

$$\mathbf{D}_{(A+B)}X = \mathbf{D}_A X + \mathbf{D}_B X, \quad (4)$$

$$\mathbf{D}_A(X + Z) = \mathbf{D}_A X + \mathbf{D}_A Z. \quad (5)$$

*Proof.* All four equations follow in a straightforward way from the definition (1) and the standard properties for  $\mathbf{D}$ .  $\square$

Observe now for  $\mathbf{T}M_f$  fields  $X$  and  $Z$  we can define the  $\mathbf{T}M_f$  field  $\text{Tor}(X, Z)$  by  $[\text{Tor}(X, Z)](p) = \text{Tor}(X_p, Z_p)$  since  $\text{Tor}$  is a tensor; moreover, the linear transformation-valued tensor  $[R(X, Z)](p) = R(X_p, Z_p)$  is defined by  $p \in Q$ .

**Proposition 10.2.** With the hypothesis of Proposition (10.1), the following equations are valid:

$$\text{Tor}(A, B) = \mathbf{D}_A B - \mathbf{D}_B A - [A, B], \quad (6)$$

$$R(A, B)X = \mathbf{D}_A \mathbf{D}_B X - \mathbf{D}_B \mathbf{D}_A X - \mathbf{D}_{[A, B]}X. \quad (7)$$

*Proof.* Using the notation developed above for equation (1), let

$A = \sum_1^n a_i Y_i$  and  $B = \sum_1^n b_j Y_j$ . Then on  $U$ ,

$$\text{Tor}(A, B) = \sum_{i,j} a_i b_j \text{Tor}(Y_i, Y_j) = \sum_{i,j} a_i b_j (\mathbf{D}_{Y_i} Y_j - \mathbf{D}_{Y_j} Y_i),$$

but

$$[A, B] = \sum_j (A' b_j) Y_j - \sum_i (B' a_i) Y_i$$

and

$$\mathbf{D}_A B - \mathbf{D}_B A = \sum_j (A' b_j) Y_j + \sum_{i,j} b_j a_i \mathbf{D}_{Y_i} Y_j - \sum_i (B' a_i) Y_i - \sum_{i,j} a_i b_j \mathbf{D}_{Y_j} Y_i;$$

hence, equation (6) follows.

A similar computation gives (7). □

**Proposition 10.3.** If  $M$  is a Riemannian manifold and  $\mathbf{D}$  is the Riemannian connexion, then with the hypothesis of Proposition (10.1), the following equations are valid:

$$A' \langle X, Z \rangle = \langle \mathbf{D}_A X, Z \rangle + \langle X, \mathbf{D}_A Z \rangle, \tag{8}$$

$$\text{Tor}(X, Z) = 0. \tag{9}$$

*Proof.* Since  $\text{Tor} = 0$  in this case, equation (9) is trivial.

To verify (8), let  $Y_1, \dots, Y_n$  be an orthonormal base field with no loss of generality. Letting  $X = \sum_1^n x_i Y_i$  and  $Z = \sum_1^n z_j Y_j$ , we have

$$A' \langle X, Z \rangle = A' \left( \sum_1^n x_i z_i \right) = \sum_1^n [(A' x_i) z_i + x_i (A' z_i)],$$

while

$$\langle \mathbf{D}_A X, Z \rangle + \langle X, \mathbf{D}_A Z \rangle = \sum_i (A' x_i) z_i + \sum_{i,j} x_i z_j \langle \mathbf{D}_A Y_i, Y_j \rangle + \sum_j x_j (A' z_j) + \sum_{i,j} x_i z_j \langle Y_i, \mathbf{D}_A Y_j \rangle.$$

But  $\langle \mathbf{D}_A Y_i, Y_j \rangle + \langle Y_i, \mathbf{D}_A Y_j \rangle = A \langle Y_i, Y_j \rangle = 0$ ; hence (8) follows.  $\square$

We specialize and let  $Q$  be an open set in  $\mathbb{R}^2$ . For convenience, let  $t$  and  $w$  be the first and second coordinate functions, respectively, on  $\mathbb{R}^2$ ; then

$$T = f_* \left( \frac{\partial}{\partial t} \right) \text{ and } W = f_* \left( \frac{\partial}{\partial w} \right)$$

are tangent  $\mathbf{T}M_f$  fields on  $Q$ . Moreover, assume the  $t$ -varying curves obtained from  $f$  by holding  $w$  constant are geodesics with respect to a connexion  $\mathbf{D}$  on  $M$ ; thus  $\mathbf{D}_T T \equiv 0$  on  $Q$ . When  $f$  and  $Q$  satisfy the conditions of the above three sentences, we call  $f$  a *one-parameter family of geodesics*. When we only assume  $Q$  is an open subset of  $\mathbb{R}^2$ , we call  $f$  a *one-parameter family of curves*.

**Theorem 10.4.** If  $f$  is a one-parameter family of geodesics on  $Q$  and  $\mathbf{D}$  is torsion free, then  $\mathbf{D}_T^2 W = R(T, W)T$  on  $Q$ .

*Proof.* Since  $[T, W] = 0$  and  $\text{Tor} = 0$ , we have  $\mathbf{D}_T W = \mathbf{D}_W T$ . Hence

$$\mathbf{D}_T^2 W = \mathbf{D}_T(\mathbf{D}_T W) = \mathbf{D}_T(\mathbf{D}_W T) = \mathbf{D}_W(\mathbf{D}_T T) + R(T, W)T = R(T, W)T$$

by (6) and (7) and the fact that  $\mathbf{D}_T T = 0$ .  $\square$

Let  $T$  be the tangent field along a geodesic for a torsion-free connexion  $\mathbf{D}$  on  $M$ . Then a  $C^\infty$  field  $Z$  along the geodesic is a *Jacobi field* if  $\mathbf{D}_T^2 Z = R(T, Z)T$ . Notice the set of Jacobi fields along a geodesic is a vector space over the real field from the linearity of the defining condition.

**Theorem 10.5.** A Jacobi field  $Z$  along a geodesic is uniquely determined by its value and the value of  $\mathbf{D}_T Z$  at one point on the geodesic.

*Proof.* Let  $e_1, \dots, e_n = T$  be a parallel base along the geodesic so  $Z(t) = \sum_{i=1}^n z_i(t)e_i$  where  $t$  is the parameter on the geodesic and  $z_i$  are

$C^\infty$  real-valued functions. Then

$$D_T Z = \sum_i z'_i e_i \text{ and } D_T^2 Z = \sum_i z''_i e_i.$$

Letting  $R(e_i, e_j)e_r = \sum_{k=1}^n R_{ijrk} e_k$ , we have

$$R(T, Z)T = R(e_n, \sum z_j e_j)e_n = \sum_{j,k} z_j R_{njnk} e_k.$$

Hence  $Z$  is a Jacobi field iff  $z''_k = \sum_{j=1}^n z_j R_{njnk}$  for all  $k$ . The conclusion of the theorem now follows from the uniqueness theorem for solutions of second-order differential equations.  $\square$

**Corollary 10.6.** The vector space of Jacobi fields along a geodesic has finite dimension equal to  $2n$ . The subspace of Jacobi fields along a geodesic that vanish at a fixed point has dimension  $n$ .

The two theorems above indicate two ways of obtaining Jacobi fields, e.g., use Theorem 10.5 and existence theory from differential equations or use Theorem 10.4 by finding a one-parameter family of geodesics. We now illustrate the latter procedure.

We first fix some notation. For any vector  $A$  in the tangent space  $\mathbf{T}_m M$  we let  $A'$  be the naturally associated “constant” vector field on  $\mathbf{T}_m M$ . We use the notation of section 9.3, for a point  $X$  in  $\mathbf{T}_m M$ ,  $A'_X = \eta_X(A)$ ; or if  $e_1, \dots, e_n$  is a base of  $\mathbf{T}_m M$  and  $w_1, \dots, w_n$  its dual base with  $A = \sum_1^n a_i e_i$ , then  $A' = \sum_1^n a_i \left( \frac{\partial}{\partial w_i} \right)$ .

**Theorem 10.7.** Let  $X$  and  $A$  be any vectors in  $\mathbf{T}_m M$ . Let

$$Q = \{(t, w) \in \mathbb{R}^2 : \exp_m \text{ is defined on } t(X + wA)\},$$

which is an open set in  $\mathbb{R}^2$ . Let  $f : Q \rightarrow M$  be defined by  $f(t, w) = \exp_m t(X + wA)$ . Then  $f$  is a one-parameter family of geodesics and

$(\exp_m)_*(tA')$  is a Jacobi field along each geodesic.

*Proof.* That  $f$  is a one-parameter family of geodesics follows from the definition of the exponential map, i.e.,  $\exp_m$  maps rays in  $\mathbf{T}_mM$  into geodesics emanating from  $m$ . Then  $W = (\exp_m)_*(tA')$  is a Jacobi field by Theorem 10.4 (see Fig. 10.1).  $\square$




Figure 10.1: Jacobi Field

A point  $X$  in  $\mathbf{T}_mM$  is a *conjugate point* if  $\exp_m$  is singular at  $X$ . The point  $(m, X)$  in  $\mathbf{T}M$  is called a *conjugate point* if  $X$  is a conjugate point in  $\mathbf{T}_mM$ . A point  $m$  in  $M$  is *conjugate* to a point  $p$  in  $M$  along a geodesic  $g$  if there is a conjugate point  $X$  in  $\mathbf{T}_mM$  such that  $\exp_m X = p$  and  $g$  is a reparametrization of the geodesic  $g_X(t) = \exp_m tX$ .

Notice there is always a neighborhood of zero in  $\mathbf{T}_mM$  that is free of conjugate points since  $(\exp_m)_*$  is non-singular at zero (9.7). For a trivial (and too special) example of conjugate points, let  $M$  be the unit sphere about the origin in  $\mathbb{R}^3$ . Then the south pole is conjugate to the north pole along any geodesic (great circle); moreover, the north pole is conjugate to itself along any geodesic. To see this let  $p$  be the north pole, then  $\exp_p$  is completely singular on circles about zero in  $\mathbf{T}_mM$  which have radius  $k\pi$  for integral  $k$ .

**Theorem 10.8.** A point  $X$  in  $\mathbf{T}_m M$  is a conjugate point iff there is a non-trivial Jacobi field along  $g_X$  that vanishes at  $m$  and  $\exp_m X$ .

*Proof.* If  $\exp_m$  is singular at  $X$  let  $A' \neq 0$  be a vector such that  $(\exp_m)_* A' = 0$ . Then, letting  $A'$  denote the associated constant vector field on  $\mathbf{T}_m M$ , the field  $(\exp_m)_* tA'$  is a non-trivial Jacobi field along  $g_X$  that vanishes at  $m(t=0)$  and  $\exp_m X(t=1)$ .

Conversely, let  $Z$  be a non-trivial Jacobi field along  $g_X$  with  $Z(0) = Z(1) = 0$ . Let  $A = \mathbf{D}_X Z$  in  $\mathbf{T}_m M$  and let  $A'$  be the associated constant field on  $\mathbf{T}_m M$ . Let  $Z' = (\exp_m)_*(tA')$ . Then

$$\mathbf{D}_X Z' = \mathbf{D}_X [t(\exp_m)_* A'] = (\exp_m)_* A' + t\mathbf{D}_X [(\exp_m)_* A'],$$

and at  $t = 0$ ,  $\mathbf{D}_X Z' = A$  since at zero  $(\exp_m)_* A'_0 = A$ . Thus by uniqueness (Theorem 10.5)  $Z = Z'$ , and hence  $Z'(1) = (\exp_m)_* A'_X = 0$ . Since  $Z$  is non-trivial  $A' \neq 0$  and thus  $\exp_m$  is singular at  $X$ .  $\square$

**Corollary 10.9.** A point  $m$  is conjugate to a point  $p$  along a geodesic  $g$  iff  $p$  is conjugate to  $m$  along  $g$ .

**Theorem 10.10.** Let  $g$  be a geodesic whose parameter domain includes  $[b, c]$  and suppose  $g(b)$  is not conjugate to  $g(c)$  along  $g$ . Then there is a unique Jacobi field  $Z$  along  $g$  with prescribed values at  $g(b)$  and  $g(c)$ .

*Proof.* Suppose  $Z(b)$  and  $Z(c)$  are given. By hypothesis, the map  $\exp_{g(b)}$  is non-singular at the point  $X$  in  $\mathbf{T}_{g(b)} M$  where  $\exp_{g(b)} X = g(c)$ , i.e.,

$$g(t) = \exp_{g(b)} \left( \frac{t-b}{c-b} \right) X;$$

hence there is a unique vector  $A'$  such that  $(\exp_{g(b)})_* A' = Z(c)$ . Let  $Z_1 = (\exp_{g(b)})_*(tA')$  along  $\exp_{g(b)} tX$  (which is along  $g$ ). Similarly, we get a unique vector  $B'$  tangent to  $\mathbf{T}_{g(c)} M$  such that  $(\exp_{g(c)})_* B' = Z(b)$ . Let  $Z_2 = (\exp_{g(c)})_*(tB')$  along  $\exp_{g(c)} tY$  where  $\exp_{g(c)} Y = g(b)$ . Then  $Z = Z_1 + Z_2$  is a Jacobi field along  $g$  with the required values at  $g(b)$  and  $g(c)$ . Furthermore  $Z$  is unique, for if  $W$  were another such

field, then  $Z - W$  would be a Jacobi field that vanishes at  $g(b)$  and  $g(c)$  and hence must be trivial, so  $Z = W$ .  $\square$

## 10.2 First and Second Variation Formulae

Throughout this section let  $M$  be a  $C^\infty$  Riemannian  $n$ -manifold which is Hausdorff, and let  $\mathbf{D}$  be the Riemannian connexion. For an alternate approach to the material of this section see [Amb60].

**Theorem 10.11.** Let  $f$  be a one-parameter family of geodesics in  $M$  which are parameterized by arc length. Then  $\langle W, T \rangle$  is constant along each geodesic.

*Proof.* The function  $\langle T, T \rangle = 1$  on the domain of  $f$ ; hence,  $0 = W' \langle T, T \rangle = 2 \langle \mathbf{D}_W T, T \rangle$  by proposition 10.3. Thus

$$T \langle W, T \rangle = \langle \mathbf{D}_T W, T \rangle + \langle W, \mathbf{D}_T T \rangle = \langle \mathbf{D}_W T, T \rangle = 0,$$

since  $\mathbf{D}_T T = 0$ .  $\square$

**Corollary 10.12** (“perpendicular lemma”). Let  $X$  be a unit vector in  $\mathbf{T}_m M$ . Let  $A$  be in  $\mathbf{T}_m M$  with  $\langle A, X \rangle = 0$  and let  $A'$  be the associated constant vector field on  $\mathbf{T}_m M$ . Then  $(\exp_m)_* A'$  is perpendicular to the geodesic  $g_X$  at all points where  $g_X$  is defined.

*Proof.* We may assume  $A$  is a unit vector and then define

$$f(t, w) = \exp_m t[(\cos w)X + (\sin w)A]$$

for  $t$  in the domain of  $g_X$  and  $w$  in an interval about zero. Then  $f$  is a one-parameter family of geodesics which are parameterized by arc length. Applying Theorem 10.11, we have  $\langle W, T \rangle$  constant along each geodesic. In this case

$$W = (\exp_m)_* t[-(\sin w)X + (\cos w)A]$$



and  $w = 0$  along  $g_X$ ; hence,

$$\langle (\exp_m)_* tA, T \rangle = t \langle (\exp_m)_* A, T \rangle = \text{constant along } g_X.$$

This vanishes at  $t = 0$ , so  $\langle (\exp_m)_* A, T \rangle = 0$  along  $g_X$ . □

Let  $f$  be a one-parameter family of curves with domain  $Q$  and assume  $Q$  contains the set  $(t, 0)$  for  $0 \leq t \leq b$ . Let  $f_w(t) = f(t, w)$  for  $(t, w)$  in  $Q$ , and let  $L(w)$  be the length of the curve  $f_w$  on  $[0, b]$ , i.e.,  $L(w) = \int_0^b \sqrt{\langle T, T \rangle} dt$ . We define the first and second variations of  $L$  in the direction  $f$  to be the numbers  $L'(0)$  and  $L''(0)$ , respectively, where  $L' = dL/dw$ . Actually, we should call  $L'(0)$  the “first derivative of  $L$  in the direction of the variation  $f$  evaluated at  $f_0$  on  $[0, b]$ ,” and a similar statement should be made for the “second variation.” Henceforth we refer to  $f_0$  as the base curve.

**Theorem 10.13.** In terms of the notation just developed,

$$L'(0) = \langle W, T \rangle \Big|_{(0,0)}^{(b,0)} - \int_0^b \langle W, \mathbf{D}_T T \rangle_{w=0} dt$$

when  $f_0$  is parameterized by arc length. Thus if  $f_0$  is a geodesic, then

$$L'(0) = \langle W, T \rangle \Big|_{(0,0)}^{(b,0)}.$$

*Proof.* We compute,

$$L'(w) = \int_0^b \frac{\partial}{\partial w} \sqrt{\langle T, T \rangle} dt = \int_0^b \langle T, T \rangle^{-1/2} \langle \mathbf{D}_W T, T \rangle dt.$$

When  $w = 0$ ,  $\langle T, T \rangle = 1$  and

$$\langle \mathbf{D}_W T, T \rangle = \langle \mathbf{D}_T W, T \rangle = \frac{d}{dt} \langle W, T \rangle - \langle W, \mathbf{D}_T T \rangle$$

which we integrate to obtain the above formula. □

Notice that theorem 10.13 shows  $L'(0)$  only depends on the vector field  $W$  along the base curve  $f_0$  and we may use the general

formula of theorem 10.13 to define the first variation of  $L$  in the direction of the field  $W$  where  $W$  is any  $C^\infty$  field on the base curve. For each such  $C^\infty$  field  $W$  on a base curve  $\sigma$  we can define a one-parameter family  $f$  such that  $W = f_*\left(\frac{\partial}{\partial w}\right)$  by letting  $f(t, w) = \exp_{\sigma(t)}(wW_{\sigma(t)})$ .

A curve  $\sigma$  between points  $p$  and  $q$  in  $M$  is called an extremal to the fixed end-point problem if  $L'(0) = 0$  for every one-parameter family of curves  $f$  such that  $f_0 = \sigma$  on  $[0, b]$  and  $f(0, w) = p$ , while  $f(b, w) = q$  for  $w$  near 0.

**Theorem 10.14.** A curve  $\sigma$  between points  $p$  and  $q$  in  $M$  is an extremal iff it is a geodesic.

*Proof.* If  $\sigma$  is a geodesic and the end-points are fixed so  $W = 0$  at  $p$  and  $q$ , then  $L'(0) = 0$  by theorem 10.13.

Conversely, if  $L'(0) = 0$  and  $W = 0$  at  $p$  and  $q$ , then  $\int_0^b \langle W, \mathbf{D}_T T \rangle dt = 0$  for all  $W$  belonging to admissible (fixed end-point) one-parameter variations  $f$  of  $\sigma$ . If at some point  $m$  on  $\sigma$  between  $p$  and  $q$  we suppose  $\mathbf{T}_m(\mathbf{D}_T T) \neq 0$ , then let  $W = h\mathbf{D}_T T$  where  $h$  is a  $C^\infty$  “bump” function such that  $h(m) = 1$ ,  $h \geq 0$ , and  $h = 0$  outside a neighborhood of  $m$  on which  $\mathbf{D}_T T$  doesn’t vanish. By the remarks after theorem 10.13, there is a one-parameter family  $f$  belonging to  $W$ . In this case  $\langle W, \mathbf{D}_T T \rangle = h \langle \mathbf{D}_T T, \mathbf{D}_T T \rangle \geq 0$  is a non-negative function which is non-zero on a neighborhood of  $t'$  where  $\sigma(t') = m$ , hence  $\int_0^b \langle W, \mathbf{D}_T T \rangle dt > 0$ , which is a contradiction. Thus  $\mathbf{D}_T T = 0$ , and  $\sigma$  is a geodesic.  $\square$

**Theorem 10.15.** For a point  $m$  in  $M$ , let  $r > 0$  be chosen so  $\exp_m$  maps the set  $\hat{B} = \{X \in \mathbf{T}_m M : |X| < r\}$  diffeomorphically onto its image  $B$ . Then  $B$  is the metric ball  $B(m, r) = \{p \in M : d(m, p) < r\}$ . Furthermore, if  $X$  in  $\hat{B}$  and  $p = \exp_m X$  then  $d(m, p) = |X|$ , and the geodesic  $g_X(t) = \exp_m tX$ , defined on  $[0, 1]$ , realizes the absolute minimum possible curve-length from  $m$  to  $p$ .

*Proof.* If  $T$  is the tangent to  $g_X$ , then  $\langle T, T \rangle$  is constant on  $g_X$  so

$|g_X|_0^1 = |X|$ . We must show any other broken  $C^\infty$  curve  $\sigma$  from  $m$  to  $p$  has a length which is greater than or equal to  $|X|$ , and the theorem will follow.

First suppose  $\sigma$  is defined on  $[0, b]$  and  $\sigma(t)$  is in  $B$  for all  $t$  in  $[0, b]$ . Furthermore, suppose  $\sigma$  never returns to  $m$  after  $t = 0$ , or we could obviously obtain a shorter curve from  $m$  to  $p$ . Let:

- $\exp = \exp_m$ ,
- $\exp^{-1}$  be the inverse map of  $\exp|_{\hat{B}}$ ,
- $f(t) = |\exp^{-1} \sigma(t)|$  for  $t$  in  $[0, b]$ , which defines a broken  $C^\infty$  function  $f$ ,
- $\bar{\sigma}(t) = \exp^{-1} \sigma(t)$ ,
- $\bar{\gamma}(t) = f(t) \frac{X}{|X|}$ ,
- and  $\gamma(t) = \exp \bar{\gamma}(t)$ .

Thus  $\gamma$  is a reparameterization of  $g_X$  which has the same “radial velocity” as  $\sigma$ . Decompose the tangent to  $\bar{\sigma}$  into a radial component  $A$  and a vector  $V$  which is orthogonal to  $A$ , thus  $T_{\bar{\sigma}} = A + V$  on  $[0, b]$ , (actually,  $A(t) = f'(t)\bar{\sigma}(t)/f(t)$  for  $t > 0$ ). Using the [perpendicular lemma](#) proved above, we know  $\exp_* A$  is perpendicular to  $\exp_* V$ , so

$$|T_\sigma| = |\exp_* A + \exp_* V| \geq |\exp_* A| = |T_\gamma|.$$

Hence,  $|\sigma|_0^b \geq |\gamma|_0^b$ . Since  $\gamma$  is a reparameterization of  $g_X$ , we have  $|\gamma|_0^b \geq |g_X|_0^1 = |X|$ , where the inequality is strict if  $f$  is not an increasing function. Thus,  $|\sigma|_0^b \geq |X|$ .

If  $\sigma(t)$  is not in  $B$  for all  $t$ , then  $|\sigma| > r > |X|$  by the above paragraph. Hence,  $|X| = d(m, p)$  for  $X$  in  $\hat{(B)}$ , and the geodesic  $g_X$  realizes this minimum. □

**Theorem 10.16.** Let  $f$  be a one-parameter family of curves such that the base curve is a geodesic  $g$  parameterized by arc length

on the interval  $[0, b]$ . Then  $L''(0)$  is equal to

$$\langle \mathbf{D}_W W, T \rangle \Big|_{(0,0)}^{(b,0)} + \int_0^b [\langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle - (T \langle W, T \rangle)^2] dt.$$

If  $\langle W, T \rangle$  is constant along  $f$ , then

$$L''(0) = \langle \mathbf{D}_W W, T \rangle \Big|_{(0,0)}^{(b,0)} + \int_0^b [\langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle] dt.$$

If  $W$  is a Jacobi field and  $\langle W, T \rangle$  is constant along  $g$ , then

$$L''(0) = W \langle T, W \rangle \Big|_{(0,0)}^{(b,0)}.$$

**Proof.** First compute

$$\frac{\partial}{\partial w} \sqrt{\langle T, T \rangle} = \langle T, T \rangle^{-1/2} \langle \mathbf{D}_W T, T \rangle.$$

Then

$$\frac{\partial^2}{\partial w^2} \sqrt{\langle T, T \rangle} = -\langle T, T \rangle^{-3/2} \langle \mathbf{D}_W T, T \rangle^2 + \langle T, T \rangle^{-1/2} (\langle \mathbf{D}_W \mathbf{D}_W T, T \rangle + \langle \mathbf{D}_W T, \mathbf{D}_W T \rangle).$$

Evaluating on  $w = 0$ , we use  $\langle T, T \rangle = 1$ ,  $\mathbf{D}_T T = 0$ , and  $\mathbf{D}_T W = \mathbf{D}_W T$ , to obtain

$$\begin{aligned} \frac{\partial^2}{\partial w^2} \sqrt{\langle T, T \rangle} &= \langle \mathbf{D}_W \mathbf{D}_T T, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle - \langle \mathbf{D}_T W, T \rangle^2 \\ &= \langle R(W, T)W + \mathbf{D}_T \mathbf{D}_W W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle - (T \langle W, T \rangle)^2 \\ &= T \langle \mathbf{D}_W W, T \rangle + \langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle - (T \langle W, T \rangle)^2, \end{aligned}$$

which gives the first formula for  $L''(0)$  by integrating.

If  $\langle W, T \rangle$  is constant along  $g$ , then  $T \langle W, T \rangle = 0$  which gives the second formula.

If  $W$  is Jacobi, then

$$\begin{aligned}\langle R(W, T)W, T \rangle &= \langle R(T, W)W, W \rangle = \langle \mathbf{D}_T^2 W, W \rangle \\ &= T \langle \mathbf{D}_T W, W \rangle - \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle.\end{aligned}$$

Hence,  $L''(0) = \left[ \langle \mathbf{D}_W W, T \rangle + \langle \mathbf{D}_W T, W \rangle \right]_{(0,0)}^{(b,0)} = W \langle W, T \rangle \Big|_{(0,0)}^{(b,0)}$ .  $\square$

Notice that the first term is the only term in the above formulae that depends on something more than the vector field  $W$  along  $g$ .

**Corollary 10.17.** If  $W$  vanishes at the end-points of  $g$ , then the second variation of  $L$  depends only on the field  $W$  along  $g$ . For any vector field  $W$  along  $g$ , let  $f_W(t, w) = \exp_{g(t)} wW$  be the natural one-parameter family associated with  $W$ , and then  $\mathbf{D}_W W = 0$ , since the  $w$ -varying curves are geodesics. Letting  $L''_W(0)$  denote the second variation of  $L$  in the direction  $f_W$ , then

$$L''_W(0) = \int_0^b \left[ \langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle - \langle T \langle W, T \rangle^2 \rangle \right] dt.$$

We next prove two lemmas which are used to prove that geodesics are not minimizing-distance curves past a first conjugate point, and later, to prove conjugate points are isolated along a geodesic in the Riemannian case.

**Lemma 10.18 (Lagrange identity).** If  $X$  and  $Y$  are Jacobi fields along a geodesic  $g$  with tangent field  $T$ , then  $\langle \mathbf{D}_T X, Y \rangle - \langle X, \mathbf{D}_T Y \rangle$  is constant along  $g$ .

*Proof.* We compute

$$\begin{aligned}T(\langle \mathbf{D}_T X, Y \rangle - \langle X, \mathbf{D}_T Y \rangle) &= \langle \mathbf{D}_T^2 X, Y \rangle - \langle X, \mathbf{D}_T^2 Y \rangle \\ &= \langle R(T, X)W, Y \rangle - \langle R(T, Y), X \rangle = 0\end{aligned}$$

by the symmetry of the Riemann-Christoffel curvature tensor.  $\square$

**Lemma 10.19.** Let  $W$  be a continuous piecewise  $C^\infty$  field along the geodesic  $g$  which is parameterized on  $[0, b]$ , and let  $W(0) = 0$ . If there is no point  $g(t)$  that is conjugate to  $g(0)$  for  $t$  in  $[0, b]$ , then

$$\int_0^b [\langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle] dt$$

is greater than

$$\int_0^b [\langle R(Z, T)Z, T \rangle + \langle \mathbf{D}_T Z, \mathbf{D}_T Z \rangle] dt,$$

unless  $W = Z$ , where  $Z$  is the unique Jacobi field along  $g$  such that  $Z(0) = 0$  and  $Z(b) = W(b)$ .

*Proof.* The field  $Z$  is well-defined by theorem 10.10. Let  $Z_1, \dots, Z_n$  be a base of  $\mathbf{T}_{g(b)}M$ , and extend these vectors by theorem 10.10 to be Jacobi fields along  $g$  that vanish at  $g(0)$ . Since there is no point  $g(t)$  conjugate to  $g(0)$ , the fields  $Z_1, \dots, Z_n$  are a base of  $\mathbf{T}_{g(t)}M$  for all  $t$  in  $(0, b]$ . Using theorem 10.7, write each  $Z_i = tA_i$  where  $A_1, \dots, A_n$  are  $C^\infty$  fields that are independent on  $[0, b]$ . Setting  $W = \sum_{i=1}^n g_i A_i$ , we define continuous piecewise  $C^\infty$  functions  $g_i$  on  $[0, b]$ . Since  $g_i(0) = 0$  we may write  $g_i = t f_i$  and thus define continuous piecewise  $C^\infty$  functions  $f_i$  on  $[0, b]$  such that  $W = \sum f_i Z_i$ . Then  $Z = \sum f_i(b) Z_i$ .

Let  $\mathbf{D}_T W = A + B$  where  $A = \sum (T f_i) Z_i$  and  $B = \sum f_i \mathbf{D}_T Z_i$ . Then  $\langle \mathbf{D}_T W, \mathbf{D}_T W \rangle = \langle A, A \rangle + 2 \langle A, B \rangle + \langle B, B \rangle$ , and

$$\begin{aligned} \langle R(T, W)T, W \rangle &= \sum f_i \langle R(T, Z_i)T, W \rangle = \sum f_i \langle \mathbf{D}_T^2 Z_i, W \rangle \\ &= \sum f_i [T \langle \mathbf{D}_T Z_i, W \rangle - \langle \mathbf{D}_T Z_i, \mathbf{D}_T W \rangle] \\ &= T \langle B, W \rangle - \sum ((T f_i) \langle \mathbf{D}_T Z_i, W \rangle) - \langle B, A \rangle - \langle B, B \rangle. \end{aligned}$$

Hence,  $\langle R(T, W)T, W \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle$  is equal to

$$T \langle B, W \rangle + \langle A, A \rangle + \langle A, B \rangle - \sum ((T f_i) \langle \mathbf{D}_T Z_i, W \rangle).$$

But

$$\langle A, B \rangle - \sum (Tf_i) \langle \mathbf{D}_T Z_i, W \rangle = \sum (Tf_i) f_j [\langle Z_i, \mathbf{D}_T Z_j \rangle - \langle \mathbf{D}_T Z_i, Z_j \rangle] = 0$$

by the **Lagrange identity**, since  $Z_k(0) = 0$  for all  $k$ . Thus

$$\int_0^b [\langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle] dt = \langle B_b, W_b \rangle + \int_0^b \langle A, A \rangle dt$$

since  $W$  is continuous and  $W_0 = 0$ . Furthermore,

$$\begin{aligned} \langle B_b, W_b \rangle &= \left\langle \sum f_i(b) (\mathbf{D}_T Z_i)_b, W_b \right\rangle \\ &= \langle (\mathbf{D}_T Z)_b, Z_b \rangle \\ &= \int_0^b [\langle R(Z, T)Z, T \rangle + \langle \mathbf{D}_T Z, \mathbf{D}_T Z \rangle] dt. \end{aligned}$$

Since  $\int_0^b \langle A, A \rangle dt \geq 0$ , the inequality in the conclusion follows unless  $A = 0$ , which implies  $f_i$  are constant so  $W = Z$ . □

**Theorem 10.20.** The arc length on a geodesic  $g$  does not equal the distance in  $M$  beyond the first conjugate point; i.e., if  $g(b)$  is the first point of  $g$  that is conjugate to  $g(0)$ , and  $g$  is parameterized by arc length, then the distance  $d(g(0), g(a)) < a$  for  $a > b$ .

*Proof.* Let  $Z$  be a non-trivial Jacobi field along  $g$  which vanishes at 0 and  $b$ . Then  $\langle Z, T \rangle = 0$  by theorem 10.11 and  $L''_Z(0) = 0$  by theorem 10.16 where  $L'$  is computed from the natural one-parameter family of curves associated with  $Z$ . By theorem 10.15 we obtain  $r > 0$ , so that the neighborhood  $B(g(b), r)$  is the diffeomorphic image of the  $r$ -ball about zero in  $\mathbf{T}_{g(b)}M$ . Choose numbers  $a$  and  $c$  such that  $0 < c < b < a$  and  $g(t)$  is in  $B(g(b), r)$  for all  $t$  in  $[c, a]$ . Thus the interval  $[c, a]$  has no pair of points that are conjugate to each other on  $g$ . Let  $Y$  be the unique Jacobi field along  $g$  with  $Y(c) = Z(c)$  and  $Y(a) = 0$ . Let  $X$  be the field on  $[0, a]$  such that  $X(t) = Z(t)$  for  $t$  in  $[0, c]$  and  $X(t) = Y(t)$  for  $t$  in  $[c, a]$ . Let  $W$  be the field on  $[0, a]$  such that

$W(t) = Z(t)$  for  $t$  in  $[0, b]$  and  $W(t) = 0$  for  $t$  in  $[b, a]$  (see Fig. 10.2).

Then

$$L''_W|_0^a = L''_Z|_0^b + L''_W|_b^a = 0 \text{ while } L''_X|_0^a = L''_W|_0^c + L''_Y|_c^a.$$

By lemma 10.19, we have  $L''_W|_c^a > L''_Y|_c^a$ , which implies  $L''_X|_0^a < L''_W|_0^a = 0$ . Hence there are broken  $C^\infty$  curves in the natural one-parameter family associated with  $X$  whose length from  $g(0)$  to  $g(a)$  is less than  $a$ .  $\square$

Actually, the arc length on a geodesic may cease to measure distance in  $M$  long before a conjugate point is reached (think of a right circular cylinder). The conjugate point is where the geodesic ceases to be a minimum-length curve among nearby curves.




Figure 10.2: Fields Along a Geodesic

**Theorem 10.21.** The conjugate points of a fixed point on a geodesic occur at isolated values of the parameter.

*Proof.* Let  $g(b)$  be any point conjugate to  $g(0)$  along the geodesic  $g$  (notice it is possible that  $g(b) = g(0)$ ). Let  $A_1, \dots, A_r$  be a base for the kernel of  $(\exp_{g(0)})_*$  at  $bT_0$  in  $\mathbf{T}_{g(0)}M$ , where  $T_t$  is the tangent to  $g$  at  $g(t)$ , and we assume  $\langle T, T \rangle = 1$ . Choose  $A_{r+1}, \dots, A_n$  so  $A_1, \dots, A_n$  are independent and let  $Z_i(t) = (\exp_{g(0)})_* tA_i$ . Then the fields  $Z_1, \dots, Z_n$  are Jacobi fields along  $g$  that vanish at 0 and are independent for all values of  $t$  except 0 and conjugate values. We show there exists an  $\epsilon > 0$  such that  $Z_1, \dots, Z_n$  are independent for  $0 < |t - b| < \epsilon$ .



## Section 10.3: Geometric Interpretation of Riemannian Curvature

---

This is done by showing  $\mathbf{D}_T Z_1, \dots, \mathbf{D}_T Z_r, Z_{r+1}, \dots, Z_n$  are independent at  $b$  and then  $Z_1/(t-b), \dots, Z_r/(t-b), Z_{r+1}, \dots, Z_n$  are independent for  $0 < |t-b| < \epsilon$ .

Since  $A_{r+1}, \dots, A_n$  are independent at  $bT_0$ , we know  $Z_{r+1}, \dots, Z_n$  are independent at  $b$ . For  $i \leq r$ ,  $(\mathbf{D}_T Z_i)_b \neq 0$ , since  $(Z_i)_b = 0$  and  $Z_i$  is non-trivial. If

$$\sum_{i=1}^r c_i (\mathbf{D}_T Z_i)_b = 0, \text{ let } W = \sum_{i=1}^r c_i Z_i.$$

Then  $W$  is a Jacobi field with  $W_b = 0$  and  $(\mathbf{D}_T W)_b = 0$ ; hence  $W = 0$ . For small  $a > 0$ , we know  $Z_1, \dots, Z_r$  are independent, and  $\sum_{i=1}^r c_i (Z_i)_a = 0$  implies  $c_i = 0$  for all  $i$ . Thus  $\mathbf{D}_T Z_1, \dots, \mathbf{D}_T Z_r$  are independent at  $b$ . We now show for  $i \leq r$  and  $j > r$ ,  $\mathbf{D}_t Z_i$  is orthogonal to  $Z_j$  at  $b$ . By the **Lagrange identity**  $\langle \mathbf{D}_T Z_i, Z_j \rangle - \langle Z_i, \mathbf{D}_T Z_j \rangle$  is constant along  $g$ . Since  $Z_i$  and  $Z_j$  vanish at 0, and  $Z_i$  vanishes at  $b$ , we have  $\langle \mathbf{D}_T Z_i, Z_j \rangle = 0$  at  $b$ . Thus  $\mathbf{D}_T Z_1, \dots, \mathbf{D}_T Z_r, Z_{r+1}, \dots, Z_n$  are independent at  $b$  and hence in some neighborhood of  $b$ . Since  $Z_i(t)/(t-b) \rightarrow (\mathbf{D}_T Z_i)_b$  as  $t \rightarrow b$ , the conclusion follows.  $\square$

### 10.3 Geometric Interpretation of Riemannian Curvature

In this section, let:

- $M$  be a Riemannian manifold.
- $g$  be a geodesic in  $M$  with unit tangent  $T$ .
- $A_0$  be a unit vector in  $\mathbf{T}_{g(0)}M$  which is orthogonal to  $T_0$ .
- $A'$  be the constant vector field on  $\mathbf{T}_{g(0)}M$  generated by  $A_0$ .
- $\exp = \exp_{g(0)}$ .
- $A = \exp_* A'$ .
- $K = \frac{\langle R(T, A)A, T \rangle}{\langle A, A \rangle}$  as a function of  $t$  along  $g$ , whenever  $A_t \neq 0$ .

We study the relationship between the Riemannian curvature  $K(t)$  of the plane section spanned by  $A_t$  and  $T_t$  and the length of the vector  $A_t$ . The field  $tA$  is used in the computation since it is a Jacobi field.

**Lemma 10.22.** If  $tA_t \neq 0$ , then

$$(1) \quad T|tA| = \frac{\langle \mathbf{D}_T tA, tA \rangle}{|tA|} = |A| + \frac{t \langle \mathbf{D}_T A, A \rangle}{|A|}$$

$$(2) \quad T^2|tA| = -|tA|K(t) + H(t) \text{ where } H(t) \geq 0.$$

$$(3) \quad |A_t| = 1 - K(0)\frac{t^2}{6} + G(t)t^3 \text{ for } t \text{ in a neighborhood of zero where } G \text{ is } C^\infty.$$

*Proof.* We compute

$$T|tA| = T\sqrt{\langle tA, tA \rangle} = \frac{\langle \mathbf{D}_T tA, tA \rangle}{|tA|} = \frac{\langle A + t\mathbf{D}_T A, tA \rangle}{|tA|} = |A| + \frac{t \langle \mathbf{D}_T A, A \rangle}{|A|}$$

Thus

$$\begin{aligned} T^2|tA| &= \frac{1}{|tA|} \left[ \langle \mathbf{D}_T^2 tA, tA \rangle + \langle \mathbf{D}_T tA, \mathbf{D}_T tA \rangle - \frac{\langle \mathbf{D}_T tA, tA \rangle^2}{\langle tA, tA \rangle} \right] \\ &= \frac{1}{|tA|^3} \left[ \langle R(T, tA)T, tA \rangle |tA|^2 + |\mathbf{D}_T tA|^2 |tA|^2 - \langle \mathbf{D}_T tA, tA \rangle^2 \right] \\ &= -|tA|K(t) + H(t) \end{aligned}$$

where

$$H(t) = \frac{1}{|tA|^3} \left[ |\mathbf{D}_T tA|^2 |tA|^2 - \langle \mathbf{D}_T tA, tA \rangle^2 \right]$$

The Schwartz inequality implies  $H(t) \geq 0$ . A straightforward computation shows, as  $t \rightarrow 0$ ,  $H(t) \rightarrow 0$ ,  $H'(t) \rightarrow 0$  since  $(\mathbf{D}_T A)_0 = 0$  (use normal coord.), hence as  $t \rightarrow 0$ ,

$$|tA| \rightarrow 0, \quad T|tA| \rightarrow |A_0| = 1, \quad T^2|tA| \rightarrow 0, \quad T^3|tA| \rightarrow -K(0)$$

Since  $A_t$  does not vanish near  $t = 0$ , the function  $|A_t|$  is  $C^\infty$  at  $0$ ,

### Section 10.3: Geometric Interpretation of Riemannian Curvature

---

and hence  $F(t) = |tA_t|$  admits a representation

$$F(t) = F(0) + F'(0)t + F''(0)\frac{t^2}{2} + F'''(0)\frac{t^3}{6} + G(t)t^4$$

for  $t$  in a neighborhood of  $0$  where  $G$  is a  $C^\infty$  function on this neighborhood. Substituting the values for the derivatives of  $F$  and cancelling a factor of  $t$  then gives (3).  $\square$

The following theorem derives its form essentially from some class notes of Ambrose.

**Theorem 10.23.** If  $K(t) \leq 0$  for  $t \in [0, b]$ , then  $|A_t| \geq |A_0| = 1$  for  $t \in [0, b]$ . Thus if  $K \leq 0$  for all plane sections at all points of  $M$ , then  $M$  has no conjugate points. If  $K(0) < 0$ , then  $|A_t| \geq 1$  for  $t$  near zero, and if  $K(0) > 0$ , then  $|A_t| \leq 1$  for  $t$  near zero.

*Proof.* Let

$$F(t) = |tA_t| - t|A_0| = |tA_t| - t$$

Then

$$F(0) = 0, \quad F'(0) = 0, \quad F''(t) = T^2 |tA_t| \geq 0 \quad \text{if} \quad K(t) \leq 0$$

Applying the Mean Value Theorem twice,

$$F(t) = F'(\bar{t})t = F''(\bar{\bar{t}})\bar{\bar{t}}t \geq 0 \quad \text{where} \quad 0 \leq \bar{\bar{t}} \leq \bar{t} \leq t \leq b$$

Hence  $|A_t| \geq 1$  for  $t \in [0, b]$ .

The second sentence of the theorem follows from the first, and the last two sentences follow from ((3)) in the lemma.  $\square$

We obtain a geometric interpretation of Riemannian curvature from the following considerations (see Fig. 10.3). The vector  $A'$  at the point  $bT_0 \in \mathbf{T}_{g(0)}M$  is tangent to the circle  $\sigma$  of radius  $b$  about the origin which lies in the plane of  $A_0$  and  $T_0$ . Hence  $A = \exp_* A'$  is the tangent at  $\exp(bT_0)$  to the curve  $\exp \circ \sigma$  in  $M$ . If  $b$  is sufficiently small, then  $\exp \circ \sigma$  passes through points that are exactly  $b$  units

distant from  $g(0)$ . If  $|A_b| > |A'|$  then the curve  $\exp \circ \sigma$  is “stretching” the curve  $\sigma$  near  $bT_0$  and the geodesics emanating from  $g(0)$  that are determined by  $\sigma$  are “spreading out”. A corresponding statement applies to the case  $|A_b| < |A'|$ .




Figure 10.3: Comparing Geodesics

## 10.4 The Morse Index Theorem

Our approach to this section is based on the notes of Bott. For further material see [JW69], [Amb61] and [Mor34]. Let  $M$  be a  $C^\infty$  manifold and let  $f$  be a real-valued  $C^\infty$  function defined on a neighborhood of a point  $m \in M$ . The point  $m$  is a *critical point* of  $f$  if  $(f_*)_m$  is the zero linear transformation on  $\mathbf{T}_m M$ . If  $m$  is a critical point of  $f$ , we define a symmetric bilinear function  $H : \mathbf{T}_m M \times \mathbf{T}_m M \rightarrow \mathbb{R}$  by

$$H(X_m, Y_m) = X_m(Yf)$$

where  $Y$  is any  $C^\infty$  vector field about  $m$  whose value at  $m$  is  $Y_m$ . It is a simple exercise to show  $H(X_m, Y_m)$  is independent of the field  $Y$  and is symmetric and bilinear (see problem 95). The function  $H$  is called the *Hessian* of  $f$  at  $m$ . The *index* of  $H$  is defined to be the dimension of a maximal subspace  $V$  of  $\mathbf{T}_m M$  on which  $H$  is

negative definite (and  $V$  is maximal if it is not properly contained in a subspace  $V'$  on which  $H$  is negative definite). The null space of  $H$  is the subspace

$$V = \{X \in \mathbf{T}_m M : H(X, Y) = 0 \text{ for all } Y \in \mathbf{T}_m M\}$$

The nullity of  $H$  is the dimension of its null space. We denote the index of  $H$  and the nullity of  $H$  by  $I(f_m)$  and  $N(f_m)$ , respectively, and call them the *index of  $f$  at  $m$*  and the *nullity of  $f$  at  $m$* , respectively. The *positivity*  $P(f_m)$  is the integer such that  $P(f_m) + I(f_m) + N(f_m)$  is the dimension of  $\mathbf{T}_m M$ . The index of  $H$  intuitively gives the number of dimensions of directions in  $\mathbf{T}_m M$  in which  $f$  is decreasing.

Next we need the definition of the conjugate degree of points along a geodesic. Let  $g$  be a geodesic in a manifold with connexion. The *conjugate degree of the point  $g(t)$*  (with respect to  $g(0)$ ) is the dimension of the kernel of  $(\exp_{g(0)})_*$  at  $tT_0$ , where  $T_0$  is the unit tangent to  $g$  at  $g(0)$  and  $g$  is parameterized by arc length. Thus the conjugate degree of the point  $g(t)$  is the maximum number of linearly independent Jacobi fields along  $g$  that vanish at  $0$  and  $t$ .

The Morse Index Theorem relates the concepts just defined. Roughly it says, for a particular geodesic segment in a Riemannian manifold  $M$ , the distance function can be used to define a  $C^\infty$  function  $L$  on a manifold  $C$ , and then the index of  $L$  at a particular critical point is equal to the sum of the degrees of conjugate points along the geodesic segment.

For the rest of the section let  $M$  be a  $C^\infty$  Riemannian Hausdorff  $n$ -manifold. If  $m \in M$ , then a *local geodesic manifold of  $M$  at  $m$*  is a submanifold  $C$  defined as follows. Let  $B$  be an open ball about the origin (zero) in  $\mathbf{T}_m M$  which  $\exp_m$  maps diffeomorphically into  $M$ , and let  $V$  be any subspace of  $\mathbf{T}_m M$ . Then the submanifold

$$C = \{\exp_m X : X \in B \wedge V\}$$

is a *local geodesic submanifold of  $M$* . Note  $C$  contains geodesic

segments of geodesics emanating from  $m$  whose tangent vectors lie in  $V$  (see Fig. 10.4).

**Lemma 10.24.** Let:

- $A$  be a convex neighborhood of  $M$ .
- $p_1, p_2 \in A$ .
- $g$  be the unique geodesic from  $p_1$  to  $p_2$  which lies in  $A$  and is parameterized by arc length.
- $T$  be the tangent field to  $g$ .
- $C_1$  and  $C_2$  be disjoint local geodesic hypersurfaces of  $A$  through  $p_1$  and  $p_2$ , respectively, that are orthogonal to  $T$ .
- $C = C_1 \times C_2$  (see Fig. 10.4).
- $d(m_1, m_2)$  be the distance from  $m_1$  to  $m_2$  whenever  $(m_1, m_2) \in C$ ; thus  $d \in C^\infty(C, R)$  (problem 96).
- $W = (W_1, W_2)$  and  $U = (U_1, U_2)$  be vectors tangent to  $C$  at  $(p_1, p_2)$ , where  $W_i, U_i \in \mathbf{T}_{p_i}M$  for  $i = 1, 2$ , and let  $U$  also denote the unique Jacobi field along  $g$  determined by  $U_1, U_2$ .

Then  $p = (p_1, p_2)$  is a critical point of  $d$  on  $C$  and

$$!!_p(U, W) = U_p(Wd) = [\langle W, \mathbf{D}_T U \rangle - !!_T(U, W)]_{p_1}^{p_2}$$

where  $!!_T$  at  $p_i$  is the second fundamental form of  $C_i$  with respect to the normal in the direction of  $T$ .

*Proof.* A two-parameter family of geodesics is a  $C^\infty$  function  $f$  mapping an open set  $Q \subset \mathbb{R}^3$  into  $M$  such that the curves

$$f_{(u_0, w_0)}(t) = f(t, u_0, w_0)$$

obtained from  $f$  by fixing the coordinates in the last two slots, are geodesics. Let  $f$  be such a map and suppose  $(t, 0, 0) \in Q$  for




Figure 10.4: Cross Manifolds

$0 \leq t \leq b$ . Call the geodesic  $g = f_{(0,0)}$  the base geodesic and assume  $g$  is parameterized by arc length. Let

$$T = f_*\left(\frac{\partial}{\partial t}\right), \quad U = f_*\left(\frac{\partial}{\partial u}\right), \quad W = f_*\left(\frac{\partial}{\partial w}\right)$$

then  $T, U, W$  are Jacobi fields along the geodesics of  $f$ , while

$$\mathbf{D}_T W = \mathbf{D}_W T, \quad \mathbf{D}_T U = \mathbf{D}_U T, \quad \mathbf{D}_U W = \mathbf{D}_W U$$

by section 10.1. We assume further that  $\langle T, U \rangle$  and  $\langle T, W \rangle$  are constant on  $g$ ; hence  $\langle \mathbf{D}_T U, T \rangle = 0$  and  $\langle \mathbf{D}_T W, T \rangle = 0$  on  $g$ . For  $(u, w)$  near  $(0, 0)$ , let

$$L(u, w) = \int_0^b \sqrt{\langle T, T \rangle} dt.$$

Notice  $\langle T, T \rangle$  is a function on  $Q$  which depends only on  $u, w$  since the  $t$ -curves are geodesics. Differentiating with respect to  $w$ ,

$$L_w = \frac{\partial L}{\partial w} = \int_0^b \langle T, T \rangle^{-1/2} \langle \mathbf{D}_W T, T \rangle dt$$

and

$$(L_w)_{(0,0)} = \int_0^b \langle \mathbf{D}_T W, T \rangle dt = 0$$

since  $\langle T, T \rangle = 1$  on  $g$ . Differentiating with respect to  $u$ ,

$$L_{wu} = \int_0^b [-\langle T, T \rangle^{-3/2} \langle \mathbf{D}_U T, T \rangle \langle \mathbf{D}_W T, T \rangle + \langle T, T \rangle^{-1/2} (\langle \mathbf{D}_U \mathbf{D}_W T, T \rangle + \langle \mathbf{D}_W T, \mathbf{D}_U T \rangle)] dt$$

Evaluating on  $g$ ,

$$\begin{aligned} (L_{wu})_{(0,0)} &= \int_0^b [\langle \mathbf{D}_U \mathbf{D}_T W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T U \rangle] dt \\ &= \int_0^b [\langle R(U, T)W, T \rangle + \mathbf{D}_T \mathbf{D}_U W, T \rangle + T \langle W, \mathbf{D}_T U \rangle - \langle W, \mathbf{D}_T^2 U \rangle] dt \end{aligned}$$

But, since  $U$  is Jacobi,

$$\langle R(U, T)W, T \rangle - \langle W, \mathbf{D}_T^2 U \rangle = \langle R(U, T)W, T \rangle - \langle W, R(T, U)T \rangle = 0$$

hence

$$\begin{aligned} (L_{wu})_{(0,0)} &= \int_0^b [\langle \mathbf{D}_T \mathbf{D}_U W, T \rangle + T \langle W, \mathbf{D}_T U \rangle] dt \\ &= \int_0^b [T \langle \mathbf{D}_U W, T \rangle + T \langle W, \mathbf{D}_T U \rangle] dt \\ &= \langle \mathbf{D}_U W, T \rangle + \langle W, \mathbf{D}_T U \rangle \Big|_{(0,0,0)}^{(b,0,0)} \end{aligned}$$

We apply the above analysis to prove the lemma. Let

$$f_{(u,w)}(t) = f(t, u, w)$$

be the unique geodesic in  $A$  from

$$\exp_{p_1}(uU_1 + wW_1) = \gamma_1(u, w)$$

to

$$\exp_{p_2}(uU_2 + wW_2) = \gamma_2(u, w)$$

which is parameterized on  $[0, b]$ . Then  $f$  is a two-parameter family



of geodesics satisfying the above requirements. Furthermore,

$$d(\gamma_1(u, w), \gamma_2(u, w)) = L(u, w)$$

hence

$$H_p(U, W) = \left\langle \mathbf{D}_{U_i} W_i, T \right\rangle + \left\langle W_i, \mathbf{D}_T U_i \right\rangle \Bigg|_{i=1}^{i=2}$$

But letting  $\mathbf{D}'$  be the induced Riemannian connexion on  $C_i$ , by the Gauss equation we get

$$\mathbf{D}_{U_i} W_i = \mathbf{D}'_{U_i} W_i - \mathbb{I}_T(U_i, W_i)T$$

hence

$$\mathbb{I}_p(U, W) = [\langle W, \mathbf{D}_T U \rangle - \mathbb{I}_T(U, W)]_{p_1}^{p_2}$$

□

**Theorem 10.25. (Morse Index Theorem).** Let  $g$  be a geodesic in  $M$  which is parameterized by arc length on the interval  $[0, b]$ . Let  $r > 0$  be chosen such that the balls  $B(g(t), 2r)$  are convex neighborhoods of  $g(t)$  for  $0 \leq t \leq b$ . Let  $\bar{m} = (m_1, \dots, m_k)$  be a sequence of points on  $g$  such that  $m_i = g(t_i)$ ,  $0 < t_i < t_{i+1} < b$ , and

$$0 < d(m_i, m_{i+1}) < r \tag{10}$$

for  $0 \leq i \leq k$  where  $m_0 = g(0)$  and  $m_{k+1} = g(b)$ . Let  $C_i$  be a local geodesic submanifold which is orthogonal to  $g$  at  $m_i$  and contained in  $B(m_i, r)$  for

$1 \leq i \leq k$ , and let  $C = C_1 \times \dots \times C_k$ . Define  $L : C \rightarrow \mathbb{R}$  by

$$L(\bar{p}) = \sum_{i=0}^k d(p_i, p_{i+1})$$

where  $\bar{p} = (p_1, \dots, p_k) \in C$ ,  $p_0 = g(0)$ , and  $p_{k+1} = g(b)$  (see Fig. 10.5).

Then  $L$  is  $C^\infty$  on  $C$ ,  $\bar{m}$  is a critical point of  $L$ , the nullity of  $L$  at

$\bar{m}$  equals the conjugate degree of  $g(b)$  (with respect to  $g(0)$ ), and

$$I(L_{\bar{m}}) = \sum_{0 \leq t \leq b} \text{deg } g(t)$$




Figure 10.5: Cross Manifolds

Before proving the theorem, we make some remarks. The fact that  $N(L_{\bar{m}})$  is the conjugate degree of  $g(b)$  is often called the Nullity Theorem. The Index Theorem shows  $I(L_{\bar{m}})$  and  $N(L_{\bar{m}})$  are independent of the position of the points  $m_i$  and the number of points  $k$  as long as condition (10) is satisfied.

*Proof.* Define  $L_i : C \rightarrow \mathbb{R}$  by

$$L_i(\bar{p}) = d(p_i, p_{i+1})$$

for  $0 \leq i \leq k$ . Then  $L$  is  $C^\infty$  since  $L = \sum L_i$  and each  $L_i$  is  $C^\infty$ . By the lemma, the point  $\bar{m}$  is a critical point of each  $L_i$  and hence is a critical point of  $L$ .

To compute the nullity of  $L$  at  $\bar{m}$ , let  $U, W \in \mathbf{T}_{\bar{m}}C$  where

$$U = (U_1, \dots, U_k), \quad W = (W_1, \dots, W_k), \quad U_i, W_i \in \mathbf{T}_{m_i}M$$

Let  $U_0 = W_0$  and  $U_{k+1} = W_{k+1}$  be the zero vectors at  $g(0)$  and  $g(b)$ ,

---

respectively. By the lemma,

$$\begin{aligned}
 U_{\bar{m}}(WL) &= \sum_{i=0}^k U_{\bar{m}}(WL_i) \\
 &= \sum_{i=1}^k [\langle W_{i+1}, \mathbf{D}_T U_{i+1}^- \rangle - !I_T(U_{i+1}, W_{i+1}) - \langle W_i, \mathbf{D}_T U_i^+ \rangle + !I_T(U_i, W_i)] \\
 &= \sum_{i=1}^k \langle W_i, \mathbf{D}_T U_i^- - \mathbf{D}_T U_i^+ \rangle
 \end{aligned}$$

where  $U_i^-$  is the Jacobi field on  $[t_{i-1}, t_i]$  agreeing with  $U$  at the end points, and  $U_i^+ = U_{i+1}^-$ . If  $U$  is in the null space of  $H_L$  at  $\bar{m}$ , then  $U_{\bar{m}}(WL) = 0$  for all  $W$ ; hence  $\mathbf{D}_T U_i^- = \mathbf{D}_T U_i^+$  for all  $i$ , which implies  $U$  is a Jacobi field along  $g$  that vanishes at  $\theta$  and  $b$ . This proves the nullity theorem.

We now work on the index of  $L$  at  $\bar{m}$ . Let us refer to a point  $(\bar{m}, b) \in M^k \times \mathbb{R}$  which satisfies the conditions stated in the third sentence of the theorem as an admissible partition. Let  $N = k(n-1)$ , and for each admissible partition  $(y, t)$  let  $C_y$  be the product of  $k$  local geodesic submanifolds crossing  $g$  at the points of  $y$ , let  $L_{(y,t)} : C_y \rightarrow \mathbb{R}$  be the function corresponding to  $L$  in the theorem, and let  $F_y$  map  $\mathbb{R}^N$  into the tangent space to  $C_y$  at  $y$  by

$$F_y(a_1, \dots, a_N) = \left( \sum_{i=1}^{n-1} a_i e_i(y_1), \sum_{j=1}^{n-1} a_{n-1+j} e_j(y_2), \dots \right)$$

where  $e_1, \dots, e_{n-1}, T$  is an orthonormal parallel base field along  $g$ . Then let  $H_{(y,t)}$ ,  $I_{(y,t)}$ ,  $P_{(y,t)}$ , and  $N_{(y,t)}$  denote the Hessian, index, positivity, and nullity, respectively, of  $H_{L_{(y,t)}} \circ F_y$ . Thus  $H_{(y,t)}$  is a symmetric bilinear form on  $\mathbb{R}^N$  which is continuous in  $y$  and  $t$ .

For each admissible partition  $(y_0, t_0)$  there is a neighborhood (in  $M^k \times \mathbb{R}$ ) such that

$$I_{(y,t)} \geq I_{(y_0,t_0)}, \quad P_{(y,t)} \geq P_{(y_0,t_0)} \tag{11}$$

for  $(y, t)$  in this neighborhood. This follows since  $I_{(y_0, t_0)}$  is the dimension of a subspace  $V \subset \mathbb{R}^N$  such that  $H_{(y_0, t_0)}(W, W) < 0$  for all non-zero  $W \in V$ , and by continuity the inequality must hold on a neighborhood of  $(y_0, t_0)$ . A similar argument handles the positivity case.

Fix  $y$  such that  $(y, b_1)$  and  $(y, b_2)$  are admissible partitions with  $b_1 \leq b_2$ . We show

$$I_{(y, b_1)} \leq I_{(y, b_2)}, \quad P_{(y, b_1)} \geq P_{(y, b_2)} \quad (12)$$

For  $x$  in the cross manifold  $C_y$ , let

$$A(x) = L_{(y, b_2)}(x), \quad B(x) = L_{(y, b_1)}(x) + d(g(b_1), g(b_2))$$

Then  $A(x) \leq B(x)$  by the triangle inequality and  $A(y) = B(y)$ . On a curve  $\gamma(w)$  with tangent  $W$  that is tangent to  $C_y$  at  $y = \gamma(0)$ ,

$$A \circ \gamma(w) = A(y) + H_{(y, b_2)}(W, W) \frac{w^2}{2} + \dots$$

while

$$B \circ \gamma(w) = B(y) + H_{(y, b_1)}(W, W) \frac{w^2}{2} + \dots$$

Thus  $H_{(y, b_2)}(W, W) \leq H_{(y, b_1)}(W, W)$  for all  $W$ , and if  $H_{(y, b_1)}$  is negative definite on a subspace  $V$  then so is  $H_{(y, b_2)}$ , which implies  $I_{(y, b_1)} \leq I_{(y, b_2)}$ , and similarly,  $P_{(y, b_1)} \geq P_{(y, b_2)}$ .

If  $g(t)$  is not a conjugate point of  $g(0)$ , then  $H_{(y, t)}$  is non-singular on a neighborhood of  $(y, t)$  since the conjugate points are isolated, hence

$$I_{(y, t)} \text{ and } P_{(y, t)} \text{ are constant on a neighborhood of } (y, t). \quad (13)$$

We now use the properties (11), (12), and (13) to compute  $I(L_y)$ . Let  $a_1, \dots, a_s$  be the points on  $[0, b)$  that are conjugate to  $\mathbf{0}$ . If  $0 < t < a_1$  we know

$$P_{(y, t)} = N, \quad I_{(y, t)} = N_{(y, t)} = 0$$

by theorem 10.15 and property (13). At  $t = a_1$ ,

$$N_{(y,a_1)} = \deg g(a_1), \quad I_{(y,a_1)} = 0$$

by (11) since  $I(y, t) = 0$  for  $t < a_1$ , hence

$$P_{(y,a_1)} = N - \deg g(a_1)$$

If  $a_1 < t < a_2$  and  $t$  is near  $a_1$ , then  $P_{(y,t)} \geq P_{(y,a_1)}$  by (11) and  $P_{(y,t)} \leq P_{(y,a_1)}$  by (12), hence

$$P_{(y,t)} = N - \deg g(a_1), \quad N_{(y,t)} = 0, \quad I_{(y,t)} = \deg g(a_1)$$

The situation then remains unchanged for  $a_1 < t < a_2$  by (13). For  $t = a_2$ , we repeat the above reasoning to compute

$$N_{(y,a_2)} = \deg g(a_2), \quad I_{(y,a_2)} = \deg g(a_1), \quad P_{(y,a_2)} = N - \sum_{0 \leq t \leq a_2} \deg g(t)$$

Continuing the argument, we obtain

$$I(L_y) = I_{(y,b)} = \sum_{0 \leq t \leq b} \deg g(t)$$

□

## 10.5 Completeness

The theorem that follows gives useful criteria for a Riemannian manifold to be complete. The analytic case was first studied by Hopf-Rinow. The approach we give essentially follows de Rham [Rha52].

**Theorem 10.26.** If  $M$  is a connected Hausdorff Riemannian manifold, then statements (a) through (d) below are equivalent, and any one of them implies (e).

- (a) The exponential map is everywhere defined on  $TM$ .

- (b)  $M$  is complete with respect to its Riemannian metric.
- (c) Bounded closed sets in  $M$  are compact.
- (d) The closed balls  $\bar{B}(m, r)$  are compact for one  $m \in M$  and all  $r > 0$ .
- (e) Any two points in  $M$  can be joined by a geodesic segment whose length equals the distance between the two points.

*Proof.* The implications (d)  $\implies$  (c)  $\implies$  (b)  $\implies$  (a) are all simple. We show (a) implies (d) and (e). Fix  $m \in M$  and let

$$B_r = B(m, r), \quad S_r = \bar{B}(m, r)$$

$$E_r = \{p \in S_r : \text{there is a geodesic segment } \gamma \text{ from } m \text{ to } p \text{ with } |\gamma| = d(m, p)\}$$

We show  $E_r$  is compact and  $E_r = S_r$  for all  $r$ , which proves (d) and (e).

**Lemma 10.27.** The set  $E_r$  is compact for all  $r$ .

*Proof.* Fix  $r$  and let  $(m_k)$  be a sequence of points in  $E_r$ . By (a) there exist points  $X_k \in \mathbf{T}_m M$  such that  $\exp_m X_k = m_k$  for all  $k$ . This follows since a geodesic can always be written as a composite map which is the exponential of a ray in a tangent space. Then  $|X_k| < r$  for all  $k$ , hence  $(X_k)$  is a sequence of points in the compact set  $\bar{B}(0, r)$  in the Euclidean space  $\mathbf{T}_m M$ . Thus we obtain a subsequence (which we reindex if necessary)  $(X_k)$  that converges to  $X \in \mathbf{T}_m M$  with  $|X| \leq r$ . The corresponding subsequence  $(m_k)$  converges to  $\exp_m X$ , which lies in  $E_r$  since  $\exp_m$  is  $C^\infty$ .  $\square$

**Lemma 10.28.** If  $E_r = S_r$  for a fixed  $r$  and  $d(m, p) > r$  then there is a point  $\bar{m}$  such that  $d(m, \bar{m}) = r$  and  $d(m, p) = r + d(\bar{m}, p)$ .

*Proof.* For each integer  $k > 0$ , let  $\gamma_k$  be a broken  $C^\infty$  curve from  $m$  to  $p$  with  $|\gamma_k| < d(m, p) + \frac{1}{k}$ . Let  $m_k$  be the last point on each  $\gamma_k$  that

lies in  $S_r$ , so  $d(m, m_k) = r$ . Since  $S_r$  is compact, the sequence  $(m_k)$  has a limit point  $\bar{m}$  and  $d(m, \bar{m}) = r$ . But

$$d(m_k, p) \leq |\gamma_k|_{m_k}^p = |\gamma_k|_m^p - |\gamma_k|_m^{m_k} \leq |\gamma_k| - r < d(m, p) + \frac{1}{k} - r$$

Hence  $d(\bar{m}, p) \leq d(m, p) - r$ , and the triangle inequality proves the opposite inequality. □

**Lemma 10.29.** For  $r \geq 0$ ,  $E_r = S_r$ .

*Proof.* The proof uses a continuous induction argument on  $r$ . By definition,  $E_r \subset S_r$  for all  $r$ . For  $r = 0$ ,  $E_0 = S_0$ . If  $E_r = S_r$ , then trivially  $E_{r'} = S_{r'}$  for all  $r' < r$ . Conversely, if  $E_{r'} = S_{r'}$  for all  $r' < r$ , then  $E_r = S_r$ . This follows by taking any point  $p \in S_r$  and then choosing  $(p_k) \rightarrow p$  such that each  $p_k$  is in some  $S_{r'}$  for  $r' < r$ . Hence  $p_k \in E_{r'} \subset E_r$ , and  $E_r$  is compact, which implies the limit  $p$  is in  $E_r$ .

Finally, if  $E_r = S_r$ , then there exists  $\epsilon > 0$  such that  $E_{r+\epsilon} = S_{r+\epsilon}$ . Since  $S_r$  is compact, we obtain a number  $2\epsilon > 0$  such that, for all  $p \in S_r$ , the map  $\exp_p$  is a diffeo from  $\{X \in \mathbf{T}_p M : |X| < 2\epsilon\}$  onto  $B(p, 2\epsilon)$ . Take  $p \in S_{r+\epsilon}$ . By lemma 10.28, there is a point  $\bar{m}$  with  $d(m, \bar{m}) = r$  and

$$d(\bar{m}, p) = d(m, p) - d(m, \bar{m}) \leq r + \epsilon - r \leq \epsilon$$

Hence there is a geodesic segment  $\gamma_1$  from  $m$  to  $\bar{m}$  with  $|\gamma_1| = r$ , and a geodesic segment  $\gamma_2$  from  $\bar{m}$  to  $p$  with  $|\gamma_2| = d(\bar{m}, p)$ . Joining  $\gamma_1$  and  $\gamma_2$  gives a broken  $C^\infty$  curve  $\gamma$  from  $m$  to  $p$  with  $|\gamma| = d(m, p)$ . Parameterizing  $\gamma$  by arc length, there can be no breaks in  $\gamma$ , so  $\gamma$  is a geodesic. Thus  $p \in E_{r+\epsilon}$ . □

□

We can now prove a classical theorem which illustrates how assumptions about the Riemannian curvature can affect the topology of a manifold.

**Theorem 10.30 (of Bonnet).** If  $M$  is a complete connected Riemannian manifold with Riemannian curvature greater than or equal to some  $K > 0$ , then  $M$  is compact and its diameter is less than or equal to  $\frac{\pi}{\sqrt{K}}$ .

*Proof.* We show on every geodesic  $g$  there is a conjugate point of  $g(0)$  on  $\left[0, \frac{\pi}{\sqrt{K}}\right]$ . If  $m \in M$ , then by completeness, every point  $p \in M$  can be joined to  $m$  by a geodesic segment whose length is  $d(p, m)$ . By Theorem 10.20, this geodesic has no conjugate point of  $m$  before  $p$ , hence  $d(m, p) \leq \frac{\pi}{\sqrt{K}}$ .

Let  $g$  be a geodesic with unit tangent  $T$ ,  $g(0) = m$ , and let  $e$  be a unit parallel field along  $g$  which is orthogonal to  $T$ . Let  $W_t = (\sin \sqrt{K}t)e_t$ . Then  $W$  is orthogonal to  $T$ ,  $W$  vanishes at  $\mathbf{0}$  and  $\frac{\pi}{\sqrt{K}}$ , and

$$\begin{aligned} \mathbf{D}_T W &= (\sqrt{K} \cos \sqrt{K}t)e_t \\ L''_W(0) &= \int_0^{\pi/\sqrt{K}} [\langle R(W, T)W, T \rangle + \langle \mathbf{D}_T W, \mathbf{D}_T W \rangle] dt \\ &= \int_0^{\pi/\sqrt{K}} [-K(t) \sin^2 \sqrt{K}t + K \cos^2 \sqrt{K}t] dt \\ &\leq K \int_0^{\pi/\sqrt{K}} [\cos^2 \sqrt{K}t - \sin^2 \sqrt{K}t] dt \\ &= 0 \end{aligned}$$

where  $K(t) = \langle R(e, t)T, e \rangle$ . If the interval  $\left[0, \frac{\pi}{\sqrt{K}}\right]$  was free of conjugate points, then by lemma 10.19,

$$L''_W(0) > L''_Z(0) = 0$$

where  $Z = 0$  is the unique Jacobi field along  $g$ , which coincides with  $W$  at  $\mathbf{0}$  and  $\frac{\pi}{\sqrt{K}}$ . This contradiction proves the theorem.  $\square$

The following theorem, due to K. Nomizu and H. Ozeki, settles



the question of the existence of complete Riemannian metrics on a paracompact (or Riemannian) manifold. A Riemannian metric is bounded if the manifold is bounded with respect to the induced metric function.

**Theorem 10.31.** Let  $M$  be a connected Hausdorff  $C^\infty$  manifold. If  $G$  is any Riemannian metric on  $M$ , then there exist Riemannian metrics  $G_1$  and  $G_2$ , both conformal to  $G$ , with  $G_1$  complete and  $G_2$  bounded.

*Proof.* Since there is more than one Riemannian metric involved, write  $G_i(X, Y)$  rather than  $\langle X, Y \rangle_i$  for the metric tensor applied to a pair of vectors,  $d_i$  for the metric, and  $B_i(m, r)$  for the corresponding  $r$ -ball neighborhoods.

Using the metric  $G$ , for each  $p \in M$ , let

$$r(p) = \sup \{r : \bar{B}(p, r) \text{ is compact}\}$$

If  $r(p) = \infty$  for some  $p$ , then  $G$  is complete by theorem 10.26. Suppose  $r(p) < \infty$  for all  $p$ , and we construct  $G_1$ .

Notice  $|r(p) - r(m)| \leq d(p, m)$  for all  $p$  and  $m$ , for if  $r(p) > r(m) + d(p, m)$ , one could increase  $r(m)$ ; hence  $r(p) \leq r(m) + d(p, m)$  for all  $p$  and  $m$ , and the inequality follows. This proves  $r$  is continuous.

Since  $M$  is paracompact, it is easy to show there exists  $f \in C^\infty(M, \mathbb{R})$  with  $f(p) > \frac{1}{r(p)}$  for all  $p$ . Let

$$G_1(X, Y) = f^2(m)G(X, Y)$$

for  $X, Y \in T_m M$ , which defines a  $C^\infty$  Riemannian metric  $G_1$  on  $M$ .

That  $G_1$  is complete will follow by showing  $B(p, \frac{1}{3}) \subset B(p, \frac{r(p)}{2})$  and hence  $\bar{B}_1(p, \frac{1}{6})$  is compact. This implies every Cauchy sequence in the  $G_1$  metric must converge. To show this, take  $p \in M$  and take  $m$  such that  $d(p, m) \geq \frac{r(p)}{2}$ . Let  $\gamma$  be a broken  $C^\infty$  curve from  $p$  to  $m$ , which is parameterized by  $G$ -arc length, i.e., if  $T$  is the tangent to  $\gamma$ , then  $G(T, T) = 1$  and  $\gamma$  is defined on  $[0, L]$  where  $L$  is the  $G$ -length

of  $\gamma$ , so  $L \geq \frac{r(p)}{2}$ . Letting  $L_1$  be the  $G_1$ -length of  $\gamma$ ,

$$L_1 = \int_0^L \sqrt{G_1(T, T)} dt = \int_0^L (f \circ \gamma) dt = f(\bar{p})L > \frac{L}{r(\bar{p})}$$

where  $\bar{p}$  is on  $\gamma$  between  $p$  and  $m$ . But

$$|r(\bar{p}) - r(p)| \leq d(p, \bar{p}) \leq L$$

hence

$$r(\bar{p}) \leq r(p) + L, \quad L' > \frac{L}{r(p+L)} > \frac{L}{3L} = \frac{1}{3}$$

Hence  $d_1(p, m) \geq \frac{1}{3}$ , so  $B_1(p, \frac{1}{3}) \subset B(p, \frac{r(p)}{2})$ .

For the second part of the theorem we may assume  $G = G_1$  is complete. Fix a point  $m \in M$  and let  $f \in C^\infty(M, \mathbb{R})$  such that  $f(p) > d(m, p)$  for all  $p$ . Let

$$G_2 = e^{-2f} G$$

and we show  $G_2$  is bounded. Take  $p \in M$  and let  $\gamma$  be a geodesic from  $m$  to  $p$  with tangent  $T$  such that  $G(T, T) = 1$ ,  $\gamma$  is defined on  $[0, L]$ , and  $L = d(m, p)$ . Then

$$f \circ \gamma(t) > d(m, \gamma(t)) = t$$

for all  $t$ . Letting  $L_2$  be the  $G_2$ -length of  $\gamma$ ,

$$L_2 = \int_0^L \sqrt{G_2(T, T)} dt = \int_0^L e^{-f} dt < \int_0^L e^{-t} dt < \int_0^\infty e^{-t} dt = 1$$

Hence  $d_2(m, p) < 1$  for all  $m$  and  $p$ . □

**Corollary 10.32.** Every Riemannian metric on a manifold is complete iff the manifold is compact.

For further work on completeness, see the papers of J. A. Wolf and P. A. Griffiths.

## 10.6 Manifolds with Constant Riemannian Curvature

**Theorem 10.33.** Let  $M$  and  $M'$  be connected Riemannian manifolds with  $M$  complete. Let  $f : M \rightarrow M'$  be an isometry. Then  $f$  is an onto covering map and  $M'$  is complete.

*Proof.* To show  $f$  is onto, we show  $f(M)$  is open (which is trivial since  $f$  is a local diffeo) and closed. Take  $m' \in f(\bar{M})$ , let  $B'$  be a convex neighborhood of  $m'$ , let  $p' = f(p)$  be in  $B'$ , and let  $g'$  be the unique geodesic in  $B'$  from  $p'$  to  $m'$  with  $g'(0) = p'$  and  $g'(1) = m'$ . Let  $g$  be the unique geodesic in  $M$  with  $g(0) = p$  and  $f_*T_g(0) = T_{g'}(0)$ . Since  $f$  is an isometry,  $f \circ g$  is a geodesic in  $M'$ , and by uniqueness,  $f \circ g = g'$ . Since  $M$  is complete,  $g(1) = m$  is defined; hence  $f(m) = m'$  and  $f$  is onto. We have also shown  $M'$  is complete.

It is trivial that  $f$  evenly covers, since  $f$  preserves locally convex neighborhoods; thus for  $m'$  we choose a convex neighborhood  $B'$ , and  $f^{-1}(B')$  is a union of disjoint convex neighborhoods, each of which  $f$  maps diffeomorphically onto  $B'$ . □

**Theorem 10.34.** Let  $M$  be a connected, simply connected, complete Riemannian manifold with constant Riemannian curvature  $K$ . Then  $M$  is isometric to Euclidean space, spherical space, or hyperbolic space, when  $K = 0$ ,  $K > 0$ , or  $K < 0$ , respectively.

*Proof.* Let  $g$  be a geodesic in  $M$  parameterized by arc length with  $g(0) = m$ . Let  $e$  be a parallel unit field along  $g$ , which is orthogonal to  $T$ , the unit tangent to  $g$ . Let  $Z(t) = a(t)e(t)$  be a  $C^\infty$  field along  $g$ . Then

$$\mathbf{D}_T Z = a'e, \quad \mathbf{D}_T^2 Z = a''e$$

Thus  $Z$  is a Jacobi field if  $\mathbf{D}_T^2 Z = R(T, Z)T$  or

$$\langle \mathbf{D}_T^2 Z, Z \rangle = \langle R(T, Z)T, Z \rangle = -K \langle Z, Z \rangle \quad \text{i.e.} \quad a''a = -Ka^2$$

or  $a'' + Ka = 0$ . This differential equation has solutions uniquely determined by  $a(0)$  and  $a'(0)$ . If  $a(0) = 0$ , then  $Z(t) = (\exp_m)_* tA'$

where  $A'$  is the constant field on  $\mathbf{T}_m M$  with  $A' = a'(0)e$ . This equality follows from the fact that the right side is a Jacobi field and a Jacobi field is determined by  $Z$  and  $\mathbf{D}_T Z$  at one point. Hence

$$\langle Z, Z \rangle = t^2 \langle \exp_* A'(t), \exp_* A'(t) \rangle = a^2(t)$$

When  $K = 0$ , then  $a'' = 0$  and  $a = ct$  where  $c = a'(0)$ . Thus

$$\langle \exp_* A'(t), \exp_* A'(t) \rangle = c^2 = \langle A'(t), A'(t) \rangle$$

and  $\exp_m$  is an isometry from  $\mathbf{T}_m M$  onto  $M$ . Apply the previous theorem to obtain that  $\exp_m$  is a covering map. Since  $M$  is simply connected,  $\exp_m$  is a diffeo, hence  $M$  is isometric to  $\mathbf{T}_m M$ , and  $\mathbf{T}_m M$  is trivially isometric to Euclidean space.

When  $K < 0$ , let  $M'$  be hyperbolic space for  $K < 0$  (section 6.7). We know  $\exp_0 : \mathbf{T}_0 M' \rightarrow M'$  is a diffeo so let

$$E = (\exp_0)^{-1}$$

Choose an orthonormal base  $e_1, \dots, e_n$  of  $\mathbf{T}_0 M'$  and an orthonormal base  $e_1, \dots, e_n$  of  $\mathbf{T}_m M$ , where  $m \in M$  is arbitrary. Define  $F : \mathbf{T}_0 M' \rightarrow \mathbf{T}_m M$  by

$$F(e'_i) = e_i$$

Define  $f : M' \rightarrow M$  by

$$f = \exp_m \circ F \circ E$$

Then  $f_* Z'(t) = Z(t)$  along corresponding geodesics in  $M'$  and  $M$ , and

$$\langle f_* Z', f_* Z' \rangle = a^2(t) = \langle Z', Z' \rangle$$

Thus  $f_*$  is an isometry. Now the apply previous to obtain that  $f$  is a diffeo.

When  $K > 0$ , then  $Z = (\sin \sqrt{K}t)e$  is a Jacobi field along any geodesic emanating from  $m$  (a fixed point in  $M$ ). Thus every ray in

---

Section 10.6: Manifolds with Constant Riemannian Curvature

---

$\mathbf{T}_m M$  has a conjugate point at  $\frac{\pi}{\sqrt{K}}$  units from the origin and  $(\exp_m)_*$  has an  $(n - 1)$ -dimensional kernel at these points. Let

$$C = \left\{ X \in \mathbf{T}_m M : |X| = \frac{\pi}{\sqrt{K}} \right\}$$

Then  $\exp_m|_C$  is completely singular and hence is a constant map since  $C$  is connected. From the nature of the Jacobi equations in the first paragraph, there are no conjugate points in

$$B = \left\{ X \in \mathbf{T}_m M : |X| < \frac{\pi}{\sqrt{K}} \right\}$$

Now let  $M'$  be spherical space of curvature  $K$  and let  $p \in M'$ . We know  $\exp_p$  is a diffeo on the set  $B'$  (corresponding to  $B$ ) in  $\mathbf{T}_p M'$ . Define  $E$  and  $F$  as in the above paragraph ( $E$  defined on  $B(p, \frac{\pi}{\sqrt{K}})$ , the open ball), and let

$$f = \exp_m \circ F \circ E$$

on  $B(p, \frac{\pi}{\sqrt{K}})$  while  $f(-p) = \exp_m(C)$ . As in the above paragraph,  $f$  is an isometry on  $B(p, \frac{\pi}{\sqrt{K}})$ . Note what should be  $f_*$  at  $-p$  is well-defined via the tangent to incoming geodesics. Thus we may define a map  $g : B(-p, \pi/\sqrt{K}) \rightarrow M$  with

$$g(-p) = f(-p)$$

and  $g_*$  at  $-p$  determined by  $f_*$ . Then  $f = g$  on their common domain, and  $g$  is  $C^\infty$  and metric preserving at  $-p$ . Hence  $f$  is an isometry of  $M'$  onto  $M$ , and by the previous theorem,  $f$  is a diffeo.  $\square$

**Corollary 10.35.** Let  $M$  and  $M'$  be Riemannian manifolds, let  $b = \{e_1, \dots, e_n\}$  be an orthonormal base at  $m \in M$ , and similarly, let  $b'$  be such a base at  $m' \in M'$ . Define  $F : \mathbf{T}_m M \rightarrow \mathbf{T}_{m'} M'$  by

$$F(e_i) = e'_i$$

The map  $F$  induces a correspondence between geodesics emanating from  $m$  and  $m'$ , respectively, and also a correspondence between plane sections  $P$  and  $P'$  along these geodesics, via parallel translation of corresponding plane sections at  $m$  and  $m'$ . Thus for a geodesic  $g$  in  $M$  with  $g(0) = m$ , let  $g'$  be the geodesic in  $M'$  with  $g'(0) = m'$  and  $T_{g'}(0) = F(T_g(0))$ ; and for a plane section  $P$  in  $T_m M$ , let  $P(t)$  be the parallel translate of  $P$  along  $g$  to  $g(t)$ , let  $P' = F(P)$ , and  $P'(t)$  be the parallel translate of  $P'$  along  $g'$ . Suppose  $K'(P'(t)) = K(P(t))$  for all geodesics and plane sections (emanating from  $m$  and  $m'$ ). Then there are neighborhoods  $B$  and  $B'$  of  $m$  and  $m'$ , respectively, and a map  $f : B \rightarrow B'$  which is an isometry (and a diffeo). Thus  $M$  and  $M'$  are locally isometric at  $m$  and  $m'$ .

*Proof.* Choose an  $r > 0$  such that  $\exp_m$  is a diffeo from  $B(0, r)$  in  $T_m M$  onto  $B = B(m, r)$  in  $M$  and  $\exp_{m'}$  is also a diffeo from  $B'(0, r)$  in  $T_{m'} M'$  onto  $B' = B'(m', r)$  in  $M'$ . Let

$$f = \exp_m \circ F \circ (\exp_{m'})^{-1}$$

on  $B'$ , so  $f$  is a diffeo. By the method of proof in the preceding theorem,  $f$  is an isometry.  $\square$

If, in the above corollary, we add the hypothesis that  $M$  and  $M'$  are complete, connected, and simply connected, then it is an open question whether  $M$  is isometric to  $M'$ . When the Riemannian curvature is preserved for corresponding plane sections on once-broken geodesics, then Ambrose [Amb56] has proven  $M$  is isometric to  $M'$ .

## 10.7 Manifolds without Conjugate Points

Most of the results of the next two sections are based on a paper by A. Preissmann and some informal notes by W. B. Housing, Jr.

Throughout this section, let  $M$  be a complete connected Hausdorff Riemannian  $n$ -manifold. If  $m \in M$  and there exists no point of  $M$  that is conjugate to  $m$ , then  $m$  is called a *pole*.

**Theorem 10.36.** If  $m \in M$  is a pole, then  $\exp_m : \mathbf{T}_m M \rightarrow M$  is a covering map. Thus the simply connected covering of  $M$  is diffeomorphic to  $\mathbb{R}^n$ , and if  $M$  is simply connected, then  $M$  is diffeomorphic to  $\mathbb{R}^n$ .

*Proof.* Letting  $E = \exp_m$ , we know  $E$  is onto since  $M$  is complete, and  $E$  is a local diffeomorphism since  $m$  has no conjugate points. The metric tensor  $G$  of  $M$  induces a Euclidean metric on  $\mathbf{T}_m M$  whose distance function we denote by  $d$ . On the other hand, by requiring  $E$  to be an isometry, we define a metric tensor  $G_1$  on  $\mathbf{T}_m M$  whose distance function we denote by  $d_1$ . The rays in  $\mathbf{T}_m M$ , emanating from the origin, are  $G_1$ -geodesics since  $E$  is a connection preserving. We now show these rays are minimizing  $G_1$ -geodesics from the origin.

Take any  $X \in \mathbf{T}_m M$ , and let  $\gamma$  be a  $C^\infty$  curve from  $\mathbf{0}$  to  $X$  with  $\gamma(t) \in \bar{B}(0, |X|)$  for all  $t$  ( $B$  is the Euclidean ball). Assume  $\gamma$  is parameterized so  $|\dot{\gamma}(t)| = 1$ , thus  $\gamma$  is defined on  $[0, |X|]$ . Let  $T$  be the tangent to  $\gamma$ , then  $T_t = R_t + V_t$  where  $R$  is the unit (outward) radial vector field on  $\mathbf{T}_m M$  (and  $R_0 = T_0$ ), and  $V_t$  is orthogonal to  $R_t$  at each point. Computing the  $G_1$ -length of  $T$ ,

$$|T|_1 = |E_*(R + V)| \geq |E_*(R)| = 1$$

by the perpendicular lemma. Hence

$$|\gamma|_1 = \int_0^{|X|} |T|_1 dt \geq |X|$$

which implies  $d_1(0, X) = |X|$ , since the ray from  $\mathbf{0}$  to  $X$  has  $G_1$ -length equal to  $|X|$ . Thus  $\bar{B}_1(0, b) = \bar{B}(0, b)$  for all  $b \geq 0$ , and since the latter is compact, so is the former. By the complete theorem (10.26),  $\mathbf{T}_m M$  is complete with respect to the  $G_1$ -metric. By theorem 10.33, the map  $E : \mathbf{T}_m M \rightarrow M$  is a covering map.  $\square$

**Corollary 10.37.** If  $M$  has non-positive Riemannian curvature, then all points are poles and  $\mathbb{R}^n$  is a simply connected covering space of  $M$ .

We now define the *universal covering manifold*  $\bar{M}$ , based at a point  $m \in M$ , in a standard way. Let  $\bar{M}$  be the set of equivalence classes of  $C^0$ -homotopic  $C^0$ -curves  $f$  defined on a finite interval such that  $f(0) = m$  (see Hocking-Young, p. 188). Let  $[f]$  denote the equivalence class of a curve  $f$ , and let  $\pi: \bar{M} \rightarrow M$  denote the covering map where  $\pi([f])$  is the endpoint of  $f$ . Define a  $C^\infty$  structure on  $\bar{M}$  by demanding  $\pi$  to be a  $C^\infty$  map, and if  $M$  is Riemannian, define a Riemannian metric on  $\bar{M}$  such that  $\pi$  is an isometry. We use repeated the fact that a  $C^0$ -curve  $f$  in  $M$  has a unique lifting  $\bar{f}$  in  $\bar{M}$  such that  $\pi \circ \bar{f} = f$  once one has prescribed  $\bar{f}(0)$ . Let  $f \sim h$  denote the fact that  $f$  is homotopic to  $h$  under a fixed end-point homotopy, and let  $\bar{m}$  be the constant path at  $m$ .

**Theorem 10.38.** Let  $f$  be a finite curve in  $M$  and let

$$b = \inf \{ |h| : h \text{ is a broken } C^\infty \text{ curve and } h \sim f \}$$

Then there exists a geodesic  $g$  such that  $g \sim f$  and  $|g| = b$ . Thus in every homotopy class of curves (with fixed end-points), there is a geodesic whose length is the absolute minimum for the lengths of all broken  $C^\infty$  curves in the homotopy class.

*Proof.* Let  $\bar{M}$  be the universal covering manifold based at  $m = f(0)$ . Since  $M$  is complete,  $\bar{M}$  is complete, and hence there exists a geodesic  $\bar{g}$  from  $[\bar{m}]$  to  $[f]$  which gives the distance in  $\bar{M}$  between these two points. Then  $g = \pi \circ \bar{g}$  is a geodesic in  $M$  since  $\pi$  is an isometry, and  $g \sim f$  since  $\bar{M}$  is simply connected. If  $h$  is a broken  $C^\infty$  curve with  $h \sim f$ , then lift  $h$  to a curve  $\bar{h}$  starting at  $[\bar{m}]$  and obtain a broken  $C^\infty$  curve  $\bar{h}$  from  $[\bar{m}]$  to  $[f]$ . Since  $\bar{g}$  gives the distance,  $|\bar{h}| \geq |\bar{g}| = |g|$ , thus  $|g| = b$ .  $\square$



**Theorem 10.39.** Let  $m \in M$  be a pole and let  $g_1, g_2$  be geodesics emanating from  $m$  that intersect later. If  $g_1 \sim g_2$ , then  $g_1 = g_2$  (when both parameterized by arc length).

*Proof.* Let  $\bar{M}$  be the universal covering manifold based at  $m$  with  $\pi$  an isometry. Define  $\bar{\exp} : \mathbf{T}_m M \rightarrow \bar{M}$  by

$$\bar{\exp}(X) = \{\exp_m tX : 0 \leq t \leq 1\}$$

Then  $\pi \circ \bar{\exp} = \exp_m$  and  $\bar{\exp}$  is  $C^\infty$ , since locally  $\bar{\exp} = \pi^{-1} \circ \exp_m$ . Moreover,  $\bar{\exp}$  is an isometry, for  $m$  is a pole. Since  $\mathbf{T}_m M$  is simply connected,  $\bar{\exp}$  is a diffeo by theorem 10.33. If  $g_1(t) = \exp tX_1$  where  $g_1(1) = g_2(1)$ , and  $g_1 \sim g_2$ , then  $[g_1] = [g_2]$ . Since  $\bar{\exp}$  is a diffeo, this implies  $X_1 = X_2$ , which implies  $g_1 = g_2$ . □

We remark that one can always define the  $C^\infty$  map  $\bar{\exp} : \mathbf{T}_m M \rightarrow \bar{M}$  (base point  $m$ ) with  $\pi \circ \bar{\exp} = \exp_m$ . The map  $\bar{\exp}$  will be onto if  $M$  is complete, but it will not in general be locally one-to-one.

**Corollary 10.40.** If  $m \in M$  is a pole and  $M$  is simply connected, then for any point  $p \in M$  there is a unique geodesic through  $m$  and  $p$ .

**Corollary 10.41.** If  $M$  is simply connected and has only non-positive Riemannian curvature, then there is a unique geodesic through any two points of  $M$ .

## 10.8 Manifolds with Non-Positive Curvature

We add to the standard hypothesis of the last section the assumption that  $K(P) \leq 0$  for all plane sections  $P$  of  $M$ .

**Lemma 10.42.** Let  $f$  be a finite curve in  $M$  parameterized by arc length, and let  $m \in M$ . Let  $\bar{f}$  be any lifting of  $f$  to the covering space  $\mathbf{T}_m M$  (see theorem 10.36). Then  $|f| \geq |\bar{f}|$ , the Euclidean length of  $\bar{f}$  in  $\mathbf{T}_m M$ . If  $K < 0$ , then  $|f| > |\bar{f}|$  unless  $\bar{f}$  is a segment of a ray emanating from zero in  $\mathbf{T}_m M$ .

*Proof.* By theorem 10.23, if  $T$  is a vector tangent to  $\mathbf{T}_m M$ , then  $|(\exp_m)_* T| \geq |T|$ . If  $K < 0$ , then  $|(\exp_m)_* T| > |T|$  unless  $T$  is a radial vector tangent to a ray through zero.  $\square$

**Theorem 10.43.** Let  $p_1, p_2, p_3 \in M$  be distinct points joined by geodesics  $g_1, g_2, g_3$  where  $g_1$  joins  $p_2$  and  $p_3$ , etc., (see Fig. 10.6). Assume the three points are not on one geodesic and the broken loop formed by the three curves is homotopic to zero. Let  $\theta_i$  be the unique angle at  $p_i$  made by the intersecting geodesics with  $0 < \theta_i < \pi$ . Then

$$|g_1|^2 \geq |g_2|^2 + |g_3|^2 - |g_2| |g_3| \cos \theta_1, \quad \theta_1 + \theta_2 + \theta_3 \leq \pi$$

If  $K < 0$  on  $M$ , these inequalities are strict.

*Proof.* Let  $m = p_1$  and let  $\bar{g}_2, \bar{g}_3$  be the rays through zero in  $\mathbf{T}_m M$  such that  $\exp_m \circ \bar{g}_i = g_i$  for  $i = 2, 3$ . Let  $X_2, X_3$  be the endpoints of  $\bar{g}_3, \bar{g}_2$  respectively. Since the loop formed by  $g_2, g_1, g_3$  is homotopic to zero, we can lift  $g_1$  to a curve  $\bar{g}_1$  joining  $X_2, X_3$ . By the preceding lemma,

$$|g_1| \geq |\bar{g}_1| \geq d(X_2, X_3)$$

where  $d$  is the Euclidean distance in  $\mathbf{T}_m M$ . By the law of cosines in  $\mathbf{T}_m M$ ,

$$d(X_2, X_3)^2 = |g_2|^2 + |g_3|^2 - |g_2| |g_3| \cos \theta_1$$

which proves the first inequality.

For the second inequality, we construct a triangle in  $\mathbb{R}^2$  whose sides have lengths  $a_i = |g_i|$  and label the angles at the appropriate corners by  $\phi_i$ . Then

$$(a_1)^2 = (a_2)^2 + (a_3)^2 - a_2 a_3 \cos \phi_1$$

hence  $\cos \theta_1 \geq \cos \phi_1$  and  $\theta_1 \leq \phi_1$ . Similarly,  $\theta_i \leq \phi_i$  for  $i = 2, 3$ , thus

$$\theta_1 + \theta_2 + \theta_3 \leq \phi_1 + \phi_2 + \phi_3 = \pi$$

If  $K < 0$ , then  $|g_1| > |\bar{g}_1|$  and the strict inequalities then follow.  $\square$

**Corollary 10.44.** The sum of the interior angles ( $0 \leq \theta_i < \pi$ ) of a geodesic quadrilateral which is homotopic to zero is less than or equal to  $2\pi$ . If  $K < 0$ , then the sum is less than  $2\pi$ .




Figure 10.6: Geodesic Triangle

**Corollary 10.45.** Let  $m \in M$ , and let  $g$  be a geodesic that does not pass through  $m$ . Then there cannot be two distinct geodesics  $g_1, g_2$  from  $m$  to  $g$  which intersect  $g$  orthogonally such that the geodesic triangle formed is homotopic to zero.

*Proof.* The sum of the interior angles of the geodesic triangle would be greater than  $\pi$ .  $\square$

**Corollary 10.46.** Let  $M$  be simply connected,  $m \in M$ , and  $g$  a geodesic that does not pass through  $m$ . Then there is a unique geodesic  $f$  from  $m$  to  $g$  which is orthogonal to  $g$  and  $|f| \leq d(m, g(t))$  for all  $t$ .

*Proof.* Let  $f_t$  be the unique geodesic from  $m$  to  $g(t)$ , let

$$L(t) = |f_t| = d(m, g(t))$$

and let  $g_t$  be  $g$  restricted to the interval  $[0, t]$  or  $[t, 0]$ , as the case may be. Let  $\theta$  be the angle between  $f_0$  and  $g_t$  for  $t > 0$ . We show that  $L(t) \rightarrow \infty$  as  $t \rightarrow \pm\infty$ . For  $t > 0$ ,

$$L^2(t) = |f_t|^2 \geq |f_0|^2 + |g_t|^2 - |f_0||g_t| \cos \theta = |f_0|^2 + |g_t|(|g_t| - |f_0| \cos \theta)$$

As  $t \rightarrow \infty$ , we have  $|g_t| \rightarrow \infty$ , hence  $L(t) \rightarrow \infty$ . Similarly,  $L(t) \rightarrow \infty$  as  $t \rightarrow -\infty$ .

By theorem 10.13, a point  $t'$  is a critical point of  $L$  iff  $f_{t'}$  is orthogonal to  $g$ . By corollary 10.45 there can be at most one critical point of  $L$ , and that must be an absolute minimum by the first paragraph.  $\square$

For further results, see Preissman [Pre43] and Helgason [Hel12].

## Problems

93. Using the notation of section 3.4,

- (a) show that  $T_v$  is a Jacobi field on a surface of revolution.
- (b) If  $G = \langle T_v, T_v \rangle$  and  $S$  is arc length along the meridians, show

$$\frac{d^2 \sqrt{G}}{ds^2} = -K \sqrt{G}$$

94. If  $M$  is a complete Riemannian 2-manifold, show the locus of first (those nearest the origin on each ray) conjugate points in  $\mathbf{T}_m M$  is a  $C^\infty$  curve (see [Mye35]).

95. Show the Hessian is well-defined, symmetric, and bilinear.

96. If  $d$  is the function defined in lemma 10.24, show  $d$  is  $C^\infty$  on  $C$ .

97. If  $M = \mathbb{R}^3$ ,  $g$  is the  $x$ -axis from  $(a, 0, 0)$  to  $(b, 0, 0)$  with  $a < b$ , and  $C_1$  and  $C_2$  are the planes  $x = a$  and  $x = b$  respectively, check lemma 10.24.

98. A submanifold  $V$  of a manifold  $M$  is totally geodesic with respect to a connexion  $D$  if any geodesic that is tangent to  $V$  at a point lies wholly in  $V$ . If  $V$  and  $W$  are compact totally geodesic submanifolds of dimension  $r$  and  $s$  respectively, lying in a Riemannian  $n$ -manifold  $M$  of positive Riemannian curvature and  $r + s \geq n$ , show  $V \cap W$  is non-empty (see [Fra61]).
99. Find a condition relating curvature and parallel translation that will ensure the existence of complete totally geodesic submanifolds in a Riemannian manifold (see [Her60] or [Hel12]).
100. If  $M$  is an oriented  $n$ -manifold and  $\alpha$  is a  $C^\infty$   $(n - 1)$ -form on  $M$  with compact support, show  $\int_M d\alpha = 0$  (see [NJ62]).



## Bibliography

- [Adl57] A. Adler. “Characteristic Classes of Homogeneous Spaces”. In: *Transactions of the American Mathematical Society* 86 (2 1957), pp. 348–365. url: <https://www.jstor.org/stable/1993020>.
- [AH37] P. Alexandroff and H. Hopf. “Topologie I”. In: *The Mathematical Gazette* 21 (1937), pp. 296–298. url: <https://www.jstor.org/stable/3607723>.
- [AL60] L.V Ahlfors and L.Sario. *Riemann Surfaces*. princeton university press, 1960. isbn: 9780691626123. url: <https://zr9558.files.wordpress.com/2013/11/lars-v-ahlfors-l-sario-riemann-surfaces.pdf>.
- [All40] C.B. Allendoerfer. “The Euler Number of a Riemann Manifold”. In: *American Journal of Mathematics* 62 (1 1940), pp. 243–248. url: [https://www.jstor.org/stable/2371450#metadata\\_info\\_tab\\_contents](https://www.jstor.org/stable/2371450#metadata_info_tab_contents).
- [All50] C.B. Allendoerfer. “Characteristic Cohomology Classes in a Riemann Manifold”. In: *Annals of Mathematics* 51 (3 1950), pp. 551–570. url: <https://www.jstor.org/stable/1969368>.
- [AM12] L. Auslander and R.E. MacKenzie. *Introduction to Differentiable Manifolds*. Dover Books on Mathematics. Dover Publications, 2012. isbn: 9780486158082. url: <https://books.google.com/books?id=J98LQTpm0wIC>.

- [Amb56] W. Ambrose. "Parallel Translation of Riemannian Curvature". In: *Annals of Mathematics* 64 (2 1956), pp. 337–363. url: <https://www.jstor.org/stable/1969978>.
- [Amb60] W. Ambrose. "The Cartan Structural Equations in Classical Riemannian Geometry". In: *The Journal of the Indian Mathematical Society* 24 (1960), pp. 23–76. url: <https://www.informaticsjournals.com/index.php/jims/article/view/16900>.
- [Amb61] W. Ambrose. "The Index Theorem in Riemannian Geometry". In: *Annals of Mathematics* 73 (1 1961), pp. 49–86. url: <https://www.jstor.org/stable/1970282>.
- [AW43] C.B. Allendoerfer and A. Weil. "The Gauss-Bonnet Theorem for Riemannian Polyhedra". In: *Transactions of the American Mathematical Society* 53 (1943), pp. 101–129. url: <https://www.jstor.org/stable/1990134>.
- [BC11] R.L. Bishop and R.J. Crittenden. *Geometry of Manifolds*. ISSN. Elsevier Science, 2011. isbn: 9780080873275. url: <https://books.google.com/books?id=sRaSuentwngC>.
- [BF34] T. Bonnesen and W. Fenchel. *Theorie der konvexen Körper*". *Ergebnisse der Mathematik Und Ihrer Grenzgebiete*. Springer, 1934. isbn: 3540062343. url: <https://www.goodreads.com/book/show/14711691-theorie-der-konvexen-korper>.
- [Buc03] R.C. Buck. *Advanced Calculus: Third Edition*. Waveland Press, 2003. isbn: 9781478608202. url: <https://books.google.com/books?id=7cYQAAAAQBAJ>.
- [Cai] S.S. Cairns. "A Simple Triangulation Method for Smooth Manifolds". In: *Bulletin of the American Mathematical Society* 67 (), pp. 389–390. url: <https://www.ams.org/journals/bull/1961-67-04/S0002-9904-1961-10631-9/S0002-9904-1961-10631-9.pdf>.



- [Car25] E. Cartan. *La géométrie des espaces de Riemann*. fr. Gauthier-Villars, 1925. url: <http://eudml.org/doc/192543>.
- [Che45] S. S. Chern. "On the Curvatura Integra in a Riemannian Manifold". In: *Annals of Mathematics* 46.4 (1945), pp. 674–684. issn: 0003486X. url: <http://www.jstor.org/stable/1969203> (visited on 07/20/2022).
- [Che46] C. Chevalley. *Theory of Lie Groups*. Princeton University Press, 1946. isbn: 9780691049908. url: <https://archive.org/details/in.ernet.dli.2015.86469/page/n13/mode/2up>.
- [Che51] S. S. Chern. *Topics in Differential Geometry*. Topics in Differential Geometry. Institute for Advanced Study, 1951. url: <https://www.intlpress.com/site/pub/pages/books/items/00000488/index.php>.
- [Che56] C. Chevalley. *Fundamental concepts of algebra*. New York, Academic Press, 1956. isbn: 9780080873152. url: [https://archive.org/details/fundamentalconce00chev\\_0](https://archive.org/details/fundamentalconce00chev_0).
- [CL90] A. Coddington and N. Levinson. *Theory of Ordinary Differential Equations*. Tata McGraw- Hill Publ. Company Limited, 1990. url: <https://ptvtpqa.files.wordpress.com/2013/12/coddington-e-levinson-n-theory-of-ordinary-differential-equations.pdf>.
- [Eis15] L.P. Eisenhart. *An Introduction to Differential Geometry with Use of the Tensor Calculus*. Creative Media Partners, LLC, 2015. isbn: 9781296828103. url: <https://books.google.com/books?id=HbU1jgEACAAJ>.
- [Fen40] Werner Fenchel. "On Total Curvatures of Riemannian Manifolds: I". In: *Journal of The London Mathematical Society-second Series* (1940), pp. 15–22. url: <https://londmathsoc.onlinelibrary.wiley.com/doi/abs/10.1112/jlms/s1-15.1.15>.

- [Fra61] Theodore Frankel. “Manifolds with positive curvature”. In: *Pacific Journal of Mathematics* 11 (1961), pp. 165–174. url: <https://msp.org/pjm/1961/11-1/pjm-v11-n1-p12-s.pdf>.
- [Gri58] John S. Griffin Jr. “On the rotation number of a normal curve”. eng. In: *Compositio Mathematica* 13 (1958), pp. 270–276. url: <http://eudml.org/doc/88841>.
- [Gug12] H.W. Guggenheimer. *Differential Geometry*. Dover Books on Mathematics. Dover Publications, 2012. isbn: 9780486157207. url: <https://archive.org/details/differentialgeom0000gugg>.
- [Hal17] P.R. Halmos. *Finite-Dimensional Vector Spaces: Second Edition*. Dover Books on Mathematics. Dover Publications, 2017. isbn: 9780486814865. url: <https://www.maa.org/press/maa-reviews/finite-dimensional-vector-spaces-0>.
- [HC99] D. Hilbert and S. Cohn-Vossen. *Geometry and the Imagination*. AMS Chelsea Publishing Series. AMS Chelsea Pub., 1999. isbn: 9780821819982. url: <https://www.maa.org/press/maa-reviews/geometry-and-the-imagination>.
- [Hel12] S. Helgason. *Differential Geometry, Lie Groups, And Symmetric Spaces*. Academic Press, 2012. isbn: 9780124157316. url: <https://bookstore.ams.org/gsm-34>.
- [Her60] Robert Hermann. “Existence in the large of totally geodesic submanifolds of Riemannian spaces”. In: *Bulletin of the American Mathematical Society* 66 (1960), pp. 59–61. url: <https://www.ams.org/journals/bull/1960-66-01/S0002-9904-1960-10391-6/S0002-9904-1960-10391-6.pdf>.
- [Hic63a] N. J. Hicks. “Connexion Preserving, Conformal, And Parallel Maps”. In: *Michigan Math Journal* 10 (1963), pp. 295–302. url: [https://projecteuclid.org/download/pdf\\_1/euclid.mmj/1028998915](https://projecteuclid.org/download/pdf_1/euclid.mmj/1028998915).

- [Hic63b] N. J. Hicks. "Submanifolds of Semi-Riemannian Manifolds". In: *Rendiconti del Circolo Matematico di Palermo* 12 (1963), pp. 137–149. url: <https://link.springer.com/article/10.1007/BF02843960>.
- [Hir76] Morris W. Hirsch. *Differential Topology*. Graduate Texts in Mathematics. Springer-Verlag, 1976. isbn: 9780486141091. url: [https://archive.org/details/springer\\_10.1007-978-1-4684-9449-5](https://archive.org/details/springer_10.1007-978-1-4684-9449-5).
- [Hop26] H. Hopf. "Über die Curvatura integra geschlossener Hyperflächen". In: *Mathematische Annalen* 95 (1926), pp. 340–367. url: <http://eudml.org/doc/159134>.
- [Hop50] H. Hopf. "Über Flächen mit einer Relation zwischen den Hauptkrümmungen. Meinem Lehrer Erhard Schmidt in Verehrung und Freundschaft zum 75. Geburtstag gewidmet." In: *Mathematische Nachrichten* 4.1-6 (1950), pp. 232–249. doi: <https://doi.org/10.1002/mana.3210040122>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/mana.3210040122>. url: <https://onlinelibrary.wiley.com/doi/abs/10.1002/mana.3210040122>.
- [HS13] D. C. Spencer H. K. Nickerson and N. E. Steenrod. *Advanced Calculus*. Dover Books on Mathematics. Dover Publications, 2013. isbn: 9780486173740. url: <https://books.google.com/books?id=KgHDagAAQBAJ>.
- [Hsi82] Wu-yi Hsiang. "Generalized rotational hypersurfaces of constant mean curvature in the Euclidean spaces. I". In: *Journal of Differential Geometry* 17 (1982), pp. 337–356. url: <https://www.semanticscholar.org/paper/Generalized-rotational-hypersurfaces-of-constant-in-Hsiang/bbea07a96aee6a20c14f718f946>
- [HY12] J.G. Hocking and G.S. Young. *Topology*. Dover Books on Mathematics. Dover Publications, 2012. isbn: 9780486141091. url: [https://download.tuxfamily.org/openmathdep/analysis\\_real/Topology-Hocking.pdf](https://download.tuxfamily.org/openmathdep/analysis_real/Topology-Hocking.pdf).

- [Jac13] N. Jacobson. *Lectures in Abstract Algebra: II. Linear Algebra*. Graduate Texts in Mathematics. Springer New York, 2013. isbn: 9781468470536. url: [https://archive.org/details/springer\\_10.1007-978-1-4684-7053-6](https://archive.org/details/springer_10.1007-978-1-4684-7053-6).
- [Joh60] H. H. Johnson. "Terminating prolongation procedures." In: *Pacific Journal of Mathematics* 10.2 (1960), pp. 577–583. url: <https://msp.org/pjm/1960/10-2/pjm-v10-n2-p16-s.pdf>.
- [JW69] M. Spivak J. Milnor and R. Wells. *Morse Theory*. (AM-51), Volume 51. Princeton University Press, 1969. isbn: 9780691080086. url: [https://drive.google.com/file/d/1yixyw3Gpzs6fvBxxWUL\\_XRamWsdFCagL/view](https://drive.google.com/file/d/1yixyw3Gpzs6fvBxxWUL_XRamWsdFCagL/view).
- [Kap03] W. Kaplan. *Advanced Calculus*. Addison-Wesley higher mathematics. Addison-Wesley, 2003. isbn: 9780201799378. url: <http://hore.dnom.fmph.uniba.sk/~demetrian/lect/biomedmat4/kaplan.pdf>.
- [Kel17] J. L. Kelley. *General Topology*. Dover Books on Mathematics. Dover Publications, 2017. isbn: 9780486815442. url: <https://archive.org/details/GeneralTopologyJohnL.Kelley/page/n21/mode/2up>.
- [KN63] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry, Volume 2*. Wiley, 1963. isbn: 9780470496480. url: <https://thunv.files.wordpress.com/2008/05/foundation-of-differential-geometry-ii-kobayashinomuzu.pdf>.
- [Kur57] M. Kuranishi. "On E. Cartan's Prolongation Theorem of Exterior Differential Systems". In: *American Journal of Mathematics* 79.1 (1957), pp. 1–47. issn: 00029327, 10806377. url: <http://www.jstor.org/stable/2372381> (visited on 07/20/2022).
- [Lan14] S. Lang. *Introduction to Differentiable Manifolds*. Springer, 2014. isbn: 9781475777444. url: <https://books.google.com/books?id=grTWzgEACAAJ>.

- [Lev77] T. Levi-Civita. *The Absolute Differential Calculus (calculus of Tensors)*. Dover Publications, 1977. isbn: 9780486634012. url: <https://archive.org/details/TheAbsoluteDifferentialCalculus>.
- [Mac04] G. W. Mackey. *Mathematical Foundations of Quantum Mechanics*. Dover Books on Mathematics. Dover Publications, 2004. isbn: 9780486435176. url: <https://books.google.com/books?id=Gcsz0c39QzgC>.
- [Mil56] J. W. Milnor. "On Manifolds Homeomorphic to the 7-Sphere". In: *Annals of Mathematics* 64 (1956), p. 399. url: <https://www.jstor.org/stable/1969983>.
- [Mor34] M. Morse. *The Calculus of Variations in the Large*. American Mathematical Society v. 18. American Mathematical Society, 1934. isbn: 9780821810187. url: <https://books.google.com/books?id=ouOVawAAQBAJ>.
- [Mun59] J. R. Munkres. "Obstructions to the smoothing of piecewise-differentiable homeomorphisms". In: *Bulletin of the American Mathematical Society* 65 (1959), pp. 332–334. url: <https://www.jstor.org/stable/1970228>.
- [Mun66] J. R. Munkres. *Elementary Differential Topology*. Annals of Mathematics Studies no. 54, pt. 1. Princeton University Press, 1966. isbn: 9780691090931. url: <https://www.maths.ed.ac.uk/~v1ranick/papers/munkresdiff.pdf>.
- [MW97] J. W. Milnor and D.W. Weaver. *Topology from the Differentiable Viewpoint*. Princeton Landmarks in Mathematics and Physics. Princeton University Press, 1997. isbn: 9780691048338. url: <https://books.google.co.in/books?id=ORCnDwAAQBAJ>.
- [Mye35] S. B. Myers. "Connections between Differential Geometry and Topology". In: *Proceedings of the National Academy of Sciences* 21.4 (1935), pp. 225–227. url: <https://www.pnas.org/doi/abs/10.1073/pnas.21.4.225>.

- [NJ62] A. Nijenhuis and R. W. Richardson Jr. "A theorem on maps with non-negative Jacobians." In: *Michigan Mathematical Journal* 9.2 (1962), pp. 173–176. url: <https://projecteuclid.org/journals/michigan-mathematical-journal/volume-9/issue-2/A-theorem-on-maps-with-non-negative-Jacobians/10.1307/mmj/1028998676.full>.
- [Nom21] K. Nomizu. *Lie Groups and Differential Geometry*. Creative Media Partners, LLC, 2021. isbn: 9781013914218. url: <https://books.google.com/books?id=3y-wzgEACAAJ>.
- [Pal54] R. S. Palais. "A Definition of the Exterior Derivative in Terms of Lie Derivatives". In: *Proceedings of the American Mathematical Society* 5.6 (1954), pp. 902–908. issn: 00029939, 10886826. url: <http://www.jstor.org/stable/2032554> (visited on 07/20/2022).
- [Pal57] R. S. Palais. "A global formulation of the Lie theory of transformation groups". In: *Memoirs of the American Mathematical Society* 22 (1957). url: <https://vmm.math.uci.edu/PalaisPapers/AGloblFormlationOfLieTheory.pdf> (visited on 07/23/2022).
- [PB14] W.J. Pervin and R.P. Boas. *Foundations of General Topology*. Academic Press textbooks in mathematics. Elsevier Science, 2014. isbn: 9781483225159. url: <https://books.google.com/books?id=QpXiBQAAQBAJ>.
- [Pog56] A.V Pogorelov. "A New Proof of Rigidity of Convex Polyhedra". In: *Uspekhi Matematicheskikh Nauk* 11 (5 1956), pp. 207–208. url: <http://mi.mathnet.ru/eng/umn/v11/i5/p207>.
- [Pre43] A. V. Preissmann. "Quelques propriétés globales des espaces de Riemann." In: *Commentarii mathematici Helvetici* 15 (1943), pp. 175–216. url: <http://eudml.org/doc/138804>.

- [Rha52] G. de Rham. "Sur la réductibilité d'un espace de Riemann." In: *Commentarii mathematici Helvetici* 26 (1952), pp. 328–344. url: <http://eudml.org/doc/139052>.
- [Sam55] H. Samelson. *Lecture Notes in Differential geometry*. 1955.
- [Sma61] S. Smale. "Differentiable and Combinatorial Structures on Manifolds". In: *Annals of Mathematics* 74 (3 1961), pp. 498–502. url: <https://www.jstor.org/stable/1970294>.
- [ST51] H. Seifert and W. Threlfall. *Variationsrechnung im Grossen: (Theorie von Marston Morse*. AMS Chelsea Publishing Series. Chelsea Publishing Company, 1951. isbn: 9780828400497. url: [https://books.google.com/books?id=1YM%5C\\_AAAAYAAJ](https://books.google.com/books?id=1YM%5C_AAAAYAAJ).
- [Ste51] N. Steenrod. *The Topology of Fibre Bundles*. Princeton University Press, 1951. isbn: 9780691005485. url: <https://press.princeton.edu/books/paperback/9780691005485/the-topology-of-fibre-bundles-pms-14-volume-14>.
- [Ste64] S. Sternberg. *Lectures on Differential Geometry*. Chelsea Publishing Company, 1964. url: <https://archive.org/details/lecturesondiffer0000ster/page/n5/mode/2up>.
- [Str61] D.J. Struik. *Lectures on Classical Differential Geometry*. Addison-Wesley series in mathematics. Addison-Wesley Publishing Company, 1961. isbn: 9780486656090. url: <https://archive.org/details/dli.ernet.469746>.
- [Tit] C. J. Titus. "A Theory of Normal Curves and Some Applications". In: *Pacific Journal Of Mathematics* 10 (), pp. 1083–1096. url: <https://projecteuclid.org/journals/pacific-journal-of-mathematics/volume-10/issue-3/A-theory-of-normal-curves-and-some-applications/pjm/1103038253.pdf>.
- [VW60] O. Veblen and J. H. C. Whitehead. *The Foundations of Differential Geometry*. Cambridge tracts in mathematics

- and mathematical physics. University Press, 1960. url: <https://books.google.com/books?id=KC9uAAAAIAAJ>.
- [Wei52] A. Weil. “Sur les théorèmes de de Rham.” In: *Commentarii mathematici Helvetici* 26 (1952), pp. 119–145. url: <http://eudml.org/doc/139040>.
- [Wen86] H. C. Wente. “Counterexample to a conjecture of H. Hopf.” In: *Pacific Journal of Mathematics* 121.1 (1986), pp. 193–243. url: <https://projecteuclid.org/journals/pacific-journal-of-mathematics/volume-121/issue-1/Counterexample-to-a-conjecture-of-H-Hopf/pjm/1102702809.full>.
- [Whi16] H. Whitney. *Geometric Integration Theory*. Princeton Legacy Library. Princeton University Press, 2016. isbn: 9780691652900. url: <https://www.maths.ed.ac.uk/~v1ranick/papers/git.pdf>.
- [Whi32] J. H. C. Whitehead. “Convex Regions In The Geometry Of Paths”. In: *The Quarterly Journal of Mathematics* 3.1 (Jan. 1932), pp. 33–42. issn: 0033-5606. url: <https://aareyanmanzoor.github.io/assets/whitehead1932.pdf>.
- [Whi37] H. Whitney. “On regular closed curves in the plane”. eng. In: *Compositio Mathematica* 4 (1937), pp. 276–284. url: <http://eudml.org/doc/88656>.
- [Wil59] T. Willmore. *An Introduction to Differential Geometry*. Clarendon, 1959. url: <https://books.google.com/books?id=dHHRswEACAAJ>.
- [WS60] R.S Palais W. Ambrose and I.M Singer. “Sprays”. In: *Anaes da Academia Brasileira de Ciencias* 32 (2 1960), pp. 163–178. url: <http://memoria.bn.br/DocReader/docreader.aspx?bib=158119&pesq=&pagfis=12553>.



# Index

- alternating tensor, 82
- area, 115, 155
- asymptotic coordinate, 200
- asymptotic vector, 119
- atlas, 16
  
- base field, 31
- beltrami operators, 204
- Bianchi Identities, 114, 145
- boundary map, 149
- bounding curve, 159
- bundle of bases, 36
  
- Cartan differential algebra, 140
- Cartan formula (geodesic curvature), 162
- catenoid, 79
- center surfaces, 78
- chains, 148
- characteristic classes, 174
- characteristic forms, 174
- characteristics, 33
- closed distributions, 183
- cochains, 150
  
- Codazzi-Mainardi equations, 51, 118, 126, 135
- cohomology group(differential), 174
- complete, 98, 112, 237
- conformal, 115
- conformal coordinates, 200
- congruent surfaces, 177
- conjugate degree, 229
- conjugate point, 214
- conjugate vectors, 47, 119
- connexion, 41, 49, 93, 104
- connexion 1-forms, 98, 105
- connexion preserving map, 97
- contraction, 140
- convex neighborhood, 197
- coordinate pair, 16
- coordinate system, 17
- coordinate systems (special), 30
- cotangent bundle, 90
- covariant derivative, 40, 41, 49, 93, 144
- critical point, 230

cube, 148  
 curve, 27  
 curve (broken), 30  
  
 developable surface, 56  
 diffeomorphism, 26  
 difference tensor, 101  
 differential of a map, 26  
 divergence, 145  
 Dupin indicatrix, 47  
  
 Euler characteristic, 164  
 excess, 163  
 exponential map, 193  
 exterior algebra, 84  
 exterior derivative, 99, 137  
 exterior product, 84  
 extremal, 218  
  
 Fermi coordinates, 196  
 fiber, 37  
 first variation formulae, 217  
 flat connexion, 107  
 flat point, 48, 119  
 flat space, 126  
 forms, 83  
 framed, 85  
 Frenet formulae, 116  
 functions,  $C^s$ , 19  
 fundamental 1-form, 90  
 fundamental forms, 46  
 fundamental  $n$ -chain, 154  
 fundamental set, 154  
  
 Gauss curvature, 46  
  
 Gauss curvature equation, 51, 118, 126, 135  
 Gauss equation, 48, 117, 124  
 Gauss map, 44  
 Gauss-Bonnet form, 171  
 general linear group  $GL(n, \mathbb{R})$ , 17  
 geodesic, 41, 49, 94  
 geodesic curvature, 115, 157, 161  
 geodesic flow field, 194  
 geodesic preserving map, 97  
 geodesic submanifold, 229, 253  
 gradient, 147  
 Grassman algebra, 84  
 Grassman product, 84  
  
 helicoid, 63  
 Hessian, 228  
 holonomy, 108  
 homology, 149  
 homothetic, 115  
 hyperbolic space, 127  
 hypersurface, 43  
  
 imbedding, 31  
 immersion, 31  
 index (function), 228  
 index (vector field), 167, 172  
 inner product, 42  
 integrable distribution, 183  
 integral curve, 29  
 integral manifold, 183

invariant polynomial, 174  
 involutive distribution, 183  
 isometric, 115  
 isometry, 115  
 isothermal coordinates, 200  
  
 Jacobi field, 212  
 Jacobi identity, 25  
 Jacobian map, 26  
  
 Lagrange identity, 221  
 Laplacian, 147, 204  
 left invariant connexion, 107  
 left invariant fields, 35  
 length (curve), 109  
 length (vector), 109  
 Lie algebra, 35  
 Lie bracket, 24  
 Lie derivative, 141  
 Lie group, 21  
 line of curvature, 47, 76  
  
 manifold, 17  
 mean curvature, 47  
 metric, 112  
 metric preserving connexion, 133  
 metric tensor, 43, 109  
 minimal surface, 70, 72, 79, 207  
  
 normal coordinates, 196  
 normal curvature, 119  
 normal deformation, 207  
 normal (of a curve), 116  
  
 one parameter family of curves, 212  
 one parameter subgroup, 35, 143  
 open set, 15  
 orientable, 153  
 orthogonal coordinates, 200  
  
 parallel hypersurface, 60  
 parallel translation, 41, 49, 94  
 parallelizable, 31, 107  
 partition of unity, 130  
 Poincaré space, 127  
 pole, 247  
 principal coordinates, 200  
 principal curvature, 47  
 principal vector, 47  
 pseudo-metric, 111  
  
 regular value, 18  
 Ricci curvature, 146  
 Ricci tensor map, 145  
 Riemann space, 127  
 Riemann surface, 205  
 Riemann-Christoffel curvature, 113, 145  
 Riemannian connexion, 43, 112  
 Riemannian curvature, 115, 226  
 Riemannian manifold, 42  
 Riemannian metric, 43  
 rigidity, 122  
 Rodrigues formula, 47

rotation group, 108  
 ruled surface, 55  
  
 scalar curvature, 146  
 second fundamental form, 46  
 second fundamental tensor, 117  
 second variation formula, 217  
 self-adjoint (symmetric) map, 44  
 semi-Riemannian manifold, 42  
 shuffle permutation, 89  
 simple closed curve, 159  
 skew-symmetric tensor, 82  
 slice, 184  
 special orthogonal group, 108  
 sphere map, 44  
 spherical space, 127  
 Stokes' map, 150  
 structural function, 37  
 subatlas, 16  
 submanifold, 31, 117  
 support, 149, 155  
 surface of revolution, 76  
 symmetric surfaces, 177  
 symmetric tensor, 82  
  
 tangent bundle, 35  
 tangent (of a curve), 27  
  
 tangent space, 22  
 tangent vector, 22  
 tensor algebra, 84  
 tensors, 81  
 torsion (of a curve), 116  
 torsion tensor, 96, 120  
 torus (flat), 34  
 total curvature, 46  
 tractrix, 79  
 two parameter family of curves, 230  
  
 umbilic, 47  
 universal covering manifold, 248  
 universal tensors, 88  
  
 vector field (on submanifold), 31  
 vector fields, 24  
 vertical vector, 37  
 volume, 155  
 volume element, 153  
  
 wedge product, 84  
 Weil homomorphism, 175  
 Weingarten map, 43, 120, 125, 135  
 Weingarten surface, 208

A concise introduction to the topics considered part of a standard background in differential geometry up until today.

The first three chapters provide a crash course in classical differential geometry, leading into an intermezzo of tensors and forms, while the rest of the book introduces the notion of connexion (connection) and various topics in Riemannian geometry. The notes are intended for a full year course, covering the first six chapters in one semester.

Contains a total of 100 problems of varied difficulty, consisting of computations, additional theorems and examples, results from the literature and supplementary topics such as Lie groups and bundle theory.

*Noel J. Hicks* (1929-1979) was born in San Antonio, Texas. His undergraduate studies took place at the University of Wyoming, and his PhD was awarded in 1957 by the Massachusetts Institute of Technology. His thesis, supervised by Warren Ambrose, was titled *On the Curvature and Torsion of Affine Connexions*.

After graduating, Hicks joined the University of Michigan faculty as instructor; he was appointed assistant professor in 1961, associate professor in 1966. Geometry was his field of research and advanced teaching: his textbook on differential geometry is known as a classic treatment of the subject, and he wrote a number of important research papers in the area.

(Sources: U. of Michigan, *Faculty History Project*; *Mathematics Genealogy Project*)