

Annals of Mathematics

On Manifolds Homeomorphic to the 7-Sphere

Author(s): John Milnor

Source: *The Annals of Mathematics*, Second Series, Vol. 64, No. 2 (Sep., 1956), pp. 399-405

Published by: Annals of Mathematics

Stable URL: <http://www.jstor.org/stable/1969983>

Accessed: 23/03/2009 18:33

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=annals>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



Annals of Mathematics is collaborating with JSTOR to digitize, preserve and extend access to *The Annals of Mathematics*.

<http://www.jstor.org>

ON MANIFOLDS HOMEOMORPHIC TO THE 7-SPHERE

BY JOHN MILNOR¹

(Received June 14, 1956)

The object of this note will be to show that the 7-sphere possesses several distinct differentiable structures.

In §1 an invariant λ is constructed for oriented, differentiable 7-manifolds M^7 satisfying the hypothesis (*) $H^3(M^7) = H^4(M^7) = 0$. (Integer coefficients are to be understood.) In §2 a general criterion is given for proving that an n -manifold is homeomorphic to the sphere S^n . Some examples of 7-manifolds are studied in §3 (namely 3-sphere bundles over the 4-sphere). The results of the preceding two sections are used to show that certain of these manifolds are topological 7-spheres, but not differentiable 7-spheres. Several related problems are studied in §4.

All manifolds considered, with or without boundary, are to be differentiable, orientable and compact. The word *differentiable* will mean differentiable of class C^∞ . A closed manifold M^n is *oriented* if one generator $\mu \in H_n(M^n)$ is distinguished.

§1. The invariant $\lambda(M^7)$

For every closed, oriented 7-manifold satisfying (*) we will define a residue class $\lambda(M^7)$ modulo 7. According to Thom [5] every closed 7-manifold M^7 is the boundary of an 8-manifold B^8 . The invariant $\lambda(M^7)$ will be defined as a function of the index τ and the Pontrjagin class p_1 of B^8 .

An orientation $\nu \in H_8(B^8, M^7)$ is determined by the relation $\partial\nu = \mu$. Define a quadratic form over the group $H^4(B^8, M^7)/(\text{torsion})$ by the formula $\alpha \rightarrow \langle \nu, \alpha^2 \rangle$. Let $\tau(B^8)$ be the index of this form (the number of positive terms minus the number of negative terms, when the form is diagonalized over the real numbers).

Let $p_1 \in H^4(B^8)$ be the first Pontrjagin class of the tangent bundle of B^8 . (For the definition of Pontrjagin classes see [2] or [6].) The hypothesis (*) implies that the inclusion homomorphism

$$i: H^4(B^8, M^7) \rightarrow H^4(B^8)$$

is an isomorphism. Therefore we can define a "Pontrjagin number"

$$q(B^8) = \langle \nu, (i^{-1}p_1)^2 \rangle.$$

THEOREM 1. *The residue class of $2q(B^8) - \tau(B^8)$ modulo 7 does not depend on the choice of the manifold B^8 .*

Define $\lambda(M^7)$ as this residue class.² As an immediate consequence we have:

COROLLARY 1. *If $\lambda(M^7) \neq 0$ then M^7 is not the boundary of any 8-manifold having fourth Betti number zero.*

¹ The author holds an Alfred P. Sloan fellowship.

Let B_1^8, B_2^8 be two manifolds with boundary M^7 . (We may assume they are disjoint.) Then $C^8 = B_1^8 \cup B_2^8$ is a closed 8-manifold which possesses a differentiable structure compatible with that of B_1^8 and B_2^8 . Choose that orientation ν for C^8 which is consistent with the orientation ν_1 of B_1^8 (and therefore consistent with $-\nu_2$). Let $q(C^8)$ denote the Pontrjagin number $\langle \nu, p_1^2(C^8) \rangle$.

According to Thom [5] or Hirzebruch [2] we have

$$\tau(C^8) = \langle \nu, \frac{1}{45} (7p_2(C^8) - p_1^2(C^8)) \rangle;$$

and therefore

$$45\tau(C^8) + q(C^8) = 7\langle \nu, p_2(C^8) \rangle \equiv 0 \pmod{7}.$$

This implies

$$(1) \quad 2q(C^8) - \tau(C^8) \equiv 0 \pmod{7}.$$

LEMMA 1. *Under the above conditions we have*

$$(2) \quad \tau(C^8) = \tau(B_1^8) - \tau(B_2^8)$$

and

$$(3) \quad q(C^8) = q(B_1^8) - q(B_2^8).$$

Formulas 1, 2, 3 clearly imply that

$$2q(B_1^8) - \tau(B_1^8) \equiv 2q(B_2^8) - \tau(B_2^8) \pmod{7};$$

which is just the assertion of Theorem 1.

PROOF OF LEMMA 1. Consider the diagram

$$\begin{array}{ccc} H^n(B_1, M) \oplus H^n(B_2, M) & \xleftarrow[\approx]{h} & H^n(C, M) \\ \downarrow i_1 \oplus i_2 & & \downarrow j \\ H^n(B_1) \oplus H^n(B_2) & \xleftarrow{k} & H^n(C) \end{array}$$

Note that for $n = 4$, these homomorphisms are all isomorphisms. If $\alpha = jh^{-1}(\alpha_1 \oplus \alpha_2) \in H^4(C)$, then

$$(4) \quad \langle \nu, \alpha^2 \rangle = \langle \nu, jh^{-1}(\alpha_1^2 \oplus \alpha_2^2) \rangle = \langle \nu_1 \oplus (-\nu_2), \alpha_1^2 \oplus \alpha_2^2 \rangle = \langle \nu_1, \alpha_1^2 \rangle - \langle \nu_2, \alpha_2^2 \rangle.$$

Thus the quadratic form of C^8 is the "direct sum" of the quadratic form of B_1^8 and the negative of the quadratic form of B_2^8 . This clearly implies formula (2).

Define $\alpha_1 = i_1^{-1}p_1(B_1)$ and $\alpha_2 = i_2^{-1}p_1(B_2)$. Then the relation

$$k(p_1(C)) = p_1(B_1) \oplus p_1(B_2)$$

implies that

² Similarly for $n = 4k - 1$ a residue class $\lambda(M^n)$ modulo $s_{k\mu}(L_k)$ could be defined. (See [2] page 14.) For $k = 1, 2, 3, 4$ we have $s_{k\mu}(L_k) = 1, 7, 62, 381$ respectively.

$$jh^{-1}(\alpha_1 \oplus \alpha_2) = p_1(C).$$

The computation (4) now shows that

$$\langle \nu, p_1^2(C) \rangle = \langle \nu_1, \alpha_1^2 \rangle - \langle \nu_2, \alpha_2^2 \rangle,$$

which is just formula (3). This completes the proof of Theorem 1.

The following property of the invariant λ is clear.

LEMMA 2. *If the orientation of M^7 is reversed then $\lambda(M^7)$ is multiplied by -1 .*

As a consequence we have

COROLLARY 2. *If $\lambda(M^7) \neq 0$ then M^7 possesses no orientation reversing diffeomorphism³ onto itself.*

§2. A partial characterization of the n -sphere

Consider the following hypothesis concerning a closed manifold M^n (where R denotes the real numbers).

(H) *There exists a differentiable function $f: M^n \rightarrow R$ having only two critical points x_0, x_1 . Furthermore these critical points are non-degenerate.*

(That is if u_1, \dots, u_n are local coordinates in a neighborhood of x_0 (or x_1) then the matrix $(\partial^2 f / \partial u_i \partial u_j)$ is non-singular at x_0 (or x_1 .)

THEOREM 2. *If M^n satisfies the hypothesis (H) then there exists a homeomorphism of M^n onto S^n which is a diffeomorphism except possibly at a single point.*

Added in proof. This result is essentially due to Reeb [7].

The proof will be based on the orthogonal trajectories of the manifolds $f = \text{constant}$.

Normalize the function f so that $f(x_0) = 0, f(x_1) = 1$. According to Morse ([3] Lemma 4) there exist local coordinates v_1, \dots, v_n in a neighborhood V of x_0 so that $f(x) = v_1^2 + \dots + v_n^2$ for $x \in V$. (Morse assumes that f is of class C^3 , and constructs coordinates of class C^1 ; but the same proof works in the C^∞ case.) The expression $ds^2 = dv_1^2 + \dots + dv_n^2$ defines a Riemannian metric in the neighborhood V . Choose a differentiable Riemannian metric for M^n which coincides with this one in some neighborhood⁴ V' of x_0 . Now the gradient of f can be considered as a contravariant vector field.

Following Morse we consider the differential equation

$$\frac{dx}{dt} = \text{grad } f / \|\text{grad } f\|^2.$$

In the neighborhood V' this equation has solutions

$$(v_1(t), \dots, v_n(t)) = (a_1(t)^{\frac{1}{2}}, \dots, a_n(t)^{\frac{1}{2}})$$

for $0 \leq t < \epsilon$, where $a = (a_1, \dots, a_n)$ is any n -tuple with $\sum a_i^2 = 1$. These can be extended uniquely to solutions $x_a(t)$ for $0 \leq t \leq 1$. Note that these solutions satisfy the identity

³ A diffeomorphism f is a homeomorphism onto, such that both f and f^{-1} are differentiable.

⁴ This is possible by [4] 6.7 and 12.2.

$$f(x_a(t)) = t.$$

Map the interior of the unit sphere of R^n into M^n by the map

$$(a_1(t)^{\frac{1}{2}}, \dots, a_n(t)^{\frac{1}{2}}) \rightarrow x_a(t).$$

It is easily verified that this defines a diffeomorphism of the open n -cell onto $M^n - (x_1)$. The assertion of Theorem 2 now follows.

Given any diffeomorphism $g: S^{n-1} \rightarrow S^{n-1}$, an n -manifold can be obtained as follows.

CONSTRUCTION (C). Let $M^n(g)$ be the manifold obtained from two copies of R^n by matching the subsets $R^n - (0)$ under the diffeomorphism

$$u \rightarrow v = \frac{1}{\|u\|} g\left(\frac{u}{\|u\|}\right).$$

(Such a manifold is clearly homeomorphic to S^n . If g is the identity map then $M^n(g)$ is diffeomorphic to S^n .)

COROLLARY 3. A manifold M^n can be obtained by the construction (C) if and only if it satisfies the hypothesis (H).

PROOF. If $M^n(g)$ is obtained by the construction (C) then the function

$$f(x) = \|u\|^2 / (1 + \|u\|^2) = 1 / (1 + \|v\|^2)$$

will satisfy the hypothesis (H). The converse can be established by a slight modification of the proof of Theorem 2.

§3. Examples of 7-manifolds

Consider 3-sphere bundles over the 4-sphere with the rotation group $SO(4)$ as structural group. The equivalence classes of such bundles are in one-one correspondence⁵ with elements of the group $\pi_3(SO(4)) \approx Z + Z$. A specific isomorphism between these groups is obtained as follows. For each $(h, j) \in Z + Z$ let $f_{h,j}: S^3 \rightarrow SO(4)$ be defined by $f_{h,j}(u) \cdot v = u^h v u^j$, for $v \in R^4$. Quaternion multiplication is understood on the right.

Let ι be the standard generator for $H^4(S^4)$. Let $\xi_{h,j}$ denote the sphere bundle corresponding to $(f_{h,j}) \in \pi_3(SO(4))$.

LEMMA 3. The Pontrjagin class $p_1(\xi_{h,j})$ equals $\pm 2(h - j)\iota$.

(The proof will be given later. One can show that the characteristic class $\bar{c}(\xi_{h,j})$ (see [4]) is equal to $(h + j)\iota$.)

For each odd integer k let M_k^7 be the total space of the bundle $\xi_{h,j}$ where h and j are determined by the equations $h + j = 1$, $h - j = k$. This manifold M_k^7 has a natural differentiable structure and orientation, which will be described later.

LEMMA 4. The invariant $\lambda(M_k^7)$ is the residue class modulo 7 of $k^2 - 1$.

LEMMA 5. The manifold M_k^7 satisfies the hypothesis (H).

Combining these we have:

⁵ See [4] §18.

THEOREM 3. For $k^2 \not\equiv 1 \pmod 7$ the manifold M_k^7 is homeomorphic to S^7 but not diffeomorphic to S^7 .

(For $k = \pm 1$ the manifold M_k^7 is diffeomorphic to S^7 ; but it is not known whether this is true for any other k .)

Clearly any differentiable structure on S^7 can be extended through $R^8 - (0)$. However:

COROLLARY 4. There exists a differentiable structure on S^7 which cannot be extended throughout R^8 .

This follows immediately from the preceding assertions, together with Corollary 1.

PROOF OF LEMMA 3. It is clear that the Pontrjagin class $p_1(\xi_{hj})$ is a linear function of h and j . Furthermore it is known that it is independent of the orientation of the fibre. But if the orientation of S^3 is reversed, then ξ_{hj} is replaced by ξ_{-j-h} . This shows that $p_1(\xi_{hj})$ is given by an expression of the form $c(h - j)\iota$. Here c is a constant which will be evaluated later.

PROOF OF LEMMA 4. Associated with each 3-sphere bundle $M_k^7 \rightarrow S^4$ there is a 4-cell bundle $\rho_k: B_k^8 \rightarrow S^4$. The total space B_k^8 of this bundle is a differentiable manifold with boundary M_k^7 . The cohomology group $H^4(B_k^8)$ is generated by the element $\alpha = \rho_k^*(\iota)$. Choose orientations μ, ν for M_k^7 and B_k^8 so that

$$\langle \nu, (i^{-1}\alpha)^2 \rangle = +1.$$

Then the index $\tau(B_k^8)$ will be $+1$.

The tangent bundle of B_k^8 is the "Whitney sum" of (1) the bundle of vectors tangent to the fibre, and (2) the bundle of vectors normal to the fibre. The first bundle (1) is induced (under ρ_k) from the bundle ξ_{hj} , and therefore has Pontrjagin class $p_1 = \rho_k^*(c(h - j)\iota) = ck\alpha$. The second is induced from the tangent bundle of S^4 , and therefore has first Pontrjagin class zero. Now by the Whitney product theorem ([2] or [6])

$$p_1(B_k^8) = ck\alpha + 0.$$

For the special case $k = 1$ it is easily verified that B_1^8 is the quaternion projective plane $P_2(K)$ with an 8-cell removed. But the Pontrjagin class $p_1(P_2(K))$ is known to be twice a generator of $H^4(P_2(K))$. (See Hirzebruch [1].) Therefore the constant c must be ± 2 , which completes the proof of Lemma 3.

Now $q(B_k^8) = \langle \nu, (i^{-1}(\pm 2k\alpha))^2 \rangle = 4k^2$; and $2q - \tau = 8k^2 - 1 \equiv k^2 - 1 \pmod 7$. This completes the proof of Lemma 4.

PROOF OF LEMMA 5. As coordinate neighborhoods in the base space S^4 take the complement of the north pole, and the complement of the south pole. These can be identified with euclidean space R^4 under stereographic projection. Then a point which corresponds to $u \in R^4$ under one projection will correspond to $u' = u/\|u\|^2$ under the other.

The total space M_k^7 can now be obtained as follows.⁵ Take two copies of $R^4 \times S^3$ and identify the subsets $(R^4 - (0)) \times S^3$ under the diffeomorphism

$$(u, v) \rightarrow (u', v') = (u/\|u\|^2, u^h v u^j / \|u\|)$$

(using quaternion multiplication). This makes the differentiable structure of M_k^7 precise.

Replace the coordinates (u', v') by (u'', v') where $u'' = u'(v')^{-1}$. Consider the function $f: M_k^7 \rightarrow R$ defined by

$$f(x) = \Re(v)/(1 + \|u\|^2)^{\frac{1}{2}} = \Re(u'')/(1 + \|u''\|^2)^{\frac{1}{2}};$$

where $\Re(v)$ denotes the real part of the quaternion v . It is easily verified that f has only two critical points (namely $(u, v) = (0, \pm 1)$) and that these are non-degenerate. This completes the proof.

§4. Miscellaneous results

THEOREM 4. *Either (a) there exists a closed topological 8-manifold which does not possess any differentiable structure; or (b) the Pontrjagin class p_1 of an open 8-manifold is not a topological invariant.*

(The author has no idea which alternative holds.)

PROOF. Let X_k^8 be the topological 8-manifold obtained from B_k^8 by collapsing its boundary (a topological 7-sphere) to a point x_0 . Let $\bar{\alpha} \in H^4(X_k^8)$ correspond to the generator $\alpha \in H^4(B_k^8)$. Suppose that X_k^8 , possesses a differentiable structure, and that $p_1(X_k^8 - (x_0))$ is a topological invariant. Then $p_1(X_k^8)$ must equal $\pm 2k\bar{\alpha}$, hence

$$2q(X_k^8) - \tau(X_k^8) = 8k^2 - 1 \equiv k^2 - 1 \pmod{7}.$$

But for $k^2 \not\equiv 1 \pmod{7}$ this is impossible.

Two diffeomorphisms $f, g: M_1^n \rightarrow M_2^n$ will be called *diffeomorphically isotopic* if there exists a diffeomorphism $M_1^n \times R \rightarrow M_2^n \times R$ of the form $(x, t) \rightarrow (h(x, t), t)$ such that

$$h(x, t) = \begin{cases} f(x) & (t \leq 0) \\ g(x) & (t \geq 1). \end{cases}$$

LEMMA 6. *If the diffeomorphisms $f, g: S^{n-1} \rightarrow S^{n-1}$ are diffeomorphically isotopic, then the manifolds $M^n(f), M^n(g)$ obtained by the construction (C) are diffeomorphic.*

The proof is straightforward.

THEOREM 5. *There exists a diffeomorphism $f: S^6 \rightarrow S^6$ of degree +1 which is not diffeomorphically isotopic to the identity.*

Proof. By Lemma 5 and Corollary 3 the manifold M_3^7 is diffeomorphic to $M^7(f)$ for some f . If f were diffeomorphically isotopic to the identity then Lemma 6 would imply that M_3^7 was diffeomorphic to S^7 . But this is false by Lemma 4.

PRINCETON UNIVERSITY

REFERENCES

1. F. HIRZEBRUCH, *Ueber die quaternionalen projektiven Räume*, S.-Ber. math.- naturw. Kl. Bayer. Akad. Wiss. München (1953), pp. 301-312.
2. ———, *Neue topologische Methoden in der algebraischen Geometrie*, Berlin, 1956.

3. M. MORSE, *Relations between the numbers of critical points of a real function of n independent variables*, Trans. Amer. Math. Soc., 27 (1925), pp. 345-396.
4. N. STEENROD, *The topology of fibre bundles*, Princeton, 1951.
5. R. THOM, *Quelques propriétés globale des variétés différentiables*, Comment. Math. Helv., 28 (1954), pp. 17-86.
6. WU WEN-TSUN, *Sur les classes caractéristiques des structures fibrées sphériques*, Actual. sci. industr. 1183, Paris, 1952, pp. 5-89.
7. G. REEB, *Sur certain propriétés topologiques des variétés feuilletées*, Actual. sci. industr. 1183, Paris, 1952, pp. 91-154.