

Math 531 Tom Tucker  
NOTES FROM CLASS 11/12

Throughout,  $L$  is as usual degree  $n$  over  $\mathbb{Q}$ ,  $h : L \rightarrow V$  is the usual embedding,  $r$  is the number of real places of  $L$  and  $s = (n - r)/2$ . Also,  $N$  is  $N_{L/\mathbb{Q}}$ .

**Proposition 30.1.**

$$\text{Vol}(X_t) = \frac{2^{r-s} \pi^s t^n}{n!}.$$

*Proof.* The proof of this is in the book on p. 66. The last step in the calculation is integration by parts, which the book neglects to mention.  $\square$

**Lemma 30.2.** *Let  $U$  be any bounded region of  $V$  and let  $\mathcal{L}$  be a full lattice in  $V$ . Then  $\mathcal{L} \cap U$  is finite.*

*Proof.* Let  $w_1, \dots, w_n$  be a basis for  $\mathcal{L}$  and let  $x_1, \dots, x_n$  be the basis for  $V$  that gives the volume form. If  $M$  is the matrix given by  $Mx_i = w_i$ , then for any integers  $m_i$  we have

$$\left| \sum_{i=1}^n m_i w_i \right|^2 = \left| M \left( \sum_{i=1}^n m_i x_i \right) \right|^2 \geq \sum_{i=1}^n m_i^2 \|M\|_{\inf}^2$$

where  $\|M\|_{\inf}$  is the minimum value of  $|M(y)|$  for  $y$  on the unit sphere centered at the origin (which is nonzero). For any constant  $C$  there are finitely many integers  $m_i$  such that

$$\sum_{i=1}^n m_i^2 \|M\|_{\inf}^2 \leq C^2$$

so there are finitely many elements of  $\lambda$  in the sphere of radius  $C$  centered at the origin. Any bounded region is contained in such a sphere, so we are done.  $\square$

**Theorem 30.3.** *Let  $I$  be a nonzero fractional ideal of  $\mathcal{O}_L$ . Then there exists a  $\neq 0$  such that*

$$|N_{L/\mathbb{Q}}(a)| \leq \frac{n!}{n^n} \left( \frac{4}{\pi} \right)^s \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N_{L/\mathbb{Q}}(I).$$

*Proof.* We want to choose  $X_t$  to which we can apply Minkowski's theorem and produce an element of  $X_t \cap h(I)$ . Recall that

$$\text{Vol}(h(I)) = \frac{1}{2^s} \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I),$$

so we need  $t$  with

$$\frac{2^{r-s} \pi^s t^n}{n!} > 2^n \frac{1}{2^s} \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I),$$

which is equivalent to

$$t > \sqrt[n]{n! \frac{1}{\pi^s} 2^{n-s-r+s} \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I)} = \sqrt[n]{n! \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I)},$$

so let

$$C := \sqrt[n]{n! \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I)}.$$

Then  $\text{Vol}(X_{C+\epsilon}) > \text{Vol}(h(I))$  for any  $\epsilon > 0$ . It follows that  $X_{C+\epsilon} \cap h(I) \neq \emptyset$  by Minkowski's theorem. If

$$X_{C+\epsilon} \cap h(I) = X_C \cap h(I),$$

then  $X_C \cap h(I) \neq \emptyset$ . Otherwise, let  $\epsilon' > 0$  be the smallest number such that

$$X_{C+\epsilon} \cap h(I) \neq X_C \cap h(I).$$

Such a number exists since  $X_{C+\epsilon} \cap h(I)$  is finite and any finite nonempty set has a minimal element. Taking  $0 < \delta < \epsilon'$ , we see that

$$X_C \cap h(I) = X_{C+\delta} \cap h(I) \neq \emptyset,$$

so there is a nonzero element  $a \in X_C \cap h(I)$ . From earlier work, we see that

$$N(a) \leq (C/n)^n = \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L/\mathbb{Z})} N(I).$$

□

Let's do an easy lemma.

**Lemma 30.4.** *Let  $I$  be a fractional ideal of a Dedekind domain  $A$  and let  $a \neq 0$  be in  $I$ . Then  $aI^{-1} \subseteq A$ .*

*Proof.* Since  $Aa \subseteq I$ , we have

$$I^{-1}Aa \subseteq II^{-1} = A.$$

□

**Theorem 30.5.** *Let  $I \subset \mathcal{O}_L$  be any fractional ideal of  $\mathcal{O}_L$ . Then there exists an ideal  $J \subset \mathcal{O}_L$  in the same ideal class as  $I$  such that*

$$|N_{L/\mathbb{Q}}(J)| \leq \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s N_{L/\mathbb{Q}}(I) \sqrt{\Delta(\mathcal{O}_L/bZ)}.$$

*Proof.* Applying the previous theorem to  $I^{-1}$ , we find that there is an element  $a \in I^{-1}$  such that

$$|N_{L/\mathbb{Q}}(a)| \leq \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L/bZ)} N(I)^{-1}.$$

Let  $J = aI$ . Since  $a \in I^{-1}$ , we see that

$$aI = a(I^{-1})^{-1} \subset \mathcal{O}_L.$$

We also have

$$N(aI) \leq \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L//bZ)} N(I)^{-1} N(I) = \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{\Delta(\mathcal{O}_L//bZ)},$$

so we are done.  $\square$

My computation for  $\mathbb{Z}[\sqrt{-13}]$  was incorrect, I'll fix it next time.