

Math 531 Problem Set #2 Due 9/15/04

1. (a) Suppose that α is integral of degree 2 over \mathbb{Z} and that there is a basis m_1, m_2 for $\mathbb{Z}[\alpha]$ over \mathbb{Z} such that

$$\begin{aligned}\alpha m_1 &= m_2 \\ \alpha m_2 &= -m_1 + m_2.\end{aligned}$$

Find the quadratic integral equation satisfied by α .

- (b) Suppose that β is integral of degree 2 over \mathbb{Z} and that there is a basis m_1, m_2 for $\mathbb{Z}[\beta]$ over \mathbb{Z} such that

$$\begin{aligned}\beta m_1 &= m_2 \\ \beta m_2 &= 5m_1.\end{aligned}$$

Find the quadratic integral equation satisfied by β .

2. Let T be an $n \times n$ matrix with coefficients in a ring A . Show that there is a matrix U for which $UT = (\det T)I$.

3. Let A , B , and C be rings with $A \subset B \subset C$. Show that if B is integral over A and C is integral over B , then C is integral over A .

4. (Ex. 4, p. 7) Let d be a squarefree integer. Show that the integral closure of \mathbb{Z} in $\mathbb{Q}[\sqrt{d}]$ is

$$\mathbb{Z}[\sqrt{d}] \quad \text{if } d \equiv 2, 3 \pmod{4},$$

and

$$\mathbb{Z}\left[\frac{1 + \sqrt{d}}{2}\right] \quad \text{if } d \equiv 1 \pmod{4}.$$

5. (a) Let $\phi : A \rightarrow B$ be a mapping of rings. Show that for any prime ideal \mathcal{P} in B , the ideal $\phi^{-1}(\mathcal{P})$ is a prime ideal in A .

- (b) Give an example of a surjective ring homomorphism $\phi : A \rightarrow B$ for which there is a prime ideal \mathcal{P} of A such that $\phi(\mathcal{P})$ is *not* a prime ideal.

6. (a) Give an example of a mapping of rings $\phi : A \rightarrow B$ for which there is an ideal I of A such that $\phi(I)$ is not an ideal.

- (b) Let $\phi : A \rightarrow B$ be a surjective mapping of rings. Show that for any ideal I of A , the set $\phi(I)$ forms an ideal in B .

- (c) Let $\phi : A \rightarrow B$ be any mapping of rings. Show that for any ideal J of B , the set $\phi^{-1}(J)$ forms an ideal in A .

7. Let $A \subset B$ where A and B are domains and let K be the field of fractions of B . Show that if B is integrally closed over A in K , then B is integrally closed over itself in K .

8. (Ex. 4, p.3) Show that if S is a multiplicative set not containing 0 in a Noetherian integral domain R , then $S^{-1}R$ is also a Noetherian integral domain.